

QSO (AGN) study with PFS

Masa Imanishi

NAOJ/Subaru Telescope



With help from

Masayuki Akiyama (Tohoku)

Yoichi Itoh (Kobe)



Toshihiro Kawaguchi (Tsukuba)

Tohru Nagao (Ehime)

science discussion inside HSC QSO team

QSO (AGN)

Type-1



Type-1 QSO = bright




High S/N spectra
even at high-z

AGN-starburst connections
Feedback to host galaxies
Origin of galaxy downsizing?

Energetically dominated
by mass-accreting
Supermassive blackhole
(SMBH) ($>10^7 M_{\text{sun}}$)

QSO science with PFS = follow-up spectroscopy of HSC-detected QSO candidates

HSC wide survey : $>1000 \text{ deg}^2$ (g,r,i,z,y)




**g=26.3, r=26.0,
i=26.0, z=24.9,
y=23.8 (AB)**

**1. numerous faint
QSOs at $z>3$**

2. $z>6.7$ QSOs

$z > 3$ QSOs



Richards+06

**QSO number density
has a peak at $z=2-3$**



**$z > 3$ = SMBH growing
phase**

**Important epoch to
understand the
cosmic evolution
of QSO phenomenon**

**SDSS found many QSOs only at $z>3$
($i<20.1$ for spectroscopy)**

What is unknown about $z>3$ QSOs ?

Faint QSO LF at $z>3$

Clustering properties of QSOs at $z>3$

Evolution of general SMBHs ($10^8 M_{\text{sun}}$)


Metallicity in chemically immaturated QSOs

$z>6.4$ QSOs




HSC survey will bring a remarkable progress

HSC $z \sim 3-5$ QSO (g,r,i-selection)




gri : $z=3.6-4.4$

riz : $z=4.6-5.1$




QSO LF SDSS : bright QSO LF only at $z > 3$
($i < 20.1$ for spectroscopy)



**Understanding the faint end of QSO LF
= origin of cosmic UV background radiation**

↳ **Effect to subsequent galaxy formation**


QSO-QSO clustering



Croom+05 (>20,000 QSOs)

$z > 3$

Halo mass
QSO lifetime



SMBH evolution

LF





Constant $L/L(\text{Edd})$
assumed

SMBH evolution
(qualitative)

$M(\text{SMBH})$ and
 $L/L(\text{Edd})$ in
individual QSOs



How about more
general QSOs with
 $M(\text{SMBH}) < 10^8 M_\odot$?

HSC




Kelly+10 (SDSS)

Cosmic chemical evolution



**Star-forming galaxies:
not applicable to $z > 3.5$**




QSOs

Cosmic chemical evolution (2)

QSOs are bright

QSOs have metal emission lines in UV




Chemical evolution can be probed at $z > 3.5$ via optical spectroscopy

Modest luminosity QSOs = chemically-immatured ?

Assumption about PFS

- **1.5 deg² FOV**
- **3900-10800 Å (extended to 13000Å ?)**
- **2400 fibers**
- **R=3000**


Sensitivity (by Dr. Iwata) (900sec, R=2500)



Number density of $z=3-5.5$ QSOs


30-650 QSOs / deg^2

$i < 24$



Candidates = 50-2000 / deg^2

Casey+08



Survey strategy ($z=3-5.5$ QSOs)

LF, clustering

ID as $z>3$ QSOs ($i<24$, $S/N>3-5$, $R\sim 300-500$)

1 hr, 75 deg², **<10 nights** (LF)

1 hr, >400 deg², **>40 nights** (clustering)

SMBH evolution

10^8 Mo at $z=4$


$S/N>10$ to estimate $M(\text{SMBH})$ $i<22.5$

1 hr, 400 deg², **~ 35 nights**



Chemical evolution

Weak metal lines $i<23$

2.5 hr, 150 deg², **~ 35 nights**



$z > 6$ QSOs in HSC wide survey



350 $z \sim 6$ QSOs
($z < 24.2$)

35 $z \sim 7$ QSOs
($y < 23.8$)

$z \sim 7$ QSO science

Currently, most distant QSO : $z \sim 6.4$

Most distant, bright QSOs

Constraints on SMBH formation theory

$z=25$: 100 Mo BH Eddington growth



$z=6.5$: 10^9 Mo $z=7.5$: 10^8 Mo


Study the physics (including neutral fraction) of more distant IGM with high S/N

High-z QSO and brown dwarf (BD)

**Galactic
BD**

flux

**Gradually
increase**




wavelength

**High-z
QSO**

flux

Sharp gap




wavelength

$z \sim 7$ QSO (z-dropout, y-detected)

$z-y > 2.3$ selection (remove BD)


$z=24.9, y=22.6$



**In VIKING field (J=22.2)AB,
 $z-y > 1.7, y-J < 1$**

$z=24.9, y=23.2$

(HSC: $z=24.9, y=23.8$)




y-J
Kashikawa10

$z \sim 6$ QSO science

Many faint $z \sim 6$ QSO

**$z \sim 6$ IGM study in
many sightlines
(neutral fraction
increases?)**





$z \sim 6$ QSO (i-dropout, z-detected)

i-z red (>1.8) : by IGM absorption


z-y blue (<0.5) : BD removal

$i=26.0, z=24.2, y=23.8$



z~7 QSOs (z-dropout,y-detected)

**35 z~7 QSOs should be included
in HSC wide 1000 deg² ($y < 23.8$)**



**We can select only 10 z~7 QSOs (from BDs)
($y < 23.2$) via Subaru FOCAS spectroscopy**



How about remaining 25 z~7 QSOs ?

**PFS spectroscopy of z-dropout sources
($z > 24.9$, $y = 23.2 - 23.8$)**

2.5hr, > 500 deg², > 100 nights

< 10 candidates / PFS FOV

z~6 QSOs (i-dropout, z-detected)


**Impossible to spectroscopically
identify 350 z~6 QSOs ($z < 24.2$)
with Subaru FOCAS**



**PFS spectroscopy of i-dropout sources
($i > 26.0$, $z < 24.9$)**

2.5hr, $>500\text{deg}^2$, >100 nights

<10 candidates / PFS FOV



Brown dwarf (BD) science


PFS spectroscopy ($z < 22.6$) of HSC-wide area will find 500 BDs



Stellar IMF in
BD regime

2.5hr, $>500 \text{ deg}^2$,
 >100 nights

<10 candidates / PFS FOV



Chabrier+03

Summary

1hr, 400 deg², 40 nights


**LF, clustering, SMBH evolution
of QSOs at $z=3-5.5$**

50-1000 / PFS FOV

2.5hr, >500 deg² (?), >100 nights

Chemical evolution, $z>6$ QSOs, BD

<500 / PFS FOV



**Combined with 4hr, 300 deg²
galaxy survey ?**

Request for PFS specification and survey design

High sensitivity at $>8000\text{\AA}$

Wide wavelength coverage (3900-10800 $\text{\AA}$)

A wider FOV ($>1.5 \text{ deg}^2$) may help

2400 fibers and $R=3000$ are sufficient

Target field must be HSC wide

HSC+PFS is most powerful until LSST (and subsequent follow-up spectroscopy) start operation

TMT

Spectroscopy of $z > 6$ QSOs to study IGM

ALMA


gas of QSO host galaxies

WFIRST/WISH

Numerous fainter $z > 7$ QSOs

Other potential QSO science with PFS

- Obscured AGNs (via galaxy survey)



← Type-2

- $z < 3$ QSOs (u-band available ?)
- Ly alpha forest toward $z < 3$ QSOs
- variability-selected low-L AGN (deep)





Other people's talk

If $z > 3$ QSO number per PFS FOV is $\ll 2400$,
 $z < 3$ PFS QSO science should be combined

End

z~7 QSO

**Number of faint QSOs :
Smaller than previously expected**



Selectable z~7 QSO:
7/1000 deg²
12 (VIKING)
(conservative)

Lim-mag:
(z,y) = (24.9, 23.8) AB


z~7 QSOs included
in y-band : ~35/1000deg²

Selection z-y>1.5

PFS (z~7 QSO)

Follow-up spectroscopy of HSC z~7 QSO candidates

Several 10s candidates/1000 deg²



Use extra several fibers in other galaxy surveys

At least cover Ly α : 9000 - 10500Å

High sensitivity required



SMBH mass  MOIRCS

IGM physics  TMT

2. $z \sim 6$ QSO (i-dropout, z-detected)

i-z red (>1.8) : by IGM absorption

z-y blue (<0.5) : BD removal




350/1000 deg²
(conservative)

Selection :
i=26.0, z=24.2, y=23.8
(AB)

PFS (z~6 QSO)

Follow-up spectroscopy of HSC z~6 QSO candidates

1000 candidates/1000 deg² (1/deg²)



Use extra several fibers
in other galaxy survey

At least cover Ly α : 8000 – 9000Å

High sensitivity required

z~6 IGM multiple sightlines  TMT