

FERMIONIC

T - DUALITY

Nathan Berkovits
(IFT-UNESP, São Paulo)

NB and Juan Maldacena , arXiv: 0807.3196
"Fermionic T-duality, dual superconformal symmetry,
and the amplitude/Wilson loop connection"

Alday and Maldacena used invariance of $AdS_5 \times S^5$ metric under T-duality to relate planar $N=4$ $d=4$ sYM amplitudes and Wilson line correlation functions at strong 'tHooft coupling.

$$\frac{1}{y^2} (dx^m dx_m + dy dy) \rightarrow y^2 d\tilde{x}^m d\tilde{x}_m + \frac{dy dy}{y^2} = \frac{1}{(y')^2} (d\tilde{x}^m d\tilde{x}_m + dy' dy')$$

$y' = \frac{1}{y}$

But full $AdS_5 \times S^5$ background was not invariant since T-duality mapped $F_{01234}^{RR} \rightarrow F_q^{RR}$ and mapped $\Phi \rightarrow \Phi + 4 \log y$.

So amplitude/Wilson line relation was only proven in limit $r_{AdS} \rightarrow \infty$ where effects of F^{RR} and Φ can be ignored.

Drummond, Henn, Korchemsky, Smirnov, Sokatchev used perturbative computations to show that planar $N=4$ $d=4$ sYM amplitudes and Wilson line correlation functions are also related at weak 'tHooft coupling.

Question: Can Alday-Maldacena result be extended to planar amplitudes at arbitrary 'tHooft coupling?

By including bosonic and fermionic T-dualities, invariance of AdS_5 metric can be extended to invariance of full $AdS_5 \times S^5$ background.

When superstring background has "abelian" fermionic isometry (i.e. $q^2 = 0$), usual bosonic T-duality has straightforward generalization to fermionic T-duality.

Fermionic T-duality leaves g_{mn} and b_{mn} invariant but transforms F^{RR} and Ψ .

After performing bosonic T-duality on four x^m variables and performing fermionic T-duality on eight $\Theta^{\alpha j}$ variables, the $AdS_5 \times S^5$ background is invariant.

Ignoring the issue of polarization dependence, this explains the relation of planar SYM amplitudes and Wilson lines at arbitrary 'tHooft coupling.

Fermionic T-duality has other applications such as mapping superstring in flat background to superstring in self-dual graviphoton background.

Review of Bosonic T-Duality (ala Buscher)

Suppose string theory background is invariant under the constant shift
 $x' \rightarrow x' + c \Rightarrow x'$ only appears with derivatives in sigma model.

$$S = \int d^2z (g_{mn} + b_{mn}) \partial x^m \bar{\partial} x^n$$

$$l_{mn} = g_{mn} + b_{mn}$$

$$= \int d^2z [g_{11} \partial x^1 \bar{\partial} x^1 + l_{1m} \partial x^1 \bar{\partial} x^m + l_{m1} \partial x^m \bar{\partial} x^1 + l_{mn} \partial x^m \bar{\partial} x^n] \quad m, n \neq 1$$

↑ Integrate out \tilde{x}' . Use that $\partial \bar{A} - \bar{\partial} A = 0 \Rightarrow A = \partial x'$ and $\bar{A} = \bar{\partial} x'$ for some x' .

$$= \int d^2z [g_{11} A \bar{A} + l_{1m} A \bar{\partial} x^m + l_{m1} \partial x^m \bar{A} + l_{mn} \partial x^m \bar{\partial} x^n + \tilde{x}' (\partial \bar{A} - \bar{\partial} A)]$$

↓ Integrate out A and \bar{A} .

$$= \int d^2z \left[\frac{1}{g_{11}} \partial \tilde{x}' \bar{\partial} \tilde{x}' + \frac{l_{1m}}{g_{11}} \partial \tilde{x}' \bar{\partial} x^m - \frac{l_{m1}}{g_{11}} \partial x^m \bar{A} + \left(l_{mn} - \frac{l_{1m} l_{m1}}{g_{11}} \right) \partial x^m \bar{\partial} x^n \right]$$

Comment: In Buscher procedure using non-compact x^i variable,
 $\bar{\partial}A - \partial\bar{A} = 0$ implies $A = \partial x^i$ and $\bar{A} = \bar{\partial}x^i$ only if
 $\int_C (A dz + \bar{A} d\bar{z}) = 0$ around all non-trivial cycles C .

This is not an issue for genus zero amplitudes which have no non-trivial cycles.

But on higher genus surfaces, T-duality is only a symmetry if x^i is a compact variable.

When x^i is compact, $\bar{\partial}A - \partial\bar{A} = 0$ implies that $A = \partial x^i$ and $\bar{A} = \bar{\partial}x^i$ where $\int_C (A dz + \bar{A} d\bar{z})$ is the winding number of x^i around the cycle C .

After T-dualizing $x^i \rightarrow \tilde{x}^i$,

$$S = \int d^2z \left[\frac{1}{g_{11}} \partial \tilde{x}^i \bar{\partial} \tilde{x}^i + \frac{l'_{im}}{g_{11}} \partial \tilde{x}^i \bar{\partial} x^m - \frac{l'_{mi}}{g_{11}} \partial x^m \bar{\partial} \tilde{x}^i + (l_{mn} - \frac{l_{in} l_{mi}}{g_{11}}) \partial x^m \bar{\partial} x^n \right]$$

implies that background fields $l_{mn} = g_{mn} + b_{mn}$ transform as

$$g'_{11} = \frac{1}{g_{11}}, \quad l'_{im} = \frac{l'_{im}}{g_{11}}, \quad l'_{mi} = -\frac{l'_{mi}}{g_{11}}, \quad l'_{mn} = l_{mn} - \frac{l_{in} l_{mi}}{g_{11}}$$

Integration over A and \bar{A} produces a Jacobian factor $(\det g_{11})^{-1}$
which is absorbed by transforming the dilaton as

$$\varphi' = \varphi - \frac{1}{2} \log g_{11}.$$

- sign in l'_{mi} implies that Dirichlet/Neumann boundary conditions for x^i variable are switched on a D-brane.

Bosonic T-duality changes dimension of D-brane.

Suppose background superfields are invariant under constant fermionic shift $\Theta' \rightarrow \Theta' + \xi \Rightarrow \Theta'$ only appears with derivatives in action.

Invariance implies background has "abelian" susy q satisfying $q^2 = 0$.

$$S = \int d^2z (G_{MN} + B_{MN}) \partial Y^M \bar{\partial} Y^N \quad L_{MN} = G_{MN} + B_{MN}$$

$$= \int d^2z [B_{11} \partial \Theta' \bar{\partial} \Theta' + L_{IM} \partial \Theta' \bar{\partial} Y^M + L_{MI} \partial Y^M \bar{\partial} \Theta' + L_{MN} \partial Y^M \bar{\partial} Y^N]$$

↑ Integrate out $\tilde{\Theta}'$. Use that $\partial \bar{a} - \bar{\partial} a = 0 \Rightarrow a = \partial \Theta'$ and $\bar{a} = \bar{\partial} \Theta'$ for some Θ' .

$$= \int d^2z [B_{11} a \bar{a} + L_{IM} a \bar{\partial} Y^M + L_{MI} \partial Y^M \bar{a} + L_{MN} \partial Y^M \bar{\partial} Y^N + \tilde{\Theta}' (\partial \bar{a} - \bar{\partial} a)]$$

↓ Integrate out a and \bar{a} .

$$= \int d^2z \left[-\frac{1}{B_{11}} \partial \tilde{\Theta}' \bar{\partial} \tilde{\Theta}' + \frac{L_{IM}}{B_{11}} \partial \tilde{\Theta}' \bar{\partial} Y^M + \frac{L_{MI}}{B_{11}} \partial Y^M \bar{\partial} \tilde{\Theta}' + \left(L_{MN} - \frac{L_{IM} L_{MI}}{B_{11}} \right) \partial Y^M \bar{\partial} Y^N \right]$$

Comment: In spacetime supersymmetric sigma models, θ' is single-valued around any non-trivial cycle.
(like a non-compact x' variable)

So $\partial\bar{a} - \bar{\partial}a = 0$ implies $a = \partial\theta'$ and $\bar{a} = \bar{\partial}\theta'$

only if $\int_c (a dz + \bar{a} d\bar{z}) = 0$ around every non-trivial cycle C .

This is not an issue for genus zero amplitudes which have no non-trivial cycles.

But on higher genus surfaces, fermionic T-duality is not a symmetry. To make it a symmetry, would need to introduce multivalued fermionic variable θ' satisfying $\theta' \rightarrow \theta' + \xi_c$ when θ' goes around cycle C .

ξ_c is a fermionic zero mode which would need to be included in the functional integral  for $\int D\theta'$.

After T-dualizing $\theta' \rightarrow \tilde{\theta}'$,

$$S = \int d^2z \left[-\frac{1}{B_{11}} \partial \tilde{\theta}' \bar{\partial} \tilde{\theta}' + \frac{L_{1M}}{B_{11}} \partial \tilde{\theta}' \bar{\partial} y^M + \frac{L_{M1}}{B_{11}} \partial y^M \bar{\partial} \tilde{\theta}' + \left(L_{MN} - \frac{L_{1N}L_{M1}}{B_{11}} \right) \partial y^M \bar{\partial} y^N \right]$$

implies that background superfields $L_{MN} = G_{MN} + B_{MN}$ transform as

$$B_{11}' = -\frac{1}{B_{11}}, \quad L_{1M}' = \frac{L_{1M}}{B_{11}}, \quad L_{M1}' = +\frac{L_{M1}}{B_{11}}, \quad L_{MN}' = L_{MN} - \frac{L_{1N}L_{M1}}{B_{11}}$$

Integration over fermionic Q and \bar{Q} produces a Jacobian factor $(\det B_{11})^{+1}$ which is absorbed by transforming the dilaton as

$$\varphi' = \varphi + \frac{1}{2} \log B_{11}$$

+ sign in L_{M1}' implies that Dirichlet/Neumann boundary conditions for θ' variable are not switched on a D-brane.

Fermionic T-duality does not change dimension of D-brane.

To derive T-duality transformations of component supergravity fields, it is very convenient to use pure spinor formalism as was done by Benichou, Policastro, Troost for bosonic T-duality.

Defining $C = B_{\alpha\beta}|_{\Theta=0}$, one finds that after T-dualization of Θ' ,

$$g'_{mn} = g_{mn}, \quad b'_{mn} = b_{mn}, \quad \varphi' = \varphi + \frac{1}{2} \log C, \quad e^{\varphi'} F'^{\alpha\hat{\beta}} = e^{\varphi} F^{\alpha\hat{\beta}} - \epsilon^\alpha \epsilon^{\hat{\beta}} C^{-1}$$

$\alpha, \hat{\beta} = 1 \text{ to } 16$ are $N=2 d=10$ spinor indices

$$F'^{\alpha\hat{\beta}} = \gamma_m^{\alpha\hat{\beta}} F^m + \frac{1}{3!} \gamma_{mnp}^{\alpha\hat{\beta}} F^{mnp} + \frac{1}{2 \cdot 5!} \gamma_{mnpqr}^{\alpha\hat{\beta}} F^{mnpqr}$$

$(\epsilon^\alpha, \epsilon^{\hat{\beta}})$ is Killing spinor associated with abelian susy

$$\text{Abelian susy} \Rightarrow \epsilon^\alpha (\gamma_m)_{\alpha p} \epsilon^p + \epsilon^{\hat{\beta}} (\gamma_m)_{\hat{\alpha}\hat{p}} \epsilon^{\hat{p}} = \cancel{\cancel{\cancel{O}}} \quad O$$

$$\text{Torsion constraints} \Rightarrow \epsilon^\alpha (\gamma_m)_{\alpha p} \epsilon^p - \epsilon^{\hat{\beta}} (\gamma_m)_{\hat{\alpha}\hat{p}} \epsilon^{\hat{p}} = \partial_m C$$

Using $\partial_m C = \epsilon^\alpha (\gamma_m)_\alpha{}^\beta \epsilon^\beta - \epsilon^{\hat{\alpha}} (\gamma_m)_{\hat{\alpha}}{}^{\hat{\beta}} \epsilon^{\hat{\beta}}$, can easily check that transformed background fields

$$g'_{mn} = g_{mn}, b'_{mn} = b_{mn}, \varphi' = \varphi + \frac{1}{2} \log C, e^{\alpha} F'^{\alpha\hat{\beta}} = e^{\alpha} F^{\alpha\hat{\beta}} - \epsilon^{\alpha} \epsilon^{\hat{\beta}} C^{-1}$$

satisfy supergravity equations of motion and are invariant under abelian susy described by Killing spinor $\epsilon'^\alpha = \frac{\epsilon^\alpha}{C}$ and $\epsilon'^{\hat{\alpha}} = \frac{\epsilon^{\hat{\alpha}}}{C}$.

Note that constant mode of C is unconstrained since when B_{11} is constant,

$$\int d^2 z B_{11} \partial \theta' \bar{\partial} \theta' = \int d^2 z \frac{1}{2} B_{11} [\partial(\theta' \bar{\partial} \theta') - \bar{\partial}(\theta' \partial \theta')] = 0$$

Assumes that surface terms can be ignored.

Example 1: d=4 Minkowski + Calabi-Yau 3-fold

Using d=4 hybrid formalism, sigma model in this Type II background is

$$S = \int d^2z [\partial x_{a\dot{a}} \bar{\partial} x^{a\dot{a}} + p_a \bar{\partial} \theta^a + \hat{p}_a \partial \hat{\theta}^a + p_{\dot{a}} \bar{\partial} \theta^{\dot{a}} + \hat{p}_{\dot{a}} \partial \hat{\theta}^{\dot{a}}] + S_{cy}$$

Choose "chiral" representation where $q_a = \frac{\partial}{\partial \theta^a}$ and $\hat{q}_a = \frac{\partial}{\partial \hat{\theta}^a}$. $a, \dot{a} = 1, 2$

To T-dualize θ^a and $\hat{\theta}^a$, add the trivial surface term

$$S \rightarrow S + \int d^2z C_{ab} (\partial \theta^a \bar{\partial} \hat{\theta}^b - \bar{\partial} \theta^a \partial \hat{\theta}^b) \text{ where } C_{ab} = C_{ba} \text{ is constant.}$$

After T-dualization of θ^a and $\hat{\theta}^a$,

$$S = \int d^2z [\partial x_{a\dot{a}} \bar{\partial} x^{a\dot{a}} + p_a \bar{\partial} \theta^a + \hat{p}_a \partial \hat{\theta}^a + p_{\dot{a}} \bar{\partial} \theta^{\dot{a}} + \hat{p}_{\dot{a}} \partial \hat{\theta}^{\dot{a}} + (C^{-1})^{ab} p_a \hat{p}_b] + S_{cy}$$

This is action for **self-dual graviphoton** background with $F^{ab} = e^{-\Psi} (C^{-1})^{ab}$.

Used in topological strings and for non-anticommutative theories (Ooguri + Vafa)
Seiberg

On higher genus surfaces, flat and self-dual graviphoton backgrounds are not equivalent because of extra fermionic zero modes S_c .

Example 2: $AdS_5 \times S^5$ background

$AdS_5 \times S^5$ sigma model is constructed from Metsaev-Tseytlin currents

$$J^A = (g^{-1} \partial g)^A \quad \text{and} \quad \bar{J}^A = (g^{-1} \bar{\partial} g)^A$$

$$\begin{aligned} A &= (c, \alpha, \hat{\alpha}) \\ c &= 0 \text{ to } 9, (\alpha, \hat{\alpha}) = 1 \text{ to } 16 \end{aligned}$$

where $g(x, \theta)$ takes values in $AdS_5 \times S^5$ coset

$$\frac{PSU(2, 2|4)}{SO(4, 1) \times SO(5)}.$$

$$S = \int d^2z (G_{MN} + B_{MN}) \partial Y^M \bar{\partial} Y^N$$

$$= \int d^2z r_{AdS}^2 [\eta_{cd} J^c \bar{J}^d + (\gamma^{01234})_{\alpha \hat{\alpha}} (J^\alpha \bar{J}^{\hat{\alpha}} - \bar{J}^\alpha J^{\hat{\alpha}})]$$

Action is invariant under global $PSU(2, 2|4)$ isometries $\delta g = \Sigma g$.

Can choose parametrization of g such that four translations

P_m ($m=0$ to 3) and eight chiral susy's q_{aj} ($a=1$ to 2 , $j=1$ to 4) act as

$$x^m \rightarrow x^m + c^m \quad \text{and} \quad \theta^{aj} \rightarrow \theta^{aj} + \xi^{aj} \quad \text{where } (c^m, \xi^{aj}) \text{ are constants.}$$

Since these shift isometries leave action invariant,
can T-dualize x^m and θ^{aj} variables.

After T-dualizing four x^m variables, AdS_5 metric is invariant since

$$\frac{r_{\text{AdS}}^2}{y^2} (dx^m dx_m + (dy)^2) \rightarrow \frac{y^2}{r_{\text{AdS}}^2} d\tilde{x}^m d\tilde{x}_m + \frac{r_{\text{AdS}}^2 (dy)^2}{y^2} = \frac{r_{\text{AdS}}^2}{(y')^2} (d\tilde{x}^m d\tilde{x}_m + (dy')^2)$$

But R-R field-strength and dilaton change as

$$y' = \frac{r_{\text{AdS}}^2}{y} \quad (\text{Kallosh} + \text{Tseytlin})$$

$$e^\varphi F^{*\hat{P}} = (\gamma_{01234})^{*\hat{P}} \rightarrow e^{\varphi'} F'^{*}\hat{P} = (i\gamma_4)^{*}\hat{P}$$

$$\varphi = \text{constant} \rightarrow \varphi' = \text{constant} + 4 \log y$$

If one now T-dualizes eight θ^{aj} variables, one finds


$$g''_{mn} = g'_m n = g_{mn}, \quad \varphi'' = \varphi' - 4 \log y = \text{constant}$$

$$e^{\varphi''} F''^{*\hat{P}} = e^{\varphi'} F'^{*}\hat{P} - e^\varphi \epsilon^{\hat{P}} C^{-1} = (i\gamma_4)^{*}\hat{P} - [(1+i\gamma_{0123})(i\gamma_4)]^{*\hat{P}} = (\gamma_{01234})^{*\hat{P}}$$

So $\text{AdS}_5 \times S^5$ background is invariant (for amplitudes on sphere or disc).

\Rightarrow Alday-Maldacena method relates planar sYM amplitudes and Wilson line correlation function at arbitrary 't Hooft coupling up to a polarization-dependent factor.

When acting on T-dualized variables $(\tilde{x}_m, \tilde{\theta}_{aj})$, original $PSU(2,2|4)$ transformations P_m and q_{aj} become trivial, and conformal and chiral superconformal boosts K^m and s^{aj} become non-local.



Dilatation D changes sign since $y' = \frac{r^2 \cos}{y}$.

In terms of T-dualized variables, "new" dual superconformal transfs are $(\tilde{P}_m, \tilde{q}^{aj}, \tilde{K}^m, \tilde{s}_{aj})$. "Old" dual superconformal transfs form supergroup $SU(2) \times SU(2|4)$. Dual supercont. group was simultaneously found by Drummond, Henn, Korchemsky, Sokatchev.

Conclusions

- Superstring background with abelian supersymmetry is related by fermionic T-duality to a superstring background with fields

$$g'_{mn} = g_{mn}, b'_{mn} = b_{mn}, \varphi' = \varphi + \frac{1}{2} \log C, e^{\psi'} F'^{\alpha\hat{\beta}} = e^{\psi} F^{\alpha\hat{\beta}} - e^{\alpha} e^{\hat{\beta}} C^{-1}$$

$$(\epsilon^x, \epsilon^z) \text{ is Killing spinor satisfying } \begin{aligned} \epsilon^x (\gamma_m)_{\alpha\hat{\beta}} \epsilon^{\hat{\beta}} + \epsilon^z (\gamma_m)_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\beta}} &= 0 \\ \epsilon^x (\gamma_m)_{\alpha\hat{\beta}} \epsilon^{\hat{\beta}} - \epsilon^z (\gamma_m)_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\beta}} &= \partial_m C \end{aligned}$$

- For superstring tree amplitudes, T-dual background is equivalent to original background.
- Under fermionic T-duality, d=4 Minkowski background is mapped to self-dual graviphoton background.
- AdS₅ × S⁵ background is mapped to itself after T-dualizing x^m ($m=0$ to 3) and Θ^{aj} ($a=1$ to 2 , $j=1$ to 4).

Applications of fermionic T-duality

- Explains "dual superconformal symmetry" of perturbative $\mathcal{N}=4$ $d=4$ super-YM amplitudes found by Drummond, Henn, Korchemsky, Sokatchev.
- Relates non-local conserved currents of $AdS_5 \times S^5$ sigma model with dual superconformal generators (Ricci, Tseytlin, Wolf)
(Beisert, Ricci, Tseytlin, Wolf)
- Except for unresolved issue of polarization dependence, extends to arbitrary 't Hooft coupling the Alday-Maldacena proof which relates planar $\mathcal{N}=4$ $d=4$ super-YM amplitudes and Wilson line correlation functions.

Possible future applications

- Use fermionic T-duality to study non-anticommutative structure of self-dual graviphoton background.
- Check invariance of other AdS superstring backgrounds under combination of bosonic and fermionic T-dualities.
- Use fermionic T-duality to enlarge U-duality group of supergravity backgrounds to a U-duality supergroup.
- If E_{10}/E_{11} models of Nicolai et al/West et al could be extended to supergroup models, fermionic supergravity fields might be related to fermionic generators just as bosonic supergravity fields have been related to bosonic generators of E_{10}/E_{11} .

