

Geometry and Topology

SHP Fall '16

Henry Liu

Last updated: December 10, 2016

Contents

0	About Mathematics	2
1	Surfaces	3
1.1	Examples of surfaces	3
1.2	Equivalence of surfaces	4
1.3	Distinguishing between surfaces	6
1.4	Planar Models	8
1.5	Cutting and Pasting	10
1.6	Orientability	13
1.7	Word representations	15
1.8	Classification of surfaces	18
2	Manifolds	19
2.1	Examples of manifolds	20
2.2	Constructing new manifolds	22
2.3	Distinguishing between manifolds	24
2.4	Homotopy and homotopy groups	25
2.5	Classification of manifolds	28
3	Differential Topology	30
3.1	Topological degree	31
3.2	Brouwer's fixed point theorem	33
3.3	Borsuk–Ulam theorem	35
3.4	Poincaré–Hopf theorem	37
4	Geometry	40
4.1	Geodesics	41
4.2	Gaussian curvature	43
4.3	Gauss–Bonnet theorem	47

0 About Mathematics

The majority of this course aims to introduce all of you to the many wonderfully interesting ideas underlying the study of geometry and topology in modern mathematics. But along the way, I want to also give you a sense of what “being a mathematician” and “doing mathematics” is really about.

There are three main qualities that distinguish professional mathematicians:

1. the ability to correctly understand, make, and prove precise, unambiguous logical statements (often using *a lot* of jargon);
2. the ability to ask meaningful questions that lead to interesting problems, tell how difficult these problems will be, and generally identify the easiest and most efficient ways to try to solve them;
3. a deep understanding and knowledge of the area in which they work.


As we go through the content of the course, I’ll point out every now and then how mathematicians apply these abilities to effectively solve problems. But right now, there is one immediate point I want to address.

Here is one of the problems that we will be able to answer later on in the course:

Can you fold a piece of paper over a sphere without crumpling it?

Some of you are probably thinking “well duh, obviously no,” possibly followed by a fairly reasonable-sounding explanation. But how do you know your explanation is actually correct? There are many examples of **intuitive and reasonable explanations that produce completely false conclusions**. Here is one.

Example 0.1. Claim: $2 = \sqrt{2}$. Take a square of side length 1 and repeat the following process to approximate the length of the diagonal:



In each of the steps, the length of the jagged approximation is exactly 2. Since the approximation gets closer and closer to the actual diagonal, the length of the actual diagonal must also be 2. But by Pythagoras’ theorem, the diagonal has length $\sqrt{2}$. So $\sqrt{2} = 2$.

What went wrong? It turns out we made the incorrect assumption that if the jagged line “approximates” the diagonal, its length also approximates the diagonal’s length. Mathematicians use jargon to express ideas precisely enough that they can catch incorrect assumptions like that. If you asked a mathematician to accurately state exactly what went wrong in the example, he/she would probably say something like

the arclength function isn't continuous on the space of piecewise-linear paths with the L^1 norm.

So although mathematical language might seem incomprehensible to you right now, there is often a very simple and intuitive idea being expressed behind the jargon. It is just that in order to express it very precisely to avoid misunderstanding and ambiguity, mathematicians must resort to technical terms that have already been very carefully defined. (For example, you can already start nitpicking at my not-very-precise use of language: what does it mean for the jagged line to “approximate” the diagonal?)

I will introduce a minimal amount of jargon in this course. We only have so much time, and I want to focus on conveying the big ideas. That said, if at any point an argument I am presenting does not make sense to you, or does not sound believable, or you think you found a counterexample, ask! It is likely that either the concepts I used are too vague, or that you have actually discovered a technicality that I have tried to hide!

Let's get started.

In these notes I will write little sidenotes like this one sometimes to make small comments on terminology and notation.

1 Surfaces

Geometry and topology is the study of spaces. Spaces generally have a dimension. We live in a three-dimensional space, but it turns out three-dimensional spaces are much, much harder to study than two-dimensional spaces.

Definition 1.1. A **surface** is a space that, if you were to zoom in on any point, looks like a bent piece of the two-dimensional plane.

Formal definition. A **surface** is a space that is *locally homeomorphic* to \mathbb{R}^2 .

We will actually see the idea of “being homeomorphic” in more detail later.

1.1 Examples of surfaces

The first thing mathematicians (should) do when given a definition is to find as many examples of it as possible! Examples help us develop intuition for what the definition is really saying. It is also important to find examples of objects that do **not** satisfy the definition, so that we develop intuition for what the definition is **not** saying.

Example 1.2. The simplest example of a surface is the two-dimensional plane itself, which we call \mathbb{R}^2 . If we zoom in at any point, it clearly still looks like a two-dimensional plane.

Example 1.3. The **sphere** S^2 and the **torus** T^2 are also surfaces:




Here S^2 does not mean “ S squared.” It is just a label we use for the sphere. The 2 means “two-dimensional.” The same goes for T^2 .

Importantly, these are **hollow**; we only care about their surfaces, not what's inside.

If we zoom in on any point on the sphere, or if we lived on a very big sphere, the region around that point looks very much like a piece of the two-dimensional plane. In fact, the resemblance is so close that for a very long time, we thought we lived on a plane, not a sphere!


Example 1.4 (Non-example). The **cylinder** is **not** a surface:



We usually say the cylinder is a “surface with boundary.” Somewhat confusingly, surfaces with boundary are not surfaces.

If we zoom in at the point p , we get something that is almost a piece of \mathbb{R}^2 , but it has a boundary and we cannot move past it, and so it is not really a piece of \mathbb{R}^2 , where we can move in any direction we want.

Example 1.5 (Non-example). The following space (which doesn't really have a name) is not a surface:




If we zoom in at the intersection point, it will always look like two planes intersecting, not a piece of \mathbb{R}^2 .

1.2 Equivalence of surfaces

One major question that mathematicians always ask whenever they define a new type of object is:


Can we **classify** all the different objects of this type?

But what do we mean by *different*? In other words, when do we consider two surfaces to be **equivalent**? For example, clearly if we take a surface and rotate it, we should consider the result to be the same surface:



Definition 1.6 (Provisional). Two surfaces are equivalent if we can get one from the other by rotating the surface.

The problem with this definition is that there are *too many* different surfaces and it becomes very hard to classify them! For example, we would have to describe exactly how the second cube bulges out on each side:




This is not easy to describe; we would have to specify what the curve of each side looks like. And that's just for a cube. There are many other shapes out there.

Example 1.7. To gain some insight into how we should really be defining equivalence, let's think about **polygons**.

One easy way to classify them is by the *number of sides*, i.e. to say two polygons are **equivalent** if they have the same number of sides.

That way we can *stretch* and *deform* a polygon however we like, and the result will be equivalent to the original polygon:



We have to keep the sides straight though, so that we end up with a polygon!

We use the symbol \sim to denote equivalence, i.e. we write $A \sim B$ to mean that “ A is equivalent to B ,” for whatever definition of equivalence we are using.

We call \sim an “equivalence relation.”

Note that if we instead said two polygons are equivalent if one can be obtained from the other by rotation, classifying polygons becomes much harder: we would have to specify each angle and each side length. We use this insight to give a better definition of equivalence for surfaces.


Definition 1.8. Two surfaces are **equivalent**, or **homeomorphic**, if we can get one from the other by stretching and squeezing and deforming the surface as if it were made from a sheet of rubber.

Formal definition. Two surfaces are **homeomorphic** if there is a *continuous function* from one to the other which has a continuous inverse.


The notion of homeomorphism is actually not specific to two dimensions. We will see later on that it applies to spaces of any dimension. The field of **topology** is the study of spaces under this equivalence.

Being “continuous” is a formalization of the idea that we can deform the surface, but cannot cut it or break it.

Example 1.9. An interesting consequence of our definition is that now a sphere and a cube are actually homeomorphic! The following diagram shows how:



Example 1.10. There is a famous mathematical saying that “topologists can’t tell the difference between a donut and a coffee cup.” This is because those two objects are homeomorphic too!



1.3 Distinguishing between surfaces

Homeomorphism is tricky to work with sometimes. How can we tell that two surfaces are **not** homeomorphic? For example, can you tell if the sphere S^2 and the torus T^2 are homeomorphic or not?



Ask a mathematician this question, and he/she will respond that

In general, to distinguish between objects, define an **invariant** of your objects that assigns the same quantity to equivalent objects.

We will define an invariant called the **Euler characteristic** that assigns an integer to each surface, such that if two surfaces are homeomorphic, they have the same Euler characteristic. That way, if we compute the Euler characteristic for the sphere and the torus and get different integers, we know for sure they are not homeomorphic.

We say the Euler characteristic is “invariant under homeomorphism.”


Definition 1.11. Here are the steps for calculating the Euler characteristic of a surface.

1. Deform the surface until it is a polyhedron, i.e. until it is made up of polygons.
2. Count the number of vertices V , edges E , and faces F on the polyhedron.
3. Calculate $V - E + F$, which is the **Euler characteristic**.

If X is the surface, its Euler characteristic is denoted by $\chi(X)$.

The symbol χ is the Greek letter chi, pronounced “kai.”

Example 1.12. To compute the Euler characteristic $\chi(S^2)$ of the sphere, we first deform it into a cube:




The cube has 8 vertices, 12 edges, and 6 faces, so

$$\chi(S^2) = 8 - 12 + 6 = 2.$$

Wait! What if we deformed the sphere into *some other polyhedron*? Since I said the Euler characteristic is the same for homeomorphic surfaces, it better be that calculating the Euler characteristic using this other deformation gives the same number. But that is not obvious at all.

Example 1.13. Let’s compute the Euler characteristic $\chi(S^2)$ of the sphere by deforming it into a tetrahedron instead:



The tetrahedron has 4 vertices, 6 edges, and 4 faces, so

$$\chi(S^2) = 4 - 6 + 4 = 2.$$

Theorem 1.14. *If two surfaces are homeomorphic, they have the same Euler characteristic.*

A proof of this theorem is actually not hard! But it is long. As we go through a sketch of the proof, it is best to have an example surface in your head to which you can apply the steps in the proof. For example, I usually think of the sphere and the cube, which are two homeomorphic surfaces.

Proof. Let’s sketch the general reason why this is true. Call the two surfaces X and Y . One key idea is that

If we want to compute the Euler characteristic of X , we don’t actually need to do the deformation in step 1 of the process to computing the Euler characteristic.


The rest of this section is somewhat optional material, because the proof of the theorem delves into some technical ideas. That said, it is very interesting and worth skimming through!

Instead, we can just draw out where the vertices and edges are on X (e.g. the diagram above for the cube draws the cube on the sphere). We can do the same for Y . But since X and Y are homeomorphic, we can deform Y until it looks

identical to X . Now we have two copies of X , but they may have different drawings of vertices and edges on them.

So it suffices to show that no matter how we draw vertices and edges on X , we get the same Euler characteristic in the end. This involves the second key idea:

Given two ways of drawing vertices and edges on X , we can “overlap” them to get another way of drawing vertices and edges, called the **refinement**.



The fact that Euler characteristic is unchanged by (or “invariant under”) refinement is an important fact we will use later.

Why does this help? Well, you can check that if we take any two polygons sharing an edge and just remove that edge, the Euler characteristic does not change. Similarly, if we take a polygon and divide it into two adjacent polygons sharing an edge, the Euler characteristic does not change. In other words, *refinement does not change Euler characteristic!* So it must be that the original two different ways of drawing vertices and edges had the same Euler characteristic to start off with. \square

1.4 Planar Models


Now that we know we can compute Euler characteristic of a surface by deforming the surface however we want, let’s compute $\chi(T^2)$, the Euler characteristic of the torus. This sounds simple now: just deform the torus into a polyhedron and count! But it’s hard to keep track of how many vertices and edges and faces there are, and for more complicated surfaces we would have to work really hard to compute the Euler characteristic this way.

It turns out there is a better way, using **planar models**. Planar models will also help us study surfaces more easily in general.

Definition 1.15. A **planar model** of a surface is a polygon whose vertices and edges are **identified**, or “glued together,” in some specified way. To specify that two edges are “glued together,” we label them with the same type of arrow.

The best way to understand a definition is through examples, so let’s look at some planar models for the surfaces we’ve seen already. Then we’ll see some new planar models corresponding to surfaces we haven’t seen yet.

Example 1.16. Here are the planar models for the sphere and the torus, respectively:



So now instead of imagining the surfaces sitting inside three-dimensional space, we can work with the planar models instead.


Example 1.17. Let's use the planar model of the torus to compute its Euler characteristic.

1. The square has 4 vertices, but in the planar model of the torus, every vertex is actually the same vertex, labeled A . So on the torus, there is 1 vertex.
2. The square has 4 edges, but in the planar model, edges a and a are glued together and become the same edge on the torus, and the same for edges b and b . So on the torus, there are 2 edges.
3. There is still only one face; that doesn't change.

We say the four vertices are "identified" to the same vertex.

Hence $\chi(T^2) = 1 - 2 + 1 = 0$.


Some of you may be confused at this point, because we haven't actually deformed the torus into a polygon of any sort. The planar model, when we draw it out on the torus, corresponds to the red and pink lines below:



But that's alright, because just as in the image above, we can perform a **refinement** by adding in new gray edges and vertices (and consequently, new faces). Now of course this deforms into some sort of polyhedron. Because Euler characteristic is unchanged by refinement, we know this procedure doesn't affect our Euler characteristic calculation: $\chi(T^2)$ is still 0.

We defined a refinement earlier, in the proof of Theorem 1.14.

Example 1.18. We can draw arrows in different directions on the square to get new planar models:




Neither of these seem to be the sphere or the torus, but we don't seem to be able to visualize them easily either. In fact they are new surfaces: the **Klein bottle** Kl^2 , and the **projective plane** \mathbb{P}^2 .

We can check that the Klein bottle and projective plane are surfaces: around any point, we can draw a little circle which looks like a bent piece of \mathbb{R}^2 . In fact, any planar model with no boundary will be a surface. The problem with the Klein bottle and the projective plane is that they don't "live inside" three-dimensional space; they are actually objects in four-dimensional space!

Here it is best to draw an analogy: just like a two-dimensional being has problems visualizing how the **Möbius strip** below does not intersect itself, we have problems visualizing how the Klein bottle and projective plane don't intersect themselves.

Later we'll see the deeper reason is Kl^2 and \mathbb{P}^2 are both "non-orientable."

The Möbius strip is actually a surface with boundary.




Exercise. Compute the Euler characteristics $\chi(Kl^2) = 0$ and $\chi(\mathbb{P}^2) = 1$ using the planar models.


1.5 Cutting and Pasting

We have a problem: the Euler characteristic can't tell apart the Klein bottle and the torus, which both have Euler characteristic 0. Maybe they are actually the same surface? The Euler characteristic allows us to tell apart different surfaces; now we will develop a tool that allows us to show two surfaces are equivalent (aside from just "seeing" the deformation).

Example 1.19. Let's look at the following planar model (it is obvious which vertices are identified just by looking at the edge identifications, so from now on I'll start omitting vertex labels):



It has Euler characteristic 0. We'll show it is actually the torus by **cutting and pasting** edges, as follows.




Cutting and pasting techniques can also construct new surfaces! The idea is to somehow glue two surfaces together in a way that produces a new surface.

Definition 1.20. Given two surfaces X and Y , their **connected sum** $X\#Y$ is the surface produced by the following steps:


1. remove a small circle from anywhere on X , and another small circle from anywhere on Y ;
2. take the boundaries where the small circles used to be and glue them together.

Think of the $\#$ as “addition” for surfaces. The notation $X\#Y$ is read “ X connected sum with Y .”

Example 1.21. The connected sum $T^2 \# T^2$ of two torii is a **two-holed torus**:



The connected sum $T^2 \# T^2 \# T^2$ of three torii is a **three-holed torus**:




Of course, we can produce **n -holed torii** for all positive integers n .

We need to make sure the n -holed torii are all inequivalent surfaces, though. We'd like to compute their Euler characteristics using planar models, but we don't know yet what the planar models for connected sums look like. The key idea is that


We can do the connected sum construction *directly on the planar models*.

The steps are as follows (illustrated with $T^2 \# T^2$):

1. Remove a little circle by choosing any vertex, drawing a loop around it, and "opening up" the loop;



2. Glue the resulting two planar models together along the newly created edges.



So the planar model for $T^2 \# T^2$ is an octagon. Now we can compute its Euler characteristic: $\chi(T^2 \# T^2) = -2$. In general, we have the following theorem to help us compute Euler characteristics of connected sums.

The notation $(T^2)^{\#n}$ is sometimes used for the n -holed torus: it stands for $T^2 \# \dots \# T^2$, i.e. T^2 connected sum with itself n times. The idea is that the n is an exponent, and the $\#$ tells us which operation is being repeated n times.

Theorem 1.22. *If X and Y are two surfaces, then $\chi(X\#Y) = \chi(X) + \chi(Y) - 2$.*

Proof. We just have to keep track of how many vertices, edges, and faces are added or removed in the process of constructing the connected sum.

1. We added two extra edges (one on X and one on Y), which are the boundaries of the small circles we remove. Note that we **did not** add extra vertices: the “two” vertices at the ends of the added edge are actually the same vertex!
2. We glued these two new edges (and therefore the vertices at their endpoints) together, i.e. removed two vertices and two edges. By doing this gluing, we also merged the face of X with the face of Y , i.e. we removed a face.

Hence in total we removed one vertex and one face from $\chi(X) + \chi(Y)$, i.e. $\chi(X\#Y) = \chi(X) + \chi(Y) - 2$. □

Corollary 1.23. *The Euler characteristic of the n -holed torus is $2 - 2n$.*

Proof. We know $\chi(T^2) = 0$. Then

$$\begin{aligned}\chi(T^2\#T^2) &= 0 + 0 - 2 = -2 \\ \chi((T^2\#T^2)\#T^2) &= -2 + 0 - 2 = -4 \\ \chi((T^2\#T^2\#T^2)\#T^2) &= -4 + 0 - 2 = -6 \\ &\dots\end{aligned}$$


□

1.6 Orientability

After all that work with cutting and pasting, we still don’t seem to be able to tell apart the Klein bottle Kl^2 and the torus T^2 . (Try cutting and pasting their planar models; it doesn’t seem like we can get from one to the other.) We need another invariant.

Definition 1.24. A surface is **non-orientable** if it contains a Möbius strip. It is **orientable** otherwise.

This is a confusing definition: why a Möbius strip, and why call it “orientable”? Here’s the idea. Imagine a person with one arm living on the Möbius strip who starts from one point, walks around the entire Möbius strip, and returns to his starting point.




Unexpectedly, he is now flipped! What used to be “his left side” is now “his right side.” We see that


On a Möbius strip, the concepts of “right” and “left” do not make sense, because walking around the strip once interchanges them.

It turns out that if a surface does *not contain* a Möbius strip, then “left” and “right” always make sense. Therefore we can pretend **non-orientable** means we cannot distinguish between “right” and “left.”

Example 1.25. The projective plane \mathbb{P}^2 is not orientable. To see this, we look at its planar model, which obviously contains a Möbius strip (whose boundary is marked by the two gray lines) by visual inspection!



Similarly, the Klein bottle Kl^2 is not orientable:



But for more complicated surfaces, we want an easier way of determining whether there is a Möbius strip contained in the surface.


Theorem 1.26. *A surface is non-orientable if and only if its planar model contains two edges glued “in opposite directions,” i.e. two edges both going clockwise, or both going counterclockwise, around the boundary of the polygonal planar model.*

“If and only if” is actually a technical term. We say “ A if and only if B ” to mean that “(if A is true, then B is true) **and** (if B is true, then A is true).”

Proof. We need to prove the following two things in order to prove the theorem:

1. if the planar model of a surface contains two edges glued in “opposite directions,” then the surface is non-orientable;
2. if a surface is non-orientable, its planar model contains two edges glued “in opposite directions”.

The first statement is easy. If a planar model contains two edges glued “in opposite directions,” we can draw a Möbius strip using those two edges. The diagram below is an example, again with gray edges marking the boundary of the Möbius strip.



Conversely, if a surface S is non-orientable, then by definition it contains a Möbius strip. The planar model of the Möbius strip contains two edges glued “in opposite directions,” and these two edges have to be somewhere in the planar model of S . \square

The “converse” of the statement “if A then B ” is “if B then A ,” which is what we’re going to prove now.

Example 1.27. The torus T^2 is orientable, because in its planar model, every pair of edges that are glued together are glued “in the same direction.” So we can finally distinguish between the Klein bottle and the torus: the Klein bottle is non-orientable, but the torus is orientable. They are *different surfaces!*

1.7 Word representations

Although we could state and prove the classification theorem for surfaces right now, we actually want to develop one more tool. The problem right now is that working with planar models, e.g. cutting and gluing, is still a little cumbersome, and we would like a better way to represent planar models without having to draw them out every time. This is an important strategy mathematicians use:

When working with complicated objects, try to represent them using symbols in a way that operations involving the complicated objects translate to very simple operations on the symbols.

(In fact one can claim that this is the strategy underlying *all* of mathematics.)

Definition 1.28. Given a planar model, its **word representation** is obtained as follows:

1. label each distinct edge of the polygon with a different letter (and of course two identified edges have the same label);
2. starting at any vertex, go clockwise around the polygon and read out the letters of each edge, and if an edge goes in the opposite direction, add an inverse sign, e.g. a^{-1} means an edge labeled a but going counterclockwise.

Example 1.29. Here are the word representations of the planar models we've been using for a few surfaces:

1. (sphere) $abb^{-1}a^{-1}$;
2. (torus) $aba^{-1}b^{-1}$;
3. (Klein bottle) $aba^{-1}b$;
4. (projective plane) $abab$.

Of course, a surface can have many different word representations. For example, for the torus, $ba^{-1}b^{-1}a$ is also a word representation, where we started reading clockwise from the top left (instead of the bottom left) corner of the planar model. As another example, we can also relabel edges arbitrarily, e.g. $aba^{-1}b^{-1} \sim cdc^{-1}d^{-1}$. We will need a list of operations we can perform on a word without changing the surface. Before we make this list, we need to make a small observation about word representations.

Lemma 1.30. *In any word representation of a surface, each letter that is used appears exactly twice.*

Proof. If a letter appears more than twice, then more than two edges of the polygon are being glued together. The resulting space is not a surface, because at any point on the edges being glued together, there are more than two directions in which we can move. \square

Now we can make a list of allowed operations on word representations. For example, the operation of "relabel the edge x as y " can be written $AxBxC \sim AyByC$, where A, B, C represent some clumps of letters, and the \sim means the resulting surface is equivalent to the original surface. By the lemma, x appears only twice, so A, B , and C do not contain x and therefore are unaffected by the relabeling. More intuitively, we write $\dots x \dots x \dots \sim \dots y \dots y \dots$.

Theorem 1.31. *The following operations on a word representation will give a word representation of an equivalent surface.*

1. **Relabeling edges:** *replace every edge named x with some other unused symbol y , i.e. $\dots x \dots x \dots \sim \dots y \dots y \dots$.*
2. **Cycle the word:** *move the first letter to the end, i.e. $x \dots \sim \dots x$.*
3. **Merge edges:** *if the same sequence of letters appears twice, merge them into one letter, e.g. $\dots xy \dots xy \dots \sim \dots z \dots z \dots$.*
4. **Cancel xx^{-1} pairs:** *if xx^{-1} occurs in a word for any letter x , then we can remove xx^{-1} from the word, i.e. $Axx^{-1}B \sim AB$.*
5. **Swap edges:** *this is really the two different operations*

$$\begin{aligned} \dots Yxx \dots &\sim \dots xY^{-1}x \dots \\ \dots Yx \dots x^{-1} \dots &\sim \dots x \dots x^{-1}Y \dots \end{aligned}$$

We choose to use this inverse notation because in many ways, a^{-1} should be treated as "1/a," because, for example, we will see that a^{-1} cancels with a when they are next to each other.


The dots are parts of the word representation that remain unchanged by the operation.

Here Y^{-1} means "take the entire clump of letters Y and flip them," e.g. $(ab)^{-1} = b^{-1}a^{-1}$.

Also, the word representation of the connected sum of two surfaces is just the concatenation of their word representations, e.g. $T^2 \# T^2$ has word representation $aba^{-1}b^{-1}cdc^{-1}d^{-1}$.

Proof. It is fairly straightforward that most of these operations give an equivalent surface, but the last operation of moving edges around needs some explanation. Both are obtained via some clever cutting and pasting. For example, the second swapping edges operation, which says $YxAx^{-1}B \sim xAx^{-1}YB$, is proved by the cutting and pasting diagram

We have to relabel the edges of the second torus so that we don't accidentally glue them onto the edges of the first torus.



followed by renaming \tilde{x} back to x . The other equivalence is left as an exercise. \square

The key thing to keep in mind is that all these operations on word representations are just “abbreviations” for operations (generally cutting and pasting operations) on the planar models corresponding to the word representations. But it is *much easier* for us, somehow, to manipulate word representations, as the following examples show.

Example 1.32. The word representation for \mathbb{P}^2 we have right now is $abab$. We write this as $\mathbb{P}^2 \sim abab$. But

$$\mathbb{P}^2 \sim abab \stackrel{3}{\sim} cc \stackrel{1}{\sim} aa,$$

so aa is also a word representation of \mathbb{P}^2 . (The numbers above the \sim specify which operation is being used.)

From now on we will use the word representation $\mathbb{P}^2 \sim aa$.

Example 1.33. The Klein bottle Kl^2 and the connected sum $\mathbb{P}^2 \# \mathbb{P}^2$ of two projective planes both have Euler characteristic 0. (Check this for yourself using the formula in Theorem 1.22 for $\chi(A \# B)$.) Are they equivalent surfaces?

$$Kl^2 \sim aba^{-1}b \stackrel{5}{\sim} baab \stackrel{2}{\sim} aabb \sim \mathbb{P}^2 \# \mathbb{P}^2$$

by swapping edges followed by cycling the word.

Example 1.34. Here is another equivalence that will be useful later on:

$$\begin{aligned} \mathbb{P}^2 \# T^2 &\sim (aab)cb^{-1}c^{-1} \stackrel{5}{\sim} a(b^{-1}acb^{-1})c^{-1} \\ &\stackrel{5}{\sim} a(c^{-1}a^{-1}b^{-1}b^{-1}c^{-1}) \stackrel{5}{\sim} abbac^{-1}c^{-1} \\ &\stackrel{1}{\sim} abba \# \mathbb{P}^2 \stackrel{2}{\sim} aabb \# \mathbb{P}^2 \sim \mathbb{P}^2 \# \mathbb{P}^2 \# \mathbb{P}^2. \end{aligned}$$

1.8 Classification of surfaces

Now we have all the necessary tools to state and prove the classification theorem for surfaces, which will conclude our study of surfaces as an introduction to topology. (We'll go on to study higher-dimensional spaces afterward.) The classification theorem is, in my opinion, one of the most beautiful results in topology, and, indeed, all of mathematics.

Theorem 1.35. *Every (compact, connected) surface is equivalent to one of the following three types of surfaces:*

1. a sphere;
2. a connected sum of projective planes (if it is non-orientable);
3. a connected sum of torii (if it is orientable, and not a sphere).

Proof. The main idea of the proof is, given a word representation, to apply the operations of Theorem 1.31 in a certain order, so that the end result is some sort of “standard form” for the word representation.

1. **Collect like terms into projective planes:** apply the (first) swapping edges operation to move any pair of letters to the beginning of the word, e.g.

$$X(aYa)Z \stackrel{5}{\sim} X(Y^{-1}aa)Z \stackrel{2}{\sim} aaZXY^{-1} \sim \mathbb{P}^2 \# ZXY^{-1}.$$

Now ignore the \mathbb{P}^2 and work with the remaining surface ZXY^{-1} .

2. Repeat the previous step until there are no more pairs of letters xx available (i.e. if x appears, the other edge is x^{-1} , not x). Hence we have written the original word representation as $\mathbb{P}^2 \# \dots \# \mathbb{P}^2 \# W$ where if a appears in the word representation W , then it is glued with a^{-1} . Now ignore the \mathbb{P}^2 and work with the remaining surface W .

3. **Collect interlinked pairs into torii:** apply the (second) swapping edges operation to move letters $\dots a \dots b \dots a^{-1} \dots b^{-1} \dots$ to the beginning of the word, e.g.

$$\begin{aligned} V(aWbXa^{-1})(Yb^{-1}Z) &\stackrel{5}{\sim} (aWbXa^{-1})V(Yb^{-1}Z) = aW(bXa^{-1}VYb^{-1})Z \\ &\stackrel{5}{\sim} a(bXa^{-1}VYb^{-1})WZ = (abXa^{-1}VYb^{-1})WZ \\ &\stackrel{2}{\sim} (bXa^{-1}VYb^{-1}a)WZ = bX(a^{-1}VYb^{-1}a)WZ \\ &\stackrel{5}{\sim} b(a^{-1}VYb^{-1}a)XWZ = (ba^{-1}VYb^{-1}a)XWZ \\ &\stackrel{2}{\sim} (b^{-1}aba^{-1}VY)XWZ = (b^{-1}aba^{-1})VYXWZ \\ &\stackrel{2}{\sim} (aba^{-1}b^{-1})VYXWZ \sim T^2 \# VYXWZ. \end{aligned}$$

Now ignore the T^2 and work with the remaining surface $VYXWZ$.

Being “compact” is a technical detail. All surfaces constructed using planar models are “compact.” Essentially, “compact” means the surface area of the surface is finite. For example, \mathbb{R}^2 is not compact. “Connected” just means the surface is “one piece.” For example, two spheres S^2 sitting side by side is a surface, but is not connected.

The whole point of this tedious computation is to show we can move $aba^{-1}b^{-1}$ to the front of the word.

4. Repeat the previous step until there are no more interlinked pairs of letters $\dots a \dots b \dots a^{-1} \dots b^{-1} \dots$. Hence we have written the original word representation as $\mathbb{P}^2 \# \dots \# \mathbb{P}^2 \# T^2 \# \dots \# T^2 \# W$ where all the letters that remain in W are already matched up in pairs $xx^{-1}yy^{-1}zz^{-1}$.
5. **Cancel** xx^{-1} : using rule 4, cancel all the pairs xx^{-1} in the remaining part W of the word representation.
6. **Conclude**: we have turned the original word representation into something of the form $\mathbb{P}^2 \# \dots \# \mathbb{P}^2 \# T^2 \# \dots \# T^2$. Now there are three cases:
 - (a) if everything canceled and there are zero \mathbb{P}^2 and T^2 , then the result is a sphere;
 - (b) if there are only T^2 and no \mathbb{P}^2 , the result is a connected sum of torii;
 - (c) otherwise if there are both \mathbb{P}^2 and T^2 , use that $\mathbb{P}^2 \# T^2 \sim \mathbb{P}^2 \# \mathbb{P}^2 \# \mathbb{P}^2$ to turn all the torii T^2 into projective planes \mathbb{P}^2 , so the result is a connected sum of projective planes only. \square

Check that the word representation of a sphere is $S^2 \sim aa^{-1} \sim 1$, where 1 represents the “empty word” consisting of no letters at all.

Of course, we should check that these three types of surfaces are not equivalent to each other. This can be done using Euler characteristic along with orientability, and is left as an exercise.

2 Manifolds

The classification theorem for surfaces shows that we understand surfaces, which are two-dimensional objects, very well. The natural question that a mathematician asks at this point (about any result, after proving it) is

Can we generalize this result somehow, and what is the most general statement we can make?

Of course, there are many things to potentially generalize. The classification theorem is a statement about two-dimensional spaces. So:

1. we could try to prove a classification theorem for two-dimensional surfaces *with boundary*;
2. we could also try to prove a classification theorem for *three or higher dimensional* spaces (maybe even with boundary).

It turns out (1) is quite straightforward using the same tools we used for the boundary-less case. But (2) is not. In order to start investigating (2), we must generalize the definition of a surface to higher dimensions.

Definition 2.1. A (topological) **manifold of dimension** n is a space that, if you were to zoom in on any point, looks like a bent piece of the n -dimensional space \mathbb{R}^n .

\mathbb{R}^n is like \mathbb{R}^2 , but instead of two coordinate axes, it has n , all in “different directions.”

Actually, now that we are giving the general definition, it is time to point out that even in our formal definition of a surface, there were several missing technical details.

Formal definition. A (topological) manifold of dimension n is a space that is *locally homeomorphic* to \mathbb{R}^n , and also *Hausdorff* and *second-countable*.

As for surfaces, we have the usual notion of **homeomorphism** for manifolds of dimension n as well.

Being “Hausdorff” and “second-countable” are technical details designed to exclude really weird spaces. It is very hard to be locally homeomorphic to \mathbb{R}^n and not be Hausdorff and second-countable, so we will ignore these details.

2.1 Examples of manifolds


As we did for surfaces, we immediately look for examples of manifolds. We start with two obvious examples.

Example 2.2. Every surface is a manifold of dimension 2.

Example 2.3. The space \mathbb{R}^n is a manifold of dimension n , called the **n -dimensional affine space** or **affine n -space** for short. Just like points in \mathbb{R}^2 are (x, y) for any real numbers x and y , the points in \mathbb{R}^n are (x_1, \dots, x_n) for any real numbers x_1, \dots, x_n .

We must also look for non-examples, so that we can tell what *isn't* a manifold. Of course, one thing that can go wrong is that we have a boundary. Another thing that can go wrong is that some parts of the space have lower or higher dimension than we want.


Example 2.4 (Non-example). The **2-dimensional disk** D^2 is the space obtained by “filling in” the inside of the 1-dimensional circle S^1 :



There is a boundary here, formed by what used to be S^1 . However, aside from the boundary, D^2 is a perfectly valid manifold, so we call it a **manifold with boundary**. The **boundary** of a manifold X with boundary is denoted ∂X .

Example 2.5 (Non-example). Affine n -space \mathbb{R}^n is **not** an m -dimensional manifold for $m \neq n$. This is actually very hard for us to prove right now.

Example 2.6 (Non-example). The following is not a 1-dimensional manifold:




Just as with surfaces, a manifold without boundary is *not* a manifold, by our definition!

No matter how much we zoom in on the origin, there will be four “branches” coming out of it, and it will look like an X.

The problem with higher-dimensional examples is that higher-dimensional manifolds are very hard to visualize. Generally, higher-dimensional manifolds are described as graphs of equations, which, although still hard to visualize, are easier to work with and to manipulate. The following example demonstrates this.

Example 2.7. We will construct the n -dimensional sphere, or “ n -sphere,” for every $n \geq 0$. It is denoted S^n .

1. ($n = 1$) The 1-sphere S^1 is the circle $x^2 + y^2 = 1$ inside \mathbb{R}^2 .
2. ($n = 2$) The 2-sphere S^2 is the sphere $x^2 + y^2 + z^2 = 1$ inside \mathbb{R}^3 .




If a space has a name, e.g. “sphere,” and is n -dimensional, we usually put the n before the name, e.g. “ n -sphere” or “affine n -space.”

Following the pattern, for n in general, we define the n -sphere S^n to be the sphere $x_1^2 + x_2^2 + \dots + x_{n+1}^2 = 1$ inside \mathbb{R}^{n+1} , whose points are of the form (x_1, \dots, x_{n+1}) .

Another method of describing higher-dimensional manifolds is via gluing. Just like we could glue sides of a square to get a 2-dimensional torus (via its planar model), we can glue sides of a cube to get a 3-dimensional torus.

Example 2.8. The following filled-in cube in \mathbb{R}^3 with the given edge (and therefore face) identifications is the 3-torus.



Exercise. A square lives in \mathbb{R}^2 , and a cube lives in \mathbb{R}^3 . How would you describe the higher-dimensional analogues, called **hypercubes**, in \mathbb{R}^n ? Using hypercubes, how would you define the “ n -torus” for larger n ? How about using n -spheres to define the “ n -disk”?

2.2 Constructing new manifolds

We are not done coming up with examples of manifolds! Just like with surfaces, we can also cut and paste with manifolds. But now, because we don't have to restrict ourselves to dimension 2, we can define more general constructions. These constructions also make it easier to describe higher-dimensional manifolds.

Definition 2.9. The **product** of two manifolds M and N is called $M \times N$. It is the manifold obtained by taking every point in M and replacing it with a copy of N (or by taking every point in N and replacing it by a copy of M).

Although this is called a product, we never write MN to mean $M \times N$.

Formal definition. The product $M \times N$ of two manifolds consists of the points


$$M \times N = \{(x, y) : x \in M \text{ and } y \in N\},$$

which is just fancy notation that means "all the points (x, y) such that x is a point in X and y is a point in Y ."

It is not obvious that $M \times N$ is still a manifold! We would have to check that if we zoom on any point in $M \times N$, it looks like a bent piece of \mathbb{R}^k for some k . What should k be? The following examples may shed some light on this.


Example 2.10. The product $\mathbb{R} \times \mathbb{R}$ is just \mathbb{R}^2 . We can see this by taking the product of two line segments, and then imagining that the segments actually extend forever in both directions.

This example gives a good reason for calling affine n -space \mathbb{R}^n : it is literally the product of n copies of \mathbb{R} .




Analogously, the product $\mathbb{R}^2 \times \mathbb{R}$ is \mathbb{R}^3 . In general, $\mathbb{R}^m \times \mathbb{R}^n = \mathbb{R}^{m+n}$, although this is harder to visualize.

Example 2.11. The product $S^1 \times \mathbb{R}$ is an infinitely long cylinder.



Example 2.12. The product $S^1 \times S^1$ is the torus! In fact, we call the product of n copies of S^1 the n -torus, and denote it $(S^1)^n$ or T^n .

Even though we never write MN for the product $M \times N$, it is very common to write M^n to mean the product of n copies of M .




Theorem 2.13. Let M be a manifold of dimension m , and N be another manifold of dimension n . Then $M \times N$ is a manifold of dimension $m + n$.

Proof idea. At every point of $M \times N$, we have m directions to move which come from M , and n extra directions to move which come from N . \square

Another very useful construction is a generalization of the idea of gluing. However, we must be very careful with this construction, because it does not always produce a manifold! In fact, it should actually be defined for any space whatsoever, not just manifolds, which is what we will do.

Definition 2.14. Let X be any space. The idea is to obtain a new space by gluing certain points of X together. The gadget that tells us which points to glue is called an **equivalence relation**, written \sim . If we want to glue two points x and y in X , we set x to be equivalent to y , which we write as $x \sim y$. The **quotient** X/\sim is the space obtained from X by gluing together every pair of points x, y that are equivalent.

Example 2.15. Take D^2 , the 2-dimensional disk. If we squish all of its boundary into a point, we get S^2 , the 2-dimensional sphere:




We say that S^2 is the quotient of D^2 by the equivalence relation where $x \sim y$ if and only if x and y are both in the boundary D^2 .

In the above example, the quotient was particularly simple: we take a part of our space X and squish it into a point. Because this situation is so common, there is a special name and a special notation for it.

Definition 2.16. Let X be any space, and A be any piece of X , called a **subspace**. The **quotient** X/A is the space obtained from X by squishing everything in A into a point. So the above example shows $D^2/\partial D^2 = S^2$.

Example 2.17. Take \mathbb{R}^2 , the 2-dimensional plane. Let the equivalence relation be $(x, y_1) \sim (x, y_2)$ for all real numbers x and y_1 and y_2 . What is the quotient X/\sim ? Well, any two points on the same vertical line are equivalent according to the equivalence relation. So the effect of quotienting is to squish each vertical line into a point.




But then we are just left with an infinite line of points, which is \mathbb{R} .

This is the reason we didn't see the product construction when we covered surfaces: the product of two surfaces is a 4-dimensional manifold.

In general, we use \sim to mean "equivalent." We already did this with surfaces, saying $X \sim Y$ if X and Y are homeomorphic.

In other words, X/A is the quotient X/\sim where $x \sim y$ if and only if x and y are both in A .

Example 2.18. Take D^2 , the 2-dimensional disk. View it inside \mathbb{R}^2 , centered at the origin $(0, 0)$. Let the equivalence relation be $x \sim -x$ for every point x on the boundary of D^2 . We have already seen the quotient D^2 / \sim . It is exactly the planar model for the projective plane \mathbb{P}^2 !



Generalizing, we define the n -dimensional projective space \mathbb{P}^n to be the quotient of D^n by the equivalence relation $x \sim -x$ for every point x on the boundary of D^n .

2.3 Distinguishing between manifolds

Now we have a bunch of manifolds (e.g. \mathbb{R}^n , S^n , T^n , \mathbb{P}^n), and just like with surfaces, we would like to have tools to tell them apart. We had two tools for surfaces: Euler characteristic and orientability. Both of them generalize, but not easily, and they are *not the most useful tools anymore*. For example, Euler characteristic for surfaces used the idea of vertices, edges, and faces, but now we can have higher-dimensional objects.

Definition 2.19. We define higher-dimensional analogues of “faces” by analogy. The idea is to see what makes an edge an edge, and a face a face, and then to extend those properties.

1. A polygon, also called a “2-polytope,” is a 2-dimensional object whose boundary consists of lines (called 1-polytopes).
2. A polyhedron, also called a “3-polytope,” is a 3-dimensional object whose boundary consists of polygons, i.e. 2-polytopes.
3. A **4-polytope** is a 4-dimensional object whose boundary consists of polyhedra.
4. In general, an n -**polytope** is an n -dimensional space whose boundary consists of $(n - 1)$ -polytopes. Vertices are 0-polytopes.

We actually need all these higher-dimensional analogues to be “convex,” which roughly means that they have “no holes.”

The **Euler characteristic** $\chi(M)$ of a manifold M of dimension n is found by deforming the manifold M into a collection of n -polytopes, and then calculating


$$\chi(M) = c_0 - c_1 + c_2 - c_3 + \cdots + (-1)^n c_n$$

where c_k is the number of k -polytopes. (For surfaces, c_0 is the number of vertices, c_1 the number of edges, and c_2 the number of faces.)

Just like Euler characteristic for surfaces using this barebones definition is hard to compute, it is also hard to compute for manifolds.

The process of deforming M into a collection of n -polytopes is (roughly) also known as “putting a CW-complex structure” on M .

Example 2.20. We will compute $\chi(S^n)$ for all $n \geq 0$. First, we look at small n to get some intuition. Here are S^0 , S^1 , and S^2 :



In other words, to get S^1 from S^0 , we add two 1-polytopes (the edges) joining the two 0-polytopes (the vertices). To get S^2 from S^1 , we add two 2-polytopes (the faces, i.e. upper and lower hemispheres) joining the two 1-polytopes (the edges). So in general, to get S^n from S^{n-1} , we just need to add two n -polytopes joining the two $(n-1)$ -polytopes in S^{n-1} .

This shows that in S^n , there are exactly two k -polytopes for every k . So the Euler characteristic is

$$\chi(S^n) = 2 - 2 + 2 - 2 + \dots + (-1)^n 2 = \begin{cases} 2 & n \text{ even} \\ 0 & n \text{ odd.} \end{cases}$$

Exercise (Challenge). This is hard: compute the Euler characteristics

$$\chi(\mathbb{R}^n) = 0, \quad \chi(T^n) = 0, \quad \chi(\mathbb{P}^n) = \begin{cases} 1 & n \text{ even} \\ 0 & n \text{ odd.} \end{cases}$$

We see that in higher dimensions, the Euler characteristic is somewhat useless: it can't even tell apart S^2 and S^4 , and it is also hard to compute. Worse, it is even harder to define orientability in an intuitive way, and it is harder still to check for orientability using only the tools we have so far.

Definition 2.21 (Rough definition). The correct way to generalize “being orientable” from surfaces to n -dimensional manifolds is something like the following. A manifold M is **orientable** if at each point we can consistently define what “clockwise” and “anti-clockwise” mean, so that regardless of how we move around M , it never happens that “clockwise” becomes “anti-clockwise.”

We will not focus on Euler characteristic or orientability anymore. The reason is that there are more powerful invariants that we have yet to talk about.

2.4 Homotopy and homotopy groups

There are two very powerful invariants of spaces in general: homotopy groups, and homology groups. Homotopy groups are hard to calculate but are relatively easier to define; homology groups are easier to calculate but are harder to define. So we begin with homotopy groups, which will enable us to tell apart many of the manifolds we have so far. First, let's see the underlying idea.


This is one possible CW complex structure that we can put on S^n . There are others. A standard one is to use the fact that $D^n / \partial D^n = S^n$, so we can take one n -polytope (homeomorphic to D^n) and glue everything on its boundary to one 0-polytope (a point), so in total there are only two polytopes: one 0-polytope and one n -polytope.

Definition 2.22. A **loop** on a manifold M is a path, with direction and possibly self-intersecting, that begins and ends at the same point, called the **basepoint** of the loop. A loop is **homotopic** to another loop if we can deform the first loop into the second loop while staying inside the manifold and without breaking the loop.

Formal definition. A loop on a manifold M is a continuous function $\gamma(t)$, for $0 \leq t \leq 1$, to the manifold M such that $\gamma(0) = \gamma(1)$, i.e. the start and end points are the same point, the basepoint, in M . A loop γ_0 is homotopic to another loop γ_1 if they are part of a collection of loops γ_s , for every $0 \leq s \leq 1$, such that $\gamma_s(t)$ is a continuous function in (s, t) .

The Greek letter γ , called “gamma,” is often used to denote paths.

Example 2.23. Here are some loops on the sphere S^2 and the torus T^2 :



The two loops drawn on the sphere are homotopic. But the two loops drawn on the torus are not homotopic: there is no way to deform one loop into the other without breaking it. In fact, with a bit more thought, we can see that *any two loops on the sphere are homotopic!*

Since we are trying to construct an *invariant*, an important observation to make at this point is that

The statement “two loops are homotopic” is invariant under homeomorphism. In other words, given two homotopic loops on a manifold M , if we deform it (along with the loops drawn on it) into an equivalent manifold N , the two resulting loops on the manifold N are still homotopic.

Immediately, this gives us a way to distinguish the sphere S^2 from the torus T^2 : any two loops on the sphere are homotopic, but not every two loops on the torus are homotopic. This idea underlies the construction of the homotopy groups. We will define the first homotopy group in the following very long and technical definition. Hopefully the examples that follow it will help.


Definition 2.24. Let M be a manifold and fix a point x_0 in M . The **first homotopy group** (or **fundamental group**) $\pi_1(M, x_0)$ of M is the collection of all non-homotopic loops on M with basepoint x_0 , with the following operations on them:

1. if γ and η are two loops, then we can **concatenate** them to get a loop $\gamma\eta$, which is the loop where we traverse the path given by γ and then traverse the path given by η ;
2. if γ is a loop, we can **reverse** it to get a loop γ^{-1} , which is the loop γ traversed backward.

Warning: $\gamma\eta$ is in general not homotopic to $\eta\gamma$, so order matters! (In other words, concatenation is not commutative.) Can you find an example?

Example 2.25. Fix a point x_0 in S^2 . There is a really silly loop which is the path that goes from x_0 to x_0 by staying fixed at x_0 ; we call this the **constant loop**. Since any two loops on the sphere S^2 are homotopic, every loop is homotopic to the constant loop. So “the collection of all non-homotopic loops on S^2 ” is just the constant loop.

Example 2.26. Let’s try to figure out what the first homotopy group of the circle S^1 is. First of all, we should try to get a sense of what loops on the circle are like. Here are two loops:



Again, we can consider the constant loop. The loop in the diagram above that goes all the way around the circle to return to x_0 is not homotopic to the constant loop, but the other loop in the diagram is indeed homotopic to the constant loop. So we have identified two non-homotopic loops:

1. the “go once around the circle counterclockwise” loop, which we call γ , and
2. the constant loop, which we call γ^0 (for a very good reason that we’ll see soon).

But we need to find *all* the non-homotopic loops! It turns out we can do that by thinking about the concatenation of loops.

What is the concatenation $\gamma\gamma$, also written γ^2 ? Well, it is the loop that goes around the circle twice. Some thought should convince you that γ^2 is not homotopic to γ or to 1. Similarly, γ^3 is not homotopic to γ^2 , γ , or 1. We can continue this pattern, to get γ^n for every $n > 0$.

What about γ^{-1} ? It is just γ , but traversed backward, so it is the “go once around the circle clockwise” loop. Again, some thinking should convince you that γ^{-1} is not homotopic to any γ^n . Now we can concatenate to get $\gamma^{-n} = \gamma^{-1} \cdots \gamma^{-1}$ for any $n > 0$.

Just like with word representations, we pretend the operation of concatenation is like multiplication, and write γ^2 to mean γ concatenated with γ .

Summary: we have constructed non-homotopic loops γ^n for every integer n . It is true that any loop on the circle S^1 is homotopic to one of these loops. So they are all the non-homotopic loops on S^1 . Hence

The first homotopy group $\pi_1(S^1, x_0)$ consists of all the loops γ^n for integers n , with the operation

$$\gamma^n \gamma^m = \gamma^{n+m} \quad \text{for integers } n, m.$$

But we can give a better description. Note that it really doesn't matter what we call the loops γ^n . So let's give them different names; call γ^n just by the integer n . Then what is $\pi_1(S^1, x_0)$? It is the collection of integers, with the concatenation being precisely addition of integers: if we concatenate the loops n and m , then we get the loop $n + m$. So $\pi_1(S^1, x_0)$ is secretly just the **integers**, called \mathbb{Z} , in disguise! Mathematicians like to write $\pi_1(S^1, x_0) = \mathbb{Z}$.

The letter \mathbb{Z} denotes the integers because of the German "Zahlen," meaning "numbers."

That was harder than computing Euler characteristic! But at least we can tell apart S^1 from S^2 and \mathbb{R}^n now, since S^1 has \mathbb{Z} as its first homotopy group, but the first homotopy groups of S^2 and (as an exercise) \mathbb{R}^n both consist of only the constant loop.

Exercise. Convince yourself that $\pi_1(S^2, x_0)$ and $\pi_1(\mathbb{R}^n, x_0)$ consists of just the constant loop (for any choice of basepoint x_0). If the first homotopy group consists of just the constant loop, we say it is **trivial**, or that it is the **trivial group**.

Exercise (Challenge). Convince yourself that $\pi_1(S^n, x_0)$ is trivial for every $n \geq 2$. (Hint: use that $S^n = D^n / \partial D^n$.)

So the first homotopy group can only tell apart S^1 from the rest of the higher-dimensional spheres. Let's see an outline of how to define higher homotopy groups, which can be used to tell apart S^n from S^m for every $n \neq m$.

Definition 2.27 (Rough definition). We can rephrase the first homotopy group $\pi_1(M, x_0)$ of a manifold M as: the collection of non-homotopic ways to draw S^1 in M such that the north pole of S^1 is at the basepoint x_0 . The generalization is the **n -th homotopy group** $\pi_n(M, x_0)$ of M , which is the collection of non-homotopic ways to draw the n -dimensional sphere S^n in M such that the north pole of S^n is at the basepoint x_0 .

A **major unsolved problem** in topology is to find tools that let us compute the n -th homotopy group of S^m for every n and m .

Mathematicians call this the problem of "computing higher homotopy groups of spheres."

2.5 Classification of manifolds

Let's think a little about what we've done so far. We started off with classifying surfaces, and this we did very effectively using planar models and their word representations to glue together, using the connected sum, the "building blocks"

of surfaces: spheres, torii, and projective planes. We have not focused on this approach in the setting of manifolds. Why is that?

The *fundamental issue* with trying to find “building blocks” for manifolds of dimension 3 or higher is the following.

For surfaces, if we cut out disks D^2 from two surfaces and glue along the resulting boundary circles S^1 , there is essentially only one way to glue the two boundary circles together. For higher dimensions, if we cut out disks D^n and try to glue along the resulting boundary circles S^{n-1} , there are many (often infinite) different ways to glue the two boundary spheres S^{n-1} together.

This makes the classification problem for manifolds very difficult in higher dimensions. In fact, it is *impossible* for dimensions 4 or higher, by the following result.

Theorem 2.28 (Proved by Markov, 1958). *There is no algorithm, i.e. series of steps, that can decide whether two given manifolds of dimension $n \geq 4$ are homeomorphic.*

This theorem immediately shows there is no hope of classifying manifolds of dimension n where $n \geq 4$: if there were such a classification of n -dimensional manifolds into a list of different types, given two manifolds we can

1. check which types they are, and
2. declare they are homeomorphic if they are of the same type, or
3. declare they are not homeomorphic if they are not of the same type.

But this is an algorithm for deciding whether two given n -manifolds are homeomorphic, contradicting the theorem! So such a classification cannot exist in dimensions $n \geq 4$.

However, we understand low-dimensional cases very well:

1. ($n = 1$) any connected 1-dimensional manifold is homeomorphic to S^1 or \mathbb{R} ;
2. ($n = 2$) any connected compact 2-dimensional manifold is homeomorphic to a connected sum of S^2 , T^2 , and \mathbb{P}^2 (this is the classification theorem for surfaces that we proved).

Remember that “compact” and “connected” are technical conditions and just mean “finite volume” and “consists of one piece” respectively.

Naturally, mathematicians were curious about what happens in $n = 3$. In 1904, Poincaré decided to tackle a simplified version of the classification problem in $n = 3$: he wanted just to be able to see if a given 3-dimensional manifold M is homeomorphic to S^3 .

Theorem 2.29 (Poincaré conjecture; posed in 1904). *Every connected compact 3-dimensional manifold which has trivial first homotopy group is actually homeomorphic to S^3 .*

Even though we call this a “conjecture,” it is actually a proven result.

Mathematicians worked on this problem for a long time, and it quickly became famous as a particularly tricky problem in the field of topology.

Theorem 2.30 (Generalized Poincaré conjecture for $n \geq 5$; proved by Smale, 1961). *For $n \geq 5$, every connected compact n -dimensional manifold whose homotopy groups agree with those of the sphere S^n is actually homeomorphic to S^n .*

Theorem 2.31 (Generalized Poincaré conjecture for $n = 4$; proved by Freedman, 1982). *Every connected compact 4-dimensional manifold whose homotopy groups agree with those of the sphere S^4 is actually homeomorphic to S^4 .*

Yet Poincaré’s original conjecture, for $n = 3$, remained unsolved, and became one of the 7 Millennium Prize Problems posed by the Clay Mathematics Institute in 2000. As of today (December 10, 2016), only one of these 7 problems has been solved: in 2003, Perelman proved the Poincaré conjecture (in $n = 3$) by actually solving the following harder problem.

Theorem 2.32 (Thurston’s geometrization conjecture; proved by Perelman, 2003). *Every connected compact 3-dimensional manifold can be split into pieces that fall into one of eight different types.*

This solves the classification problem in $n = 3$ as well.

3 Differential Topology


A major theme in modern mathematics is to study not only an object, but also continuous functions defined on the object. For example, instead of studying a manifold M , we can study continuous functions from S^1 to M .

Definition 3.1. A **continuous function** from a manifold M to another manifold N sends each point in M to a point in N , in a way such that if the point x in M is sent to y in N , then points “close to” x in M are sent to points “close to” y in N .

We generally call functions f or g or h . If f maps from M to N , we write $f: M \rightarrow N$.

Formal definition (ϵ - δ definition of continuity). A function $f: M \rightarrow N$ is **continuous** if at every point x in M , given a small number $\epsilon > 0$, you can find another small number $\delta > 0$ such that every point within a distance δ of x is sent to a point within a distance ϵ of y .

Example 3.2. A continuous function f from \mathbb{R} to \mathbb{R} is essentially any function that can be graphed on the two-dimensional plane “without lifting the pencil.”




For example, the function on the left is continuous, but the function on the right is discontinuous.

Example 3.3. A continuous function f from the circle S^1 to a manifold M is just a loop inside M ! Note that we implicitly assumed loops to be continuous when we first started talking about them.

We have already seen that studying loops inside manifolds is very useful, by defining the first homotopy group and using it as an invariant. In general, studying continuous functions on manifolds is very useful. However, we often want to restrict ourselves to a specific kind of continuous function.

Definition 3.4. A **differentiable function** is a continuous function where at each point it makes sense to talk about the slope at that point.

Example 3.5. Here are functions from S^1 to \mathbb{R}^2 , i.e. two loops in the plane \mathbb{R}^2 .



The first one is differentiable, because at every point the slope is well-defined. The second one is not differentiable at the tip of the spike, because there is no meaningful “slope” there. However note that both loops are still continuous.

The reason we want to work with differentiable functions instead of just continuous functions is that calculus works for differentiable functions and not for continuous functions. Being able to do calculus with differentiable functions on manifolds is very helpful in investigating properties of manifolds in general. In fact, there is an entire subfield of topology, called **differential topology**, where we investigate differentiable functions on (differentiable) manifolds.

In this section, we will develop some basic tools of differential topology and use them to prove some cool results. For example, one of the results we will prove, called Brouwer’s fixed point theorem, can be used to show that

If I take two pieces of paper of equal size, crumple one up and put it on top of the other one, then there is at least one point on the crumpled sheet of paper that lies directly above the same point on the flat sheet.

3.1 Topological degree

The first tool we will develop is called the topological degree of a continuous function. Since this will be hard to define for arbitrary continuous functions,

For the idea of “differentiable function” to make sense on a manifold, we actually need that the manifold itself is “differentiable,” which roughly means it has no spikes or jumps or disconnections.


we'll focus on continuous functions from S^1 to S^1 . These will be the functions we need to use in the proof of Brouwer's fixed point theorem.

Example 3.6. We have already looked at continuous functions from S^1 to S^1 : these are precisely loops on S^1 ! When we computed the first homotopy group $\pi_1(S^1)$ of the circle, we identified many such functions. For example, there were:

1. the constant loop γ^0 ;
2. the loop γ^1 that goes once around the circle counterclockwise;
3. the loop γ^{-1} that goes once around the circle clockwise;
4. concatenations of these loops.

Definition 3.7. We computed earlier that the first homotopy group of the circle is $\pi_1(S^1) = \mathbb{Z}$. This shows that given a loop $f: S^1 \rightarrow S^1$, it is homotopic to some γ^n for exactly one integer n . The **(topological) degree** of the function f is this integer n . We write $\deg f = n$. Note that two functions of different degree cannot be homotopic.

Example 3.8. The degrees of the two loops in the image below are 1 and 0:



Topological degree can actually be defined for more arbitrary continuous functions. In the special case of functions $S^1 \rightarrow S^1$, it is also commonly called “winding number,” because it is the number of times a loop winds around the circle.

This is because the loop on the left is exactly γ^1 , while the loop on the right is homotopic to γ^0 .

Although this should be a familiar concept by now because we are used to working with loops, it is sometimes hard to interpret functions from S^1 to S^1 as loops and to compute their degree. Let's do two very important examples.

Example 3.9. Treat S^1 as the unit (i.e. radius 1) circle centered at the origin in the two-dimensional plane.

1. Consider the **identity function** $f(x) = x$ on S^1 . This is the loop γ^1 , and therefore has degree 1.
2. Consider the **antipodal function** $f(x) = -x$ on S^1 . This is again the loop γ^1 , and therefore has degree 1.
3. Consider the **constant function** $f(x) = c$ on S^1 , for some constant c . This is the constant loop γ^0 , and therefore has degree 0.

The identity function is so commonly used that it has its own symbol: we denote it by id .


In particular, the identity function is not homotopic to the antipodal function, because they have different degree.

Note that we have actually done nothing new so far; all this is just a rephrasing of what we did in our computation of $\pi_1(S^1) = \mathbb{Z}$. But topological degree can be defined more generally, and so it is important to know it exists as a concept.

3.2 Brouwer's fixed point theorem

Using the idea of topological degree, we can prove Brouwer's fixed point theorem, one of the fundamental theorems in differential topology. Throughout this subsection, it helps to think of D^2 inside \mathbb{R}^2 centered at the origin.

Theorem 3.10 (Brouwer's fixed point theorem). *Any continuous function $f: D^2 \rightarrow D^2$ from the disk D^2 to itself has a fixed point, i.e. there exists a point x in D^2 such that $f(x) = x$.*




Proof. This proof is a bit more complicated than the ones we have seen so far. So, first, here is a rough outline of how the proof goes.

1. Suppose that there exists a function $f: D^2 \rightarrow D^2$ that has no fixed point.
2. Use it to construct a function $F: D^2 \rightarrow \partial D^2$ that is the identity map on ∂D^2 , i.e. $F(x) = x$ for every x on the boundary of D^2 .
3. Prove, using topological degree, that such a function F cannot exist.
4. Since this is a contradiction, our initial assumption that the function f exists must be false. Then we are done the proof.

Okay, let's do it!

Suppose there exists $f: D^2 \rightarrow D^2$ with no fixed point. That means $f(x) \neq x$ for every x in D^2 . Then we can draw a line segment from $f(x)$ to x , and continue drawing until it hits the boundary of D^2 .



Call the point where it hits the boundary $F(x)$. Hence we have constructed a function $F: D^2 \rightarrow \partial D^2$, sending every point in D^2 to some point on its boundary ∂D^2 . In particular, we see that $F(x) = x$ for every point x on the boundary of D^2 .

Now we show this map F cannot exist. First, since $\partial D^2 = S^1$, we can write F as $F: D^2 \rightarrow S^1$. If we look at F only on ∂D^2 , it is the identity map $S^1 \rightarrow S^1$, and therefore has degree 1. But ∂D^2 can be continuously deformed into, i.e. is homotopic to, a constant loop inside D^2 . If we look at F only on this constant loop, it is the constant map $S^1 \rightarrow S^1$ and therefore has degree 0. Hence F gives a way to continuously deform a degree 1 function into a degree 0 function. But this is impossible, since functions of different degree cannot be homotopic.

We have reached a contradiction. So our initial assumption that f exists must be incorrect. Hence every function $f: D^2 \rightarrow D^2$ must have a fixed point. \square

We used a key fact in this proof that is an important technical tool in its own right. These technical tools, to be used later in proofs of bigger things, are generally called **lemmas**. Here is the important lemma we used in the above proof.

Lemma 3.11. *If a function $F: S^1 \rightarrow S^1$ actually comes from looking at a part of a function $F: D^2 \rightarrow S^1$, then the degree of F must be zero.*

Proof. Here's a brief summary of how we proved this lemma in the above proof: such a function $F: D^2 \rightarrow S^1$ tells us how to make a homotopy from $F: S^1 \rightarrow S^1$ to the constant loop, which is degree 0. Since degree is unchanged by homotopy and the constant loop has degree 0, the function F has degree 0 as well. \square

An interesting application of Brouwer's fixed point theorem is to the Nash equilibrium theorem, which very roughly says that given any game with multiple players, there exists a strategy for each player such that, assuming the other players don't change their strategies, each player cannot gain anything by changing their own strategy, i.e. their own strategy is in some sense "the best possible."

3.3 Borsuk–Ulam theorem

Next we will use topological degree to prove the Borsuk–Ulam theorem.

Theorem 3.12 (Borsuk–Ulam). *If f is a continuous function from S^2 to \mathbb{R}^2 , then there exists a point x on the sphere S^2 such that $f(x) = f(-x)$.*

Here is a more intuitive way to see what the Borsuk–Ulam theorem says. Imagine the surface of the Earth; this is a two-dimensional sphere S^2 . At each point on the surface, we can measure two values: temperature T , and pressure P . Together, (T, P) is a point in \mathbb{R}^2 . So we have a continuous function that, for each point on the surface of the Earth, gives the temperature and pressure at that point. This is a continuous function from S^2 to \mathbb{R}^2 . The Borsuk–Ulam theorem then says that

At any moment, there exist two points on the surface of the Earth, directly opposite each other, where the temperature and pressure are the same.

This is somewhat non-intuitive! We will see, however, that the same ideas that go into proving Brouwer’s fixed point theorem can also be used to prove the Borsuk–Ulam theorem. In fact, the proof technique will be very similar.

Proof of Borsuk–Ulam. As with the proof of Brouwer’s fixed point theorem, here is an outline of the proof of Borsuk–Ulam.

1. Suppose that there exists a function $f: S^2 \rightarrow \mathbb{R}^2$ such that $f(x) \neq f(-x)$ for every x on S^2 .
2. Use it to construct a function $F: S^1 \rightarrow S^1$ such that $F(-x) = -F(x)$ for every x on S^1 .
3. Prove, using topological degree, that such a function F cannot exist.
4. Since this is a contradiction, our initial assumption that the function f exists must be false. Then we are done the proof.

Okay, let’s do it!

Suppose that there exists a function $f: S^2 \rightarrow \mathbb{R}^2$ such that $f(x) \neq f(-x)$ for every x on S^2 . Then let

$$h(x) = \frac{f(x) - f(-x)}{\|f(x) - f(-x)\|}.$$

Note that $h(x)$ has two important properties:


1. the value of $h(x)$ for any x is on the circle S^1 ;
2. $h(-x) = -h(x)$, since

$$h(-x) = \frac{f(-x) - f(x)}{\|f(-x) - f(x)\|} = -\frac{f(x) - f(-x)}{\|f(x) - f(-x)\|} = -h(x).$$

Note that we really require $f(x) \neq f(-x)$ for every x , so that the denominator of $h(x)$ is never zero; we can’t divide by zero.

But we wanted a function $F: S^1 \rightarrow S^1$, and right now we only have a function $h: S^2 \rightarrow S^1$. This is easy to solve: just look at h only on the equator of S^2 . The equator is just a circle S^1 , so if we focus only on the equator, we obtain a function $S^1 \rightarrow S^1$ which we will call F . Since $h(-x) = -h(x)$ is true everywhere on S^2 , it is true in particular on the equator, so $F(-x) = -F(x)$ as well.

Now we show such a function $F: S^1 \rightarrow S^1$ satisfying $F(-x) = -F(x)$ cannot exist. First we show the degree of F must be odd. This means we have to interpret F as a loop, and see how many times it goes around the circle. Consider F on half the circle S^1 , starting from y and ending at $-y$:



Since $F(-y) = -F(y)$, whatever the resulting path is, it ends halfway across the circle. But also because $F(-x) = -F(x)$, the other half of the loop, starting from $F(-y)$ and ending at $F(y)$, is just a mirror image of the first half of the loop. Hence F , as a loop, goes around the circle an odd number of times, i.e. it has odd degree. However, we can also apply lemma 3.11 to show F has degree 0. This is because we can look at $h: S^2 \rightarrow S^1$ only on the upper hemisphere of the sphere S^2 , so that it becomes a function $g: D^2 \rightarrow S^1$. The equator is part of the upper hemisphere (and also the lower hemisphere, for that matter), so $F: S^1 \rightarrow S^1$ actually arises as a little part of the function $g: D^2 \rightarrow S^1$. Then the lemma shows us that the degree of F is zero. Hence the degree of F is both odd and zero at the same time.

We have reached a contradiction. So our initial assumption that f exists must be incorrect. Hence for every function $f: S^2 \rightarrow S^1$, there exists a point x on S^2 such that $f(x) = f(-x)$. \square

There are some interesting consequences of the Borsuk–Ulam theorem once we write down its most general form, in n dimensions (instead of in 2). Here is one such consequence, in two different flavors.

Corollary 3.13 (Ham sandwich theorem). *Given 3 finite-volume (closed) subsets of \mathbb{R}^3 , there exists a plane dividing the volume of the three pieces exactly in half simultaneously.*




Consequences of theorems are called “corollaries.”

Here, “closed” is a very technical restriction that we are going to ignore.

This is called the ham sandwich theorem because we can imagine two of the subsets being pieces of bread, and the other subset being a slice of ham. Then the theorem says there always exists a way we can slice the sandwich in half, so that there is exactly the same amount of bread and ham in the two pieces.

Corollary 3.14 (Discrete ham sandwich theorem). *Given any number of red dots and blue dots arranged in any way in the plane, there exists a line dividing the plane into two regions such that the number of red dots on each side is equal, and the number of blue dots on each side is equal.*



It turns out, actually, that Brouwer’s fixed point theorem is also a consequence of the Borsuk–Ulam theorem: that’s how powerful the Borsuk–Ulam theorem is!


3.4 Poincaré–Hopf theorem

Let’s think about the surface of the Earth some more. The Borsuk–Ulam theorem shows that there always exist antipodal points on the surface of the Earth with the same temperature and pressure. Temperature and pressure are both functions that assign a *number* to each point on the surface of the Earth. What if we wanted to model something like wind? We would need a function that assigns a direction to each point. More precisely, we want to assign a *vector* to each point, so that we know both the direction and the magnitude of the wind.


Two points on the sphere are “antipodal” if they are directly opposite each other.

Definition 3.15. Let M be a manifold. A vector is **tangent** to the manifold M , if the direction in which it points is a direction in which you can move and stay in M . A **vector field** on a manifold M is a function that assigns a tangent vector to each point in M in a continuous fashion. (In other words, given a point x in M , the vectors at points around x should not be too different from the vector at x .)

Example 3.16. Here are examples of tangent vectors on the circle S^1 and the sphere S^2 :



Here are examples of non-tangent vectors on the circle S^1 and the sphere S^2 :




A vector field on the sphere S^2 therefore models wind on the surface of the Earth: at each point, there is a vector telling us which direction the wind is blowing at that point, and how strongly it is blowing. So one question we can ask is:

on a given manifold M , does there exist a vector field that is non-zero everywhere?

If our vector field represents wind, then this question is: can there be wind everywhere on the surface of the Earth at the same time?

Example 3.17. There does exist a vector field on the torus T^2 that is non-zero everywhere, shown (somewhat crudely) by the following diagram.

Of course, the important (and realistic) assumption is that wind is continuous.



What about the sphere S^2 ? After a few tries, it may start to seem impossible to draw a vector field on S^2 that is non-zero everywhere. It turns out that it is indeed impossible. So there exists a vector field that is non-zero everywhere on the torus T^2 , but there does not exist such a vector field on the sphere S^2 . Whether or not such a vector field exists actually depends on the topology of the manifold!

Theorem 3.18 (Hairy ball theorem). *Every vector field on the sphere S^2 is zero somewhere.*

Proof sketch. Since we have already seen two very long proofs (of Brouwer’s fixed point theorem and of the Borsuk–Ulam theorem), let’s just go through the outline of this proof.

1. Suppose there exists a vector field on the sphere that is non-zero everywhere.
2. Define a function $f: S^2 \rightarrow S^2$ which takes a point x and maps it to the point y given by moving a little bit, from x in the direction of the vector at x .
3. Show that f is homotopic to the identity map.
4. Since the vector field is non-zero everywhere, $f(x) \neq x$ for every x , i.e. f has no fixed points. Use this to construct a homotopy from f to the antipodal map.
5. Use topological degree to show this is a contradiction: the degree of the identity map is 1, but the degree of the antipodal map is 2. \square

This is called the “hairy ball theorem” because we can imagine a ball with lots of hair on its surface. If we can comb all the hair flat, then each hair can be interpreted as a tangent vector, so we would have a vector field that is non-zero everywhere. So sometimes people state the hairy ball theorem as “you can’t comb a hairy ball flat.”

Corollary 3.19. *At any moment in time, there is a point on the surface of the Earth with no wind.*

Actually, the hairy ball theorem is a special case of a much more general theorem. To state this more general theorem, we need to look more carefully at the points where the vector field is zero; it turns out there are “different ways” in which the vector field can be zero.

Definition 3.20. Consider a vector field V on a manifold M . Suppose that at p , the vector field is zero. The **index of V at p** can be calculated as follows:

1. draw a small circle around p , so that at each point of the circle is a vector;
2. count the number of times N the vector “turns” as we move counterclockwise around the circle;
3. if the vector turned N times counterclockwise, the index is N ; otherwise if it turned N times clockwise, the index is $-N$.

In fact, this definition works even if the vector field is not zero at the point p . But it turns out that for such p (where the vector field is non-zero), the index is always 0, so we generally don’t care about these points.

Example 3.21. The indices of the following vector fields are 0, 1, -1 , -2 .



For the purposes of computing index, the direction of the vectors don’t matter.

Theorem 3.22 (Poincaré–Hopf). *Let V be a vector field on M . The sum of the indices at every point where V is zero is equal to the Euler characteristic $\chi(M)$.*

This is usually written as $\sum_p \text{ind}_p(V) = \chi(M)$.

Corollary 3.23 (Hairy ball theorem). *Every vector field on the sphere S^2 is zero somewhere.*

Proof. Assume V is a vector field on S^2 which is non-zero everywhere, then the sum specified in the Poincaré–Hopf theorem is zero. So $\chi(S^2) = 0$. This is a contradiction, because we know $\chi(S^2) = 2$. So such a vector field V cannot exist: it must be zero somewhere. \square

But Poincaré–Hopf can give us more than just the hairy ball theorem. By the same argument as above, we have proved the following.

Corollary 3.24. *If $\chi(M) \neq 0$, then every vector field on M is zero somewhere.*

In particular, let’s think about surfaces. By the classification theorem for surfaces, there are only two surfaces with Euler characteristic zero: the torus T^2 , and the Klein bottle Kl^2 . Vector fields on every other surface must all be zero somewhere.

Exercise. Find a vector field on the Klein bottle that is non-zero everywhere.


4 Geometry

We are now going to upgrade the idea of “homeomorphism.” Remember that two spaces are homeomorphic if one can be deformed into the other. However, this process of deformation may change the *distance* between two points in the space, e.g. a big sphere is homeomorphic to a smaller sphere, but the distance between any two given points changes.

Sometimes we care about the distances between points, and don’t want the distance to change when we consider “equivalent” spaces. So we need a stricter notion of whether two spaces are “equivalent.”

Definition 4.1. In the field of **geometry**, two spaces are equivalent if they are homeomorphic in a way that preserves distances. Such a homeomorphism is called an **isometry**.

Example 4.2. Let’s look at isometries of the two-dimensional plane \mathbb{R}^2 . The best way to visualize whether a deformation is an isometry for \mathbb{R}^2 is to imagine \mathbb{R}^2 as a flat piece of paper. Then any transformation we can imagine applying to the paper must preserve the distances between points, because the paper itself can’t stretch or shrink.



So rolling up the paper to get a cylinder $S^1 \times \mathbb{R}$ is an isometry, but intuitively it does not seem possible to get a sphere S^2 .

Before we continue, note that distance is not *intrinsic* to a space. If I give you an arbitrary space by telling you what the points in the space are, what is the distance between two given points? This information is not intrinsically contained in the space. Instead, there are generally two ways to specify distances on a space X .

1. The first way is to embed X into \mathbb{R}^n for some n , and then use the usual idea of distance on \mathbb{R}^n to measure distances on X . The **distance** between two points p and q on X is the length of the shortest curve in X from p to q . For example, the distance between the north and south poles on a sphere S^2 of radius 1 is exactly π .
2. The second way is to specify a **distance function** $d(p, q)$ on X that explicitly tells you the distance from point p to point q for any two points p and q . Such a distance function must satisfy some properties:
 - (a) $d(p, q) \geq 0$ for any points p, q , and equality holds if and only if $p = q$ (“the distance between any two points cannot be negative, and is zero only when the two points are the same”);
 - (b) $d(p, q) = d(q, p)$ (“the distance from p to q is the same as the distance from q to p ”);
 - (c) $d(p, r) \leq d(p, q) + d(q, r)$ (“the distance from p to r is at most the distance from p to an intermediate point q plus the additional distance from q to r ”).

We will stick with the first method, which is much simpler to work with.

4.1 Geodesics

From now on, M is a manifold embedded into \mathbb{R}^N . So the distance between two points p and q on M is the length of the shortest curve in M connecting p and q . However, such a shortest path is often very hard to find, and very hard

Although we are calling it a cylinder, there is no actual gluing happening; we are not allowed to glue for isometries. Similarly, the sphere is not actually a sphere.

Actually, for a manifold, it is more appropriate to specify an “infinitesimal distance function” called a *metric* that acts like a dot product on tangent vectors. Then the length of a path is the integral along the path of the lengths of the tangent vectors of the path, and the distance is the length of the shortest path.

to work with. Imagine if you were an ant on a surface. You don't know the shortest path to a point a million miles away, but at least you can try to walk "in a straight line" toward it.

Definition 4.3. A **geodesic** on M is a curve on M that is *locally* distance-minimizing. In other words, if we zoom in far enough, the geodesic is the shortest path between two points (which are close to each other).


This definition illustrates the main philosophy behind doing things with manifolds:

Because manifolds locally look like \mathbb{R}^n , and we understand \mathbb{R}^n very well, we try to define objects and concepts only locally as much as possible.

A good way to imagine geodesics on a surface is to imagine taking a rubber band and nailing it to the surface at p and q . The path it shrinks to is always a geodesic.


Example 4.4. Geodesics on \mathbb{R}^n are just straight lines. So a good way to think about geodesics is that they are generalizations to arbitrary manifolds of the idea of a straight line in \mathbb{R}^n .

Example 4.5. Geodesics on the sphere S^2 are **great circles**.




This is why airplanes fly in "curved" paths around the Earth. Though on a flat map they may seem like longer paths, these "curved" paths are actually the shortest paths!

Example 4.6. Here are two different kinds of geodesics on the cylinder $S^1 \times \mathbb{R}$:




Note that the one that goes around the cylinder is indeed locally distance-minimizing, but is clearly longer than the distance r between the two points. The straight line geodesic between the two points is the shortest path.

Example 4.7. Note that geodesics may not always exist! Imagine the plane \mathbb{R}^2 with a hole in it:



Then there is no geodesic between p and q , since any locally distance-minimizing path must cross the hole.

In general relativity, geodesics are very important: the paths that objects take through spacetime are always geodesics (as long as no external forces other than gravity act on them), with the distance function determined by gravity.



You may have seen pictures of the “spacetime fabric”: these are visualizations of what spacetime would look like if it were embedded into \mathbb{R}^n . The gravity of massive objects like planets and stars “curve” the spacetime around them. Objects orbit other objects because in those curved regions of spacetime, orbits are the geodesics, not straight lines.


4.2 Gaussian curvature

From what we have seen so far, the geometry of a space is affected by how “curved” it is, while in topology it really didn’t matter, because all the curves could be made flat via deformations. So it makes sense that the idea of “curvature” is a big deal in geometry. In fact, we will see soon that curvature is an isometry invariant, so it helps us distinguish non-isometric manifolds.

We want to somehow assign a number to each point of a manifold that says how curved the manifold is at that point. There are actually many ways this can be done. We will see one of them, for surfaces only. From now on, M is an oriented surface in \mathbb{R}^3 .

Definition 4.8. To define curvature for surfaces, we must first define curvature for curves, i.e. 1-dimensional manifolds, say inside \mathbb{R}^3 . Let C be a curve and p be a point on C . The **curvature** of C at p calculated using the following steps.

1. Draw the **osculating circle** of C at p : this is the circle which is tangent to the curve at p , and which “hugs” the curve the tightest.




If the curve is a line, then the osculating circle has “infinite” radius.

2. Let r be the radius of the osculating circle. The curvature is $1/r$. (If the osculating circle has “infinite” radius and the curvature is 0.)

Example 4.9. Let M be a circle with radius r . Then the curvature of M at any point p is just $1/r$, because the osculating circle at p is precisely M itself and therefore has radius r .

Definition 4.10. Let p be a point on M . The **Gaussian curvature** of M at p is calculated using the following steps.

1. Pick a **normal vector** at every point close to p . A normal vector at the point p is a vector perpendicular to the surface at p ; more precisely, it is perpendicular to every tangent vector at the point p .
2. Consider a cross-section of the surface M that contains p and the normal vector at p . In this cross-section, M looks like a curve. Let κ be the curvature of this curve.




The **normal curvature** of M with respect to this cross-section is κ if the normal vector points “into” the osculating circle, and $-\kappa$ otherwise.

3. Let κ_1 and κ_2 be the maximal and minimal normal curvatures at p , across all possible cross-sections.
4. The Gaussian curvature is the product $K = \kappa_1\kappa_2$.

Example 4.11. The Gaussian curvature K of a sphere S^2 with radius r is $1/r^2$. This is because every normal curvature at any point is $\pm 1/r$, because every cross-section is a (portion of a) circle of radius r . In particular, $K > 0$ for any sphere.


Example 4.12. The Gaussian curvature of the plane \mathbb{R}^2 is 0. This is because every normal curvature at any point is 0, because every cross-section is a straight line. So $K = 0$.

Example 4.13. The Gaussian curvature of the cylinder $S^1 \times \mathbb{R}$ is also 0.



This is because at any point, the maximal normal curvature $\pm 1/r$ is given by the cross section along S^1 , and the minimal normal curvature 0 is given by the cross section along \mathbb{R} . Hence $K = (\pm 1/r) \cdot 0 = 0$.

Example 4.14. The Gaussian curvature of the following surface at the marked point is negative.



This is because regardless of which direction the chosen normal vector points, the maximal normal curvature is positive, and the minimal normal curvature is negative. So $K < 0$.

Note that the cylinder and the plane have the same Gaussian curvature, and we know they are isometric (by “rolling up” the plane). This is some evidence for the following theorem.

Theorem 4.15 (Gauss’s Theorema Egregium). *Gaussian curvature is invariant under isometries, i.e. two isometric surfaces always have the same Gaussian curvature.*

In particular, the theorem implies that the sphere is not isometric to the plane, just as we suspected. This means a piece of paper can’t be made into a sphere without creasing it or ripping it somewhere.

Equivalently, a sphere can’t be somehow “flattened” without distorting some part of its surface. This is why maps of the Earth never accurately represent distance everywhere.

Mapmakers have many different systems for approximating this “flattening” process so that only certain regions (e.g. near the north or south pole) are distorted.

“Theorema egregium” literally means “remarkable theorem” in Latin.




For example, in the Mercator projection system (shown above), areas around the poles seem much bigger than they should be; Greenland looks huge in comparison to Australia, even though in reality Australia is much bigger.

4.3 Gauss–Bonnet theorem

It is amazing that the Gaussian curvature is invariant under isometry, since it is not true that normal curvatures are invariant under isometry. It is even more amazing that the Gaussian curvature, which is a number associated to the geometry of the surface, is directly connected to the Euler characteristic, which is a number associated to the topology of the surface. This connection arises by looking at the “average value” of the Gaussian curvature over a polygon on a surface.

Definition 4.16. A polygon in the two-dimensional plane \mathbb{R}^2 is just a region bounded by straight lines. For a general orientable surface S , a **polygon** in S is a region bounded by geodesics, which are supposed to be the generalizations of straight lines to arbitrary manifolds.


Example 4.17. What is a triangle on a sphere? Here is one example.



Note an interesting property of this triangle: each of its internal angles is $\pi/2$, so the sum of its internal angles is actually $3\pi/2$.

On the two-dimensional plane \mathbb{R}^2 , we know from basic Euclidean geometry that the sum of internal angles for any triangle is π . The reason we can get more than π on the sphere seems to be because the sphere is curved. So perhaps we can find a relationship between curvature and the sum of some internal angles.

Definition 4.18. Given a polygon P , the **exterior angle** at a vertex of P is π minus the interior angle at P .



Theorem 4.19 (Local Gauss–Bonnet formula). *Let S be a (compact) orientable surface, and $K(x)$ be the Gaussian curvature at the point x in S . If P is a polygon in S , then*

$$\int_P K(x) dx + (\text{sum of exterior angles of } P) = 2\pi\chi(P).$$

The technical condition “compact” essentially just means “finite surface area.”


Example 4.20. Let's check the Gauss–Bonnet formula for polygons on the plane. Let \mathbb{R}^2 be the two-dimensional plane, and let P be an n -sided polygon on it. We will compute each of the terms in the formula.

1. (The $\int_P K(x) dx$ term) The normal curvatures at every point of the plane are 0, since every cross-section is just a straight line. Hence $K(x) = 0$ for any point on \mathbb{R}^2 . So $\int_P K(x) dx = 0$.
2. (The (sum of exterior angles of P) term) We will leave this term as it is.
3. (The $2\pi\chi(P)$ term) An n -sided polygon has n vertices, n edges, and 1 face, so $2\pi\chi(P) = 2\pi(n - n + 1) = 2\pi$.

Hence the Gauss–Bonnet formula simplifies to

$$(\text{sum of exterior angles of } P) = 2\pi.$$


This is a true fact about polygons in the plane.



At each vertex, the interior angle plus the exterior angle is exactly π . So a consequence of this formula is

$$\begin{aligned} (\text{sum of interior angles of } P) &= n\pi - (\text{sum of exterior angles of } P) \\ &= (n - 2)\pi. \end{aligned}$$

Example 4.21. Let's check the Gauss–Bonnet formula for triangles on spheres. Let S^2 be the sphere of radius 1, and let P be the triangle in the diagram below.



We will compute each of the terms in the formula.

1. (The $\int_P K(x) dx$ term) The normal curvatures at every point of the sphere are 1, since the osculating circle of every cross-section is of radius exactly 1. Hence $K(x) = 1$ for every x . In particular,

$$\int_P K(x) dx = \int_P 1 dx = \text{area}(P).$$

What is the area of P ? Well, 8 copies of P would cover up the entire surface of the sphere, so the area of P must be 1/8-th of the surface area of the sphere, which is 4π . Hence $\int_P K(x) dx = \pi/2$.

2. (The (sum of exterior angles of P) term) Clearly the sum of exterior angles of P is $3\pi/2$.
3. (The Euler characteristic of P) The triangle has 3 vertices, 3 edges, and 1 face, giving $\chi(P) = 3 - 3 + 1 = 1$. Hence $2\pi\chi(P) = 2\pi$.

Since $\pi/2 + 3\pi/2 = 2\pi$, the Gauss–Bonnet formula works in this case!

In fact, this example shows we can use the Gauss–Bonnet formula to calculate areas of triangles on the sphere S^2 : we know $2\pi\chi(P) = 2\pi$ for any triangle P , so as long as we know the sum of exterior angles of the triangle, we get

$$\text{area}(P) = \int_P 1 dx = \int_P K(x) dx = 2\pi - (\text{sum of exterior angles of } P).$$

Theorem 4.22 (Global Gauss–Bonnet formula). *Let S be a (compact) orientable surface, and $K(x)$ be the Gaussian curvature at the point x in S . Then*

$$\int_S K(x) dx = 2\pi\chi(S).$$

Note that this global Gauss–Bonnet formula isn’t anything new: just imagine taking the polygon P to be the entire surface S , so that there are no exterior angles to sum. The result of the local Gauss–Bonnet formula in this case is precisely the global Gauss–Bonnet formula.

Example 4.23. Let S^2 be the sphere of radius r . Then $K(x) = 1/r^2$, so the Gauss–Bonnet formula simplifies to

$$\int_S \frac{1}{r^2} dx = \frac{1}{r^2} \text{area}(S) = 2\pi(2) = 4\pi.$$

Indeed, the surface area of a sphere of radius r is $4\pi r^2$.

Example 4.24. Since the torus T^2 has Euler characteristic 0, the global Gauss–Bonnet formula says $\int_{T^2} K(x) dx = 0$.