

# Computation of the 1-loop diagrams

## Propagators

We shall work with  $\xi = 1$

$$\overbrace{A_{\mu a}(x) A_{\nu b}(y)} = \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \left( \begin{array}{c} p \\ \leftarrow \\ \text{wavy line} \\ \mu a \quad \nu b \end{array} \right) = e^2 \delta_{ab} \left( \frac{\delta_{\mu\nu}}{p^2} + (\xi - 1) \frac{p_\mu p_\nu}{p^4} \right)$$

$$\overbrace{\Psi(x) \bar{\Psi}(y)} = \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \left( \begin{array}{c} p \\ \leftarrow \\ \text{solid line} \end{array} \right) = \frac{1}{\not{p} + m_f} = \frac{\not{p} + m_f}{p^2 + m_f^2}$$

$$\overbrace{C(x) \bar{C}(y)} = \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \left( \begin{array}{c} p \\ \leftarrow \\ \text{dashed line} \end{array} \right) = \frac{1}{-p^2}$$

$$\overbrace{\Phi(x) \bar{\Phi}(y)} = \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \left( \begin{array}{c} p \\ \leftarrow \\ \text{dashed line} \end{array} \right) = \frac{1}{p^2 + m_b^2}$$

## Vertices

$$-S_{E, \text{int}} = \int d^4 x \left\{ -\frac{1}{2e^2} (\partial^\mu A^\nu - \partial^\nu A^\mu) [A_\mu, A_\nu] \right. \quad \left. \begin{array}{l} \text{wavy line} \\ \text{crossing} \end{array} \right.$$

$$-\frac{1}{4e^2} [A^\mu, A^\nu] \cdot [A_\mu, A_\nu] \quad \begin{array}{l} \text{wavy line} \\ \text{crossing} \end{array}$$


$$+ i \bar{\Psi} \not{A} \Psi \quad \begin{array}{l} \text{solid line} \\ \text{wavy line} \end{array}$$

$$+ \partial^\mu \bar{C} \cdot [A_\mu, C] \quad \begin{array}{l} \text{dashed line} \\ \text{wavy line} \end{array}$$

$$+ \bar{\Phi} A^\mu \partial_\mu \Phi - \partial^\mu \bar{\Phi} A_\mu \Phi \quad \begin{array}{l} \text{dashed line} \\ \text{wavy line} \end{array}$$

$$+ \bar{\Phi} A^\mu A_\mu \Phi \quad \begin{array}{l} \text{dashed line} \\ \text{wavy line} \end{array} \quad \left. \right\}$$

• Contributions to  $\langle A_{\mu_0 a_0}(x) A_{\nu_0 b_0}(y) \rangle$



$$= A_{\mu_0 a_0}(x) \frac{1}{2} \int d^4 z_1 d^4 z_2 \overbrace{i \bar{\psi} \gamma^\mu \psi(z_1) i \bar{\psi} \gamma^\nu \psi(z_2)}^{z_1 \leftrightarrow z_2} A_{\nu_0 b_0}(y) \times 2$$

↓


↑ move w (-1)


$$= i^2 (-1) \int d^4 z_1 d^4 z_2 \underbrace{A_{\mu_0 a_0}(x) A_{\mu a}(z_1)}_{\int \frac{d^4 p_1}{(2\pi)^4} e^{-i p_1(x-z_1)} \leftarrow_{\mu_0 a_0} p_1} \text{tr}_{V_f \otimes S} \left( \gamma^\mu e^a \underbrace{\psi(z_1) \bar{\psi}(z_2)}_{\int \frac{d^4 k_1}{(2\pi)^4} e^{-i k_1(z_1-z_2)} \leftarrow_{k_1}} \gamma^\nu e^b \underbrace{\psi(z_2) \bar{\psi}(z_1)}_{\int \frac{d^4 k_2}{(2\pi)^4} e^{-i k_2(z_2-z_1)} \leftarrow_{k_2}} \right) \underbrace{A_{\nu_0 b}(z_2) A_{\nu_0 b_0}(y)}_{\int \frac{d^4 p_2}{(2\pi)^4} e^{-i p_2(z_2-y)} \leftarrow_{\nu_0 b_0} p_2}$$

$$\int d^4 z_1 d^4 z_2 \sim (2\pi)^4 \delta(p_1 - k_1 + k_2) \cdot (2\pi)^4 \delta(k_1 - k_2 - p_2)$$

$$= (2\pi)^8 \delta(p_1 - p_2) \delta(k_2 - (k_1 - p_1)) \sim \begin{matrix} p_1 = p_2 = p \\ k_1 = k, k_2 = k - p \end{matrix}$$

$$= \int \frac{d^4 p}{(2\pi)^4} e^{-i p x} \leftarrow_{\mu_0 a_0} p \left[ \int \frac{d^4 k}{(2\pi)^4} \text{tr}_{V_f \otimes S} \left( \gamma^\mu e^a \leftarrow_{k} \gamma^\nu e^b \leftarrow_{k-p} \right) \right] \leftarrow_{\nu_0 b_0} p e^{i p y}$$





$z_1 \leftrightarrow z_2$

$$= A_{\mu_0 a_0}(x) \frac{1}{2} \int d^4 z_1 d^4 z_2 \overbrace{\partial^\mu \bar{C} \cdot [A_\mu, C](z_1)} \overbrace{\partial^\nu \bar{C} \cdot [A_\nu, C](z_2)} \overbrace{A_{\nu b_0}(y)} \times 2$$

move  $\leftrightarrow (-1)$


$$= (-1) \int d^4 z_1 d^4 z_2 \underbrace{A_{\mu_0 a_0}(x) A_{\mu a}(z_1)}$$


$$\text{tr}_g \left( \underbrace{\text{ad} e^a C(z_1) \partial^\nu \bar{C}(z_2)} \underbrace{\text{ad} e^b C(z_2) \partial^\mu \bar{C}(z_1)} \right) \underbrace{A_{\nu b}(z_2) A_{\nu b_0}(y)}$$

$$\int \frac{d^4 k_1}{(2\pi)^4} e^{-ik_1(z_1-z_2)} \quad \begin{matrix} k_1 \\ \leftarrow \leftarrow \leftarrow ik_1^\nu \end{matrix} \quad \int \frac{d^4 k_2}{(2\pi)^4} e^{-ik_2(z_2-z_1)} \quad \begin{matrix} k_2 \\ \leftarrow \leftarrow \leftarrow ik_2^\mu \end{matrix}$$

$$= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \underbrace{\mu_0 a_0}_{\mu a}$$

$$(-1) \int \frac{d^4 k}{(2\pi)^4} \text{tr}_g \left( \text{ad} e^a \underbrace{\leftarrow \leftarrow \leftarrow k}_{} \text{ad} e^b \underbrace{\leftarrow \leftarrow \leftarrow k-p}_{} i(k-p)^\mu \right) \underbrace{\mu_b}_{\nu b_0} e^{ip y}$$





$$= A_{\mu_0 a_0}(x) \frac{1}{2} \int d^4 z_1 d^4 z_2 \underbrace{V_3^B(z_1) V_3^B(z_2)}_{z_1 \leftrightarrow z_2} A_{\nu_0 b_0}(y) \times 2$$

$$V_3^B = \bar{\phi} A^\mu \partial_\mu \phi - \partial^\mu \bar{\phi} A_\mu \phi = \bar{\phi} (\vec{\partial}^\mu - \vec{\partial}^\mu) e^a \phi \cdot A_{\mu a}$$

$$= \int d^4 z_1 d^4 z_2 \overbrace{A_{\mu_0 a_0}(x) A_{\mu a}(z_1)}$$

$$\underbrace{\bar{\phi}(z_1) (\vec{\partial}^\mu - \vec{\partial}^\mu) e^a \phi(z_1) \bar{\phi}(z_2) (\vec{\partial}^\nu - \vec{\partial}^\nu) e^b \phi(z_2)}_{\text{move}} \overbrace{A_{\nu b}(z_2) A_{\nu_0 b_0}(y)}$$

$$= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \underbrace{\mu_0 a_0}_{\mu a}$$

$$\int \frac{d^4 k}{(2\pi)^4} \text{tr}_{V_b} \left( (-ik^\mu - i(k-p)^\mu) e^a \underbrace{-\frac{k}{\epsilon}}_{\nu b} (-i(k-p)^\nu - ik^\nu) e^b \right) \underbrace{e^{ipy}}_{\nu_0 b_0}$$





$$= A_{\mu_0 \alpha_0}(x) \int d^4 z \overbrace{\bar{\phi} A_\mu \delta^{\mu\nu} A_\nu \phi(z)} \overbrace{A_{\nu_0 \beta_0}(y)} \times 2$$

$A_\mu \leftrightarrow A_\nu$


(red arrow labeled "move" points from the  $\phi(z)$  term to the  $A_{\nu_0 \beta_0}(y)$  term)

$$= 2 \int d^4 z A_{\mu_0 \alpha_0}(x) A_{\mu\alpha}(z_1) \delta^{\mu\nu} \text{tr}_{V_b} (e^a e^b \overbrace{\phi(z) \bar{\phi}(z)}) \overbrace{A_{\nu\beta}(z) A_{\nu_0 \beta_0}(y)}$$

$$= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \overbrace{\mu_0 \alpha_0} \overbrace{\mu\alpha}^p$$

$$2 \int \frac{d^4 k}{(2\pi)^4} \delta^{\mu\nu} \text{tr}_{V_b} (e^a e^b \overbrace{---k---})$$

$$\overbrace{\nu\beta}^p \overbrace{\nu_0 \beta_0} e^{ipy}$$





3-3-2 terms

$$= A_{\mu\alpha_0}(x) \frac{1}{2} \int d^4z_1 d^4z_2 \underbrace{V_B^G(z_1) V_B^G(z_2)}_{z_1 \leftrightarrow z_2} A_{\alpha_0\beta_0}(y) \times 2 ;$$

$$V_B^G := -\frac{1}{2e^2} (\partial^\mu A^\nu - \partial^\nu A^\mu) \cdot [A_\mu, A_\nu] = \frac{1}{e^2} \partial^\nu A^\mu \cdot [A_\mu, A_\nu]$$

$$= \frac{1}{2} \int d^4z_1 d^4z_2 A_{\mu\alpha_0}(x) \underbrace{V_3[A, A, A](z_1) V_3[A, A, A](z_2)}_{z_1 \leftrightarrow z_2} A_{\alpha_0\beta_0}(y) ;$$

$$V_3[A_1, A_2, A_3] := \frac{1}{e^2} \partial^\nu A_1^\mu \cdot [A_{2\mu}, A_{3\nu}] + \text{permutations (3! terms)}$$

If we put  $A_I = e^{a_I} dx^\mu e^{ik_I x} A_{\mu\alpha_I}(k_I)$   $I=1,2,3$ , then

$$V_3[A_1, A_2, A_3]$$

$$= \frac{1}{e^2} ik_1^\nu e^{a_1} A_{\alpha_1}^\mu(k_1) \cdot [e^{a_2} A_{\mu\alpha_2}(k_2), e^{a_3} A_{\nu\alpha_3}(k_3)] e^{i(k_1+k_2+k_3)x} + \text{permutations}$$

$$= \frac{i}{e^2} e^{a_1} \cdot [e^{a_2}, e^{a_3}] k_1^\nu A_{\alpha_1}^\mu(k_1) A_{\mu\alpha_2}(k_2) A_{\nu\alpha_3}(k_3) e^{i(k_1+k_2+k_3)x} + \text{permutations}$$

$$= V_3 \begin{matrix} \mu_1 \alpha_1 & \mu_2 \alpha_2 & \mu_3 \alpha_3 \\ k_1, & k_2, & k_3 \end{matrix} A_{\mu_1 \alpha_1}(k_1) A_{\mu_2 \alpha_2}(k_2) A_{\mu_3 \alpha_3}(k_3) e^{i(k_1+k_2+k_3)x} ;$$

$$\bigvee_{\substack{M_1 a_1, M_2 a_2, M_3 a_3 \\ k_1, k_2, k_3}} := \frac{i}{e^2} e^{a_1} \cdot [e^{a_2}, e^{a_3}] \delta^{M_1 M_2} k_1^{M_3} + \text{permutations}$$

Note: The inner product  $\cdot$  in  $\mathfrak{g}$  is adjoint invariant

$$[X, Y] \cdot Z + Y \cdot [X, Z] = 0$$

and we are assuming  $\{e^a\} \subset \mathfrak{g}$  is an orthonormal basis,

$$e^a \cdot e^b = \delta^{ab}, \quad \text{Suppose } [e^a, e^b] = \sum_c e^c \underbrace{f^{cab}}$$

the structure constant.

$$\text{Then, } e^a \cdot [e^b, e^c] = f^{abc}$$

It is antisymmetric in  $b \leftrightarrow c$  exchange,  $f^{abc} = -f^{acb}$ .

By the adjoint invariance of  $\cdot$ , we have

$$[e^b, e^a] \cdot e^c + e^a \cdot [e^b, e^c] = 0. \quad \text{i.e. } f^{cba} + f^{abc} = 0.$$

$$\Rightarrow f^{bac} = -f^{bca} = f^{acb} = -f^{abc}$$

I.e.  $f^{abc}$  is totally antisymmetric in exchanges of  $a, b, c$ .

$\Rightarrow$  it is also cyclic invariant,  $f^{abc} = f^{bca} = f^{cab}$ .


Using this property of  $f^{a_1 a_2 a_3} = e^{a_1} \cdot [e^{a_2}, e^{a_3}]$ , we find

$$\begin{aligned} & \bigvee_{\substack{M_1 a_1, M_2 a_2, M_3 a_3 \\ k_1, k_2, k_3}} \\ &= \frac{i}{e^2} f^{a_1 a_2 a_3} \left\{ \delta^{M_1 M_2} (k_1 - k_2)^{M_3} + \delta^{M_2 M_3} (k_2 - k_3)^{M_1} + \delta^{M_3 M_1} (k_3 - k_1)^{M_2} \right\} \end{aligned}$$

Back to 


$$= \frac{1}{2} \int d^4 z_1, d^4 z_2 \overbrace{A_{\mu_1 a_1}(x)} \overbrace{V_3[A, A, A](z_1)} \overbrace{V_3[A, A, A](z_2)} \overbrace{A_{\nu_2 b_2}(y)}$$

$$= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \overbrace{\mu_1 a_1}^p \times \overbrace{\nu_2 b_2}^p e^{ipy} \times$$

$$\frac{1}{2} \int \frac{d^4 k}{(2\pi)^4} V \begin{matrix} \mu_1 a_1 & \mu_2 a_2 & \mu_3 a_3 \\ p, & -k, & k-p \end{matrix} \begin{matrix} \mu_2 a_2 & \nu_2 b_2 \\ \mu_3 a_3 & \nu_3 b_3 \end{matrix} V \begin{matrix} \nu_2 b_2 & \nu_3 b_3 & \nu b \\ k, & -(k-p), & -p \end{matrix}$$








$$= A_{\mu_0 a_0}(x) \int d^4 z \underbrace{V_4^G(z)}_{\boxed{}} \underbrace{A_{\nu_0 b_0}(y)}_{\boxed{}} \quad (4 \cdot 3 \text{ terms}) ;$$

$$V_4^G := -\frac{1}{4e^2} [A^\mu, A^\nu] \cdot [A_\mu, A_\nu]$$

$$V_4[A, A, A, A]$$

$$= \int d^4 z \underbrace{A_{\mu_0 a_0}(x)}_{\boxed{}} \underbrace{V_4[A, A, A, A](z)}_{\boxed{}} \underbrace{A_{\nu_0 b_0}(y)}_{\boxed{}} \times \frac{1}{2} ;$$

$$V_4[A_1, A_2, A_3, A_4] := -\frac{1}{4e^2} [A_1^\mu, A_2^\nu] \cdot [A_{3\mu}, A_{4\nu}] + \text{permutations}$$

$$(4! = 24 \text{ terms})$$

If we put  $A_I = e^a dx^\mu e^{ik_I x} A_\mu(k_I)$   $I=1, 2, 3, 4$ , then

$$V_4[A_1, A_2, A_3, A_4]$$

$$= -\frac{1}{4e^2} (e^{a_1}, e^{a_2}) \cdot (e^{a_3}, e^{a_4}) A_{a_1}^\mu(k_1) A_{a_2}^\nu(k_2) A_{\mu a_3}(k_3) A_{\nu a_4}(k_4) e^{i(k_1 + \dots + k_4)x} \\ + \text{permutations}$$

$$= -\frac{1}{4e^2} (e^{a_1}, e^{a_2}) \cdot (e^{a_3}, e^{a_4}) \delta^{\mu_1 \mu_3} \delta^{\mu_2 \mu_4} \\ \cdot A_{\mu_1 a_1}(k_1) \dots A_{\mu_4 a_4}(k_4) e^{i(k_1 + \dots + k_4)x}$$


$$+ \text{permutations}$$

(12) ↔ (34) ↗ ↘  
 ↗ ↘  
 ↗ ↘

$$\begin{aligned}
 &= -\frac{1}{4e^2} A_{\mu_1 a_1}(h_1) \dots A_{\mu_4 a_4}(h_4) e^{i(h_1 + \dots + h_4)z} \times 2 \times 2 \\
 &\quad \left\{ [e^{a_1}, e^{a_2}] [e^{a_3}, e^{a_4}] (\delta^{\mu_1 \mu_3} \delta^{\mu_2 \mu_4} - \delta^{\mu_1 \mu_4} \delta^{\mu_2 \mu_3}) \right. \\
 &\quad + [e^{a_1}, e^{a_3}] [e^{a_2}, e^{a_4}] (\delta^{\mu_1 \mu_2} \delta^{\mu_3 \mu_4} - \delta^{\mu_1 \mu_4} \delta^{\mu_3 \mu_2}) \\
 &\quad \left. + [e^{a_1}, e^{a_4}] [e^{a_3}, a_2] (\delta^{\mu_1 \mu_3} \delta^{\mu_4 \mu_2} - \delta^{\mu_1 \mu_2} \delta^{\mu_4 \mu_2}) \right\} \\
 &= V^{\mu_1 a_1, \dots, \mu_4 a_4} A_{\mu_1 a_1}(h_1) \dots A_{\mu_4 a_4}(h_4) e^{i(h_1 + \dots + h_4)z} ;
 \end{aligned}$$


$V^{\mu_1 a_1, \dots, \mu_4 a_4}$


$$\begin{aligned}
 &:= -\frac{1}{e^2} \left\{ \sum_b f^{b a_1 a_2} f^{b a_3 a_4} (\delta^{\mu_1 \mu_3} \delta^{\mu_2 \mu_4} - \delta^{\mu_1 \mu_4} \delta^{\mu_2 \mu_3}) \right. \\
 &\quad + \sum_b f^{b a_1 a_3} f^{b a_2 a_4} (\delta^{\mu_1 \mu_2} \delta^{\mu_3 \mu_4} - \delta^{\mu_1 \mu_4} \delta^{\mu_3 \mu_2}) \\
 &\quad \left. + \sum_b f^{b a_1 a_4} f^{b a_3 a_2} (\delta^{\mu_1 \mu_3} \delta^{\mu_4 \mu_2} - \delta^{\mu_1 \mu_2} \delta^{\mu_4 \mu_2}) \right\}
 \end{aligned}$$



$$= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \begin{array}{c} \leftarrow p \\ \text{wavy line} \\ \mu_1 a_1 \quad \mu_2 a_2 \end{array} \cdot \begin{array}{c} \leftarrow p \\ \text{wavy line} \\ \mu_3 a_3 \quad \mu_4 a_4 \end{array} e^{ip y}$$

$$\times \frac{1}{2} \int \frac{d^4 k}{(2\pi)^4} V^{\mu_1 a_1, \mu_2 a_2, \mu_3 a_3, \mu_4 a_4} \begin{array}{c} \leftarrow k \\ \text{wavy line} \\ \mu_2 a_2 \quad \mu_3 a_3 \end{array}$$




• Contribution to  $\langle \psi(x) \bar{\psi}(y) \rangle$  of 

$$= \psi(x) \frac{1}{2} \int d^4z_1 d^4z_2 \underbrace{i\bar{\psi} A \psi(z_1)}_{\substack{\uparrow \\ z_1 \leftrightarrow z_2}} \underbrace{i\bar{\psi} A \psi(z_2)}_{\substack{\uparrow \\ z_1 \leftrightarrow z_2}} \bar{\psi}(y) \times 2$$

$$= \int \frac{d^4p}{(2\pi)^4} e^{-ipx} \left[ \int \frac{d^4k}{(2\pi)^4} i\gamma^\mu e^a \overset{h-p}{\curvearrowright} i\gamma^\nu e^b \right] e^{ipy}$$




• Contribution to  $\langle C(x) \bar{C}(y) \rangle$  of 

$$= C(x) \frac{1}{2} \int d^4z_1 d^4z_2 \partial^\mu \bar{C} [A_\mu, C](z_1) \partial^\nu \bar{C} [A_\nu, C](z_2) \bar{C}(y) \times 2$$

$$= \int \frac{d^4p}{(2\pi)^4} e^{-ipx} \left[ \int \frac{d^4k}{(2\pi)^4} ip^\mu a d e^a \overset{h-p}{\curvearrowright} ik^\nu a d e^b \right] e^{ipy}$$



• Contributions to  $\langle C(x) A_{\mu_0 a_0}(w) \bar{C}(y) \rangle$

 (tree diagram)


$$= C(x) \int d^4 z \partial^\mu \bar{C} [A_\mu, C](z) A_{\mu_0 a_0}(w) \bar{C}(y)$$


$$= \int d^4 z \underbrace{C(x) \partial^\mu \bar{C}(z)}_{ad} e^a \underbrace{A_{\mu a}(z) A_{\mu_0 a_0}(w)}_{\mu a} \underbrace{C(z) \bar{C}(y)}_{P_2}$$

$$\int \frac{d^4 p_1}{(2\pi)^4} e^{-i p_1(x-z)} \underbrace{p_1}_{\dots \leftarrow \dots} i p_1^\mu \int \frac{d^4 q}{(2\pi)^4} e^{-i q(z-w)} \underbrace{q}_{\mu a} \int \frac{d^4 p_2}{(2\pi)^4} e^{-i p_2(z-y)} \underbrace{p_2}_{\dots \leftarrow \dots}$$

$$\int d^4 z \rightarrow (2\pi)^4 \delta(p_1 - q - p_2) \rightsquigarrow p_2 = p, p_1 = p + q$$

$$= \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} e^{-i(p+q)x} \underbrace{p+q}_{\dots \leftarrow \dots} \boxed{i(p+q)^\mu ad e^a} \dots \leftarrow \dots e^{iqy}$$






$$= \frac{1}{3!} \int \prod_{i=1}^3 d^4 z_i$$


$$c(x) \delta^{\mu_1} \bar{c} [A_{\mu_1}, c](y) \delta^{\mu_2} \bar{c} [A_{\mu_2}, c](z) \delta^{\mu_3} \bar{c} [A_{\mu_3}, c](w) A_{\mu_0 a_0}(y) \bar{c}(y)$$

$\times 3!$   
 ↗ permutation of  $z_1, z_2, z_3$

$$= \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} e^{-i(p+q)x} e^{i q w} \dots$$

$$\int \frac{d^4 k}{(2\pi)^4} i(p+q)^{\mu_1} a_1 e^{i a_1 (k+q)} \dots i(k+q)^{\mu} a e^{i a q} \dots i k^{\mu_3} a_3 e^{i a_3 k} \dots e^{i p y}$$





$$= \frac{1}{2!} \int d^4z_1 d^4z_2 d^4z_3$$



$$\underbrace{C(x)} \underbrace{\partial^{M_1} \bar{C} \cdot [A_{\mu_1}, C]}_{(1)} \underbrace{V_3[A, A, A]}_{(3)} \underbrace{\partial^{M_2} \bar{C} \cdot [A_{\mu_2}, C]}_{(2)} \underbrace{A_{\mu_3 a_3}(w)}_{(4)} \underbrace{\bar{C}(y)}$$

$\times 2 \leftarrow z_1 \leftrightarrow z_2$

$$= \int \frac{d^4p}{(2\pi)^4} \frac{d^4q}{(2\pi)^4} e^{-i(p+q)x} e^{iqw}$$

$\xrightarrow{p+q} \dots \leftarrow 0$

$\mu_3 a_3$   
 $q \downarrow$   
 $\mu a$



## Some useful facts on simple Lie algebra $\mathfrak{g}$

$\{e^a\} \subset \mathfrak{g}$  as before  $\left( \begin{array}{l} \text{an orthonormal basis wrt an adjoint inv} \\ \text{inner product } e^a \cdot e^b = \delta^{ab}, [e^a, e^b] = \sum_c e^c f^{cab} \\ \rightarrow f^{abc} \text{ totally antisymmetric} \end{array} \right)$

For any representation  $V$  of  $\mathfrak{g}$

$$\text{tr}_V(e^a e^b) = -T_V \delta^{ab}, \quad T_V \in \mathbb{R}_{\geq 0}$$

$$\sum_a e^a e^a = -C_2(V): \quad \text{a scalar (quadratic Casimir)} \\ \text{on each irreducible component}$$

• If  $V$  is irreducible,  $T_V \dim \mathfrak{g} = C_2(V) \dim V$

•  $V = \mathfrak{g}$  (adjoint rep),  $T_{\mathfrak{g}} = C_2(\mathfrak{g}) =: h^\vee$  is called the

dual Coxeter number of  $\mathfrak{g}$  for a suitable normalization of "•".

$$\begin{aligned} \cdot \sum_b e^b e^a e^b &= \underbrace{\sum_b e^b [e^a, e^b]}_{\text{red}} + \underbrace{\sum_b e^b e^b e^a}_{\text{blue}} \stackrel{-C_2(V)}{=} -f^{bdc} \\ &= \sum_{bc} e^b e^c f^{cab} = \frac{1}{2} \sum_{b,c} [e^b, e^c] f^{cab} = \frac{1}{2} \sum_{b,c,d} e^d \overbrace{f^{dbc}}^{= -f^{bdc}} f^{cab} \\ &= -\frac{1}{2} \sum_d e^d \underbrace{\text{tr}_{\mathfrak{g}}(\text{ad } e^d \text{ ad } e^a)}_{\text{green}} = \frac{1}{2} h^\vee e^a \\ &= \left( \frac{h^\vee}{2} - C_2(V) \right) e^a \stackrel{\text{green}}{=} -T_{\mathfrak{g}} \delta^{da} = -h^\vee \delta^{da} \end{aligned}$$

$$\stackrel{V=\mathfrak{g}}{\rightsquigarrow} \sum_b c^d e^b \text{ad } e^a \text{ad } e^b = -\frac{h^\vee}{2} \text{ad } e^a$$

We need to evaluate the integrals of the form

$$I(f) = \int \frac{d^4 k}{(2\pi)^4} \frac{f(k)}{(k^2+m^2)((k-p)^2+\mu^2)},$$

$$J(g) = \int \frac{d^4 k}{(2\pi)^4} \frac{g(k)}{(k+q)^2 k^2 (k-p)^2},$$

polynomials of  $k^\mu$ 's

which are often divergent. We shall employ the dimensional regularization in which these are replaced by

$$I_{DR}(f) = M_{DR}^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{f(k)}{(k^2+m^2)((k-p)^2+\mu^2)},$$

$$J_{DR}(g) = M_{DR}^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{g(k)}{(k+q)^2 k^2 (k-p)^2},$$

with  $d = 4 - \epsilon$  (eventually we take  $\epsilon \rightarrow 0$ ). We use

$$\frac{1}{AB} = \int_0^1 \frac{dx}{((1-x)A + xB)^2}$$

$$\frac{1}{ABC} = \int_{\triangle} \frac{2 dy dz}{((1-y-z)A + yB + zC)^3}$$

where  $\triangle := \{ (y, z) \in \mathbb{R}^2 \mid y \geq 0, z \geq 0, y+z \leq 1 \}$



Computation goes as follows

$$\begin{aligned}
 \Gamma_{\text{DR}}(f) &= M_{\text{DR}}^{4-d} \int \frac{d^d k}{(2\pi)^d} \int_0^1 \frac{dx f(k)}{\left( \underbrace{(1-x)(k^2+m^2) + x(k-p)^2 + \mu^2}_{k^2 - 2xpk + xp^2 + (1-x)m^2 + x\mu^2} \right)^2} \\
 &= \int_0^1 dx M_{\text{DR}}^{4-d} \int \frac{d^d \ell}{(2\pi)^d} \frac{f(\ell+xp)}{(\ell^2 + \Delta)^2}
 \end{aligned}$$

- We expand  $f(\ell+xp)$  in  $\ell^m$ 's, drop odd power terms and replace even power terms by a function of  $\ell^2$

$$f(\ell+xp) \rightarrow \tilde{f}(\ell^2, xp). \quad (\text{e.g. } \ell^\mu \ell^\nu \rightarrow \frac{1}{d} \delta^{\mu\nu} \ell^2)$$

- Use  $\int \frac{d^d \ell}{(2\pi)^d} F(\ell^2) = \frac{\text{Vol}(S^{d-1})}{(2\pi)^d} \int_0^\infty \ell^{d-1} d\ell F(\ell^2)$

$$= \frac{\text{Vol}(S^{d-1})}{2(2\pi)^d} \int_0^\infty \ell^{d-2} d\ell^2 F(\ell^2)$$

$$= \frac{1}{(4\pi)^{d/2} \Gamma(d/2)} \int_0^\infty t^{\frac{d}{2}-1} dt F(t)$$

$$\mathcal{I}_{DR}(f) = \frac{M_{DR}^{4-d}}{(4\pi)^{d/2} \Gamma(d/2)} \int_0^1 dx \int_0^\infty \frac{t^{\frac{d}{2}-1} dt \tilde{f}(t, xP)}{(t+\Delta)^2}$$

$$\text{We may use } \int_0^\infty \frac{t^{p-1} dt}{(t+\Delta)^{p+q}} = \frac{B(p, q)}{\Delta^q} = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)\Delta^q}$$

In this way we find

$$\mathcal{I}_{DR}(f) = \frac{1}{(4\pi)^2} \int_0^1 dx \left( \frac{4\pi M_{DR}^2}{\Delta} \right)^{2-\frac{d}{2}} \Gamma(2-\frac{d}{2}) \hat{f}$$

where  $\Delta = x(1-x)p^2 + (1-x)m^2 + x\mu^2$  and

$$\hat{1} = 1, \quad \hat{k}^m = x p^m$$

$$\widehat{k^m k^\nu} = \delta^{m\nu} \frac{\Delta}{2-d} + x^2 p^m p^\nu$$

$$\widehat{k^m (k-p)^\nu} = \delta^{m\nu} \frac{\Delta}{2-d} - x(1-x) p^m p^\nu$$

$$\widehat{(k-p)^m (k-p)^\nu} = \delta^{m\nu} \frac{\Delta}{2-d} + (1-x)^2 p^m p^\nu$$

Similarly (exercise):

$$J_{DR}(g) = \frac{1}{(4\pi)^2} \int_{\triangle} dy dz \left( \frac{4\pi M_{DR}^2}{\Delta} \right)^{2-\frac{d}{2}} \frac{\Gamma(3-\frac{d}{2})}{\Delta} \hat{g}$$

where  $\Delta = y(1-y)q^2 + z(1-z)p^2 + 2yzqp$  and

$$\hat{1} = 1, \quad \hat{k}^m = -yq^m + zp^m$$

$$\hat{k}^m \hat{k}^\nu = \delta^{\mu\nu} \frac{\Delta}{4-d} + (-yq^m + zp^m)(-yq^\nu + zp^\nu)$$

Note:  $\Gamma(3-\frac{d}{2}) = (2-\frac{d}{2})\Gamma(2-\frac{d}{2}) = \frac{1}{2}(4-d)\Gamma(2-\frac{d}{2})$

$$\therefore J_{DR}(g) = \frac{1}{(4\pi)^2} \int_{\triangle} dy dz \left( \frac{4\pi M_{DR}^2}{\Delta} \right)^{2-\frac{d}{2}} \Gamma(2-\frac{d}{2}) \hat{\hat{g}}$$

$$\hat{\hat{1}} = \frac{4-d}{2\Delta}, \quad \hat{\hat{k}}^m = \frac{4-d}{2\Delta} (-yq^m + zp^m)$$

$$\hat{\hat{k}}^m \hat{\hat{k}}^\nu = \frac{1}{2} \delta^{\mu\nu} + \frac{4-d}{2\Delta} (-yq^m + zp^m)(-yq^\nu + zp^\nu)$$