

$$\langle \psi(x) \bar{\psi}(y) \rangle_{\text{conn}} = \overbrace{\psi(x) \bar{\psi}(y)}$$

$$+ \frac{1}{2} (-e)^2 \int d^4 z_1 d^4 z_2 \langle \psi(x) (\bar{\psi} A \psi)(z_1) (\bar{\psi} A \psi)(z_2) \bar{\psi}(y) \rangle_{\text{free}} + \dots$$

⊛

$$\textcircled{\star} = \frac{e^2}{2} \int d^4 z_1 d^4 z_2 \psi(x) \overbrace{(\bar{\psi} A \psi)(z_1)} \overbrace{(\bar{\psi} A \psi)(z_2)} \bar{\psi}(y) \times 2$$

$$= e^2 \int d^4 z_1 d^4 z_2 \underbrace{\psi(x) \bar{\psi}(z_1)} \underbrace{\gamma^\mu \psi(z_1) \bar{\psi}(z_2)} \underbrace{\gamma^\nu \psi(z_2) \bar{\psi}(y)} \cdot \underbrace{A_\mu(z_1) A_\nu(z_2)}$$

$$\int \frac{d^4 p_1}{(2\pi)^4} \frac{e^{-i p_1 (x - z_1)}}{-\not{p}_1 + m} \int \frac{d^4 k_1}{(2\pi)^4} \frac{e^{-i k_1 (z_1 - z_2)}}{-\not{k}_1 + m} \int \frac{d^4 p_2}{(2\pi)^4} \frac{e^{-i p_2 (z_2 - y)}}{-\not{p}_2 + m} \int \frac{d^4 k_2}{(2\pi)^4} \frac{\delta_{\mu\nu} e^{-i k_2 (z_1 - z_2)}}{k_2^2}$$

$$(2\pi)^4 \delta(p_1 - k_1 - k_2) (2\pi)^4 \delta(k_1 - p_2 + k_2)$$

$$= (2\pi)^8 \delta(p_1 - p_2) \delta(k_2 + k_1 - p_2)$$

$$= \int \frac{d^4 p}{(2\pi)^4} \frac{e^{-i p x}}{-\not{p} + m} \underbrace{e^2 \int \frac{d^4 k}{(2\pi)^4} \gamma^\mu \frac{1}{-\not{k} + m} \gamma^\nu \frac{\delta_{\mu\nu}}{(k-p)^2} \frac{e^{i p y}}{-\not{p} + m}}_{=: -\Sigma_2(p)}$$



$$\Sigma_2(p) = -e^2 \int \frac{d^4 k}{(2\pi)^4} \gamma^\mu \frac{1}{-k+m} \gamma^\nu \frac{\delta_{\mu\nu}}{(k-p)^2}$$

$$= -e^2 \int \frac{d^4 k}{(2\pi)^4} \frac{\gamma^\mu (k+m) \gamma_\mu}{(k^2+m^2)(k-p)^2}$$

$$\left[ \begin{array}{l} \gamma^\mu (k+m) \gamma_\mu = 2k - 4m \\ \left( \begin{array}{l} \gamma^\mu \gamma_\mu = -4 \\ \gamma^\mu \gamma^\rho \gamma_\mu = \{\gamma^\mu, \gamma^\rho\} \gamma_\mu - \gamma^\rho \gamma^\mu \gamma_\mu \\ = -2\delta^{\mu\rho} \gamma_\mu + 4\gamma^\rho = 2\gamma^\rho \end{array} \right) \end{array} \right]$$

$$= e^2 \int \frac{d^4 k}{(2\pi)^4} \frac{-2k + 4m}{(k^2+m^2)(k-p)^2}$$

Pauli-Villars regularization

$$\frac{1}{(k-p)^2} \rightarrow \frac{1}{(k-p)^2} - \frac{1}{(k-p)^2 + \Lambda^2}$$

$\Leftrightarrow$  introduce a regulator field  $B_\mu$  with Lagrangian

$$\Delta \mathcal{L} = \frac{1}{2} B^\mu (-\partial^2 + \Lambda^2) B_\mu + ie \bar{\Psi} \not{B} \Psi$$

gauge invariance is preserved with  $\delta B = 0$ .

$$\begin{aligned}
 \frac{1}{(k^2+m^2)((k-p)^2+\Lambda^2)} &= \int_0^1 \frac{dx}{\left(\underbrace{((1-x)(k^2+m^2)+x((k-p)^2+\Lambda^2))}_{k^2-2xpk+xp^2+(1-x)m^2+x\Lambda^2}\right)^2} \\
 &= \underbrace{(k-xp)^2}_2 + \underbrace{x(1-x)p^2+(1-x)m^2+x\Lambda^2}_\Delta
 \end{aligned}$$

$$\text{denominator} = -2(k+xp) + 4m \approx -2xp + 4m$$

$$\Sigma_{2, PV}(p) = \sum_{i=0}^1 C_i \Sigma_2(p, \Lambda_i); \quad (C_0, \Lambda_0) = (1, 0), \quad (C_1, \Lambda_1) = (-1, \Lambda)$$

$$= 2e^2 \int_0^1 dx \int \frac{d^4 \ell}{(2\pi)^4} \sum_i C_i \frac{-x\not{p} + 2m}{(\ell^2 + \Delta_i)^2}$$

$$= \frac{2e^2}{(4\pi)^2} \int_0^1 dx \int_0^\infty t dt \sum_i C_i \frac{-x\not{p} + 2m}{(t + \Delta_i)^2}$$

$$= \frac{2e^2}{(4\pi)^2} \int_0^1 dx \left[ \sum_i C_i \left( \log(t + \Delta_i) + \frac{\Delta_i}{t + \Delta_i} \right) \right]_0^\infty (-x\not{p} + 2m)$$

$$= -\frac{2e^2}{(4\pi)^2} \int_0^1 dx \sum_i C_i (\log \Delta_i + \cancel{X}) (-x\not{p} + 2m)$$

$$= -\frac{2e^2}{(4\pi)^2} \int_0^1 dx \log\left(\frac{\Delta_0}{\Delta_1}\right) (-x\not{p} + 2m)$$

$$\begin{aligned} \log \frac{\Delta_0}{\Delta_1} &= \log \left( \frac{x(1-x)p^2 + (1-x)m^2}{x(1-x)p^2 + (1-x)m^2 + x\Lambda^2} \right) \\ &= \log \left( \frac{x(1-x)p^2 + (1-x)m^2}{x\Lambda^2} \right) + O\left(\frac{p^2}{\Lambda^2}, \frac{m^2}{\Lambda^2}\right) \end{aligned}$$

$$\Sigma_{2, PV}(p) = \frac{2e^2}{(4\pi)^2} \int_0^1 dx (-x\not{p} + 2m) \log \left( \frac{x\Lambda^2}{x(1-x)p^2 + (1-x)m^2} \right)$$

$$\langle \psi(x) \bar{\psi}(y) A_\nu(z) \rangle_{PI}$$

$$= -e \int d^4 w \overbrace{\psi(x) \bar{\psi}(w)} \gamma^\mu \overbrace{A_\mu(w) \psi(w) \bar{\psi}(y)} A_\nu(z) \quad \left. \vphantom{\int} \right\} \leftarrow \leftarrow$$

$$+ \frac{1}{3!} (-e)^3 \int d^4 w_1 d^4 w_2 d^4 w_3 \underbrace{\overbrace{\psi(x) (\bar{\psi} A \psi)(w_1)} \overbrace{(\bar{\psi} A \psi)(w_2)} \overbrace{(\bar{\psi} A \psi)(w_3)} \bar{\psi}(y) A_\nu(z)} \times 3! \quad \left. \vphantom{\int} \right\} \leftarrow \leftarrow \leftarrow$$

+ ...

$$\left. \vphantom{\int} \right\} \leftarrow \leftarrow = -e \int d^4 w \int \frac{d^4 p_1}{(2\pi)^4} \frac{e^{-i p_1(x-w)}}{-\not{p}_1 + m} \gamma^\mu \int \frac{d^4 p_2}{(2\pi)^4} \frac{e^{-i p_2(w-y)}}{-\not{p}_2 + m} \int d^4 q \frac{e^{-i q(w-z)}}{q^2} \delta_{\mu\nu}$$

$$= \int \frac{d^4 p}{(2\pi)^4} \frac{e^{-i(p+q)x}}{-\not{p+q} + m} (-e \gamma^\mu) \frac{e^{i p y}}{-\not{p} + m} \frac{\delta_{\mu\nu} e^{i q z}}{q^2}$$

$$\left. \vphantom{\int} \right\} \leftarrow \leftarrow \leftarrow = (-e)^3 \int d^4 w_1 d^4 w_2 d^4 w_3 \overbrace{\psi(x) \bar{\psi}(w_1) \gamma^\rho A_\rho(w_1) \psi(w_1)} \overbrace{\bar{\psi}(w_2) \gamma^\mu A_\mu(w_2) \psi(w_2)} \overbrace{\bar{\psi}(w_3) \gamma^\lambda A_\lambda(w_3) \psi(w_3)} \bar{\psi}(y) A_\nu(z)$$

$$= (-e)^3 \int d^4 w_1 d^4 w_2 d^4 w_3 \int \frac{d^4 p_1}{(2\pi)^4} \frac{e^{-i p_1 (x-w_1)}}{-\not{p}_1 + m} \gamma^\rho \int \frac{d^4 k_1}{(2\pi)^4} \frac{e^{-i k_1 (w_1-w_2)}}{-\not{k}_1 + m}$$

$$\gamma^\mu \int \frac{d^4 k_2}{(2\pi)^4} \frac{e^{-i k_2 (w_2-w_3)}}{-\not{k}_2 + m} \gamma^\lambda \int \frac{d^4 p_2}{(2\pi)^4} \frac{e^{-i p_2 (w_2-y)}}{-\not{p}_2 + m}$$

$$\int \frac{d^4 k_3}{(2\pi)^4} \frac{e^{-i k_3 (w_1-w_3)}}{k_3^2} \delta_{\rho\lambda} \int \frac{d^4 q}{(2\pi)^4} \frac{e^{-i q (w_2-y)}}{q^2} \delta_{\mu\nu}$$


$$\left[ \int d^4 w_1 d^4 w_2 d^4 w_3 \Rightarrow (2\pi)^4 \delta(p_1 - k_1 - k_3) (2\pi)^4 \delta(k_1 - k_2 - q) (2\pi)^4 \delta(k_2 - p_2 + k_3) \right.$$

$$\left. = (2\pi)^{4 \cdot 3} \delta(p_1 - p_2 - q) \delta(k_1 - k_2 - q) \delta(k_2 - p_2 + k_3) \right]$$

$$= \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} \frac{e^{-i(p+q)x}}{-(p+q) + m} \left( -e \Gamma_2^\mu(p, q) \right) \frac{e^{i p y}}{-\not{p} + m} \frac{e^{i q z} \delta_{\mu\nu}}{q^2},$$

where

$$\Gamma_2^\mu(p, q) = e^2 \int \frac{d^4 k}{(2\pi)^4} \gamma^\rho \frac{1}{-(k+q) + m} \gamma^\mu \frac{1}{-\not{k} + m} \gamma^\lambda \frac{\delta_{\rho\lambda}}{(k-p)^2}$$



$$\Gamma_2^{\mu}(p, q) = e^2 \int \frac{d^4 k}{(2\pi)^4} \frac{\gamma^{\rho} ((k+q)+m) \gamma^{\mu} (k+m) \gamma_{\rho}}{((k+q)^2+m^2)(k^2+m^2)(k-p)^2}$$

$$\gamma^{\rho} \gamma^{\mu} \gamma_{\rho} = \underbrace{\{\gamma^{\rho}, \gamma^{\mu}\}}_{-2\delta^{\rho\mu}} \gamma_{\rho} - \gamma^{\mu} \underbrace{\gamma^{\rho} \gamma_{\rho}}_{-4} = 2\gamma^{\mu}$$


$$\begin{aligned} \gamma^{\rho} \gamma^{\nu} \gamma^{\mu} \gamma_{\rho} &= \underbrace{\{\gamma^{\rho}, \gamma^{\nu}\}}_{-2\delta^{\rho\nu}} \gamma^{\mu} \gamma_{\rho} - \gamma^{\nu} \underbrace{\{\gamma^{\rho}, \gamma^{\mu}\}}_{-2\delta^{\rho\mu}} \gamma_{\rho} + \gamma^{\nu} \gamma^{\mu} \underbrace{\gamma^{\rho} \gamma_{\rho}}_{-4} \\ &= -2\delta^{\rho\nu} \gamma^{\mu} \gamma_{\rho} + 2\delta^{\rho\mu} \gamma^{\nu} \gamma_{\rho} - 4\gamma^{\nu} \gamma^{\mu} \\ &= -2\gamma^{\mu} \gamma^{\nu} - 2\gamma^{\nu} \gamma^{\mu} = 4\delta^{\mu\nu} \end{aligned}$$

$$\begin{aligned} \gamma^{\rho} \gamma^{\nu} \gamma^{\mu} \gamma^{\sigma} \gamma_{\rho} &= \gamma^{\rho} \gamma^{\nu} \gamma^{\mu} \underbrace{\{\gamma_{\rho}, \gamma^{\sigma}\}}_{-2\delta_{\rho}^{\sigma}} - \underbrace{\gamma^{\rho} \gamma^{\nu} \gamma^{\mu} \gamma_{\rho}}_{4\delta^{\mu\nu}} \gamma^{\sigma} \\ &= -2\gamma^{\sigma} \gamma^{\nu} \gamma^{\mu} - 4\delta^{\mu\nu} \gamma^{\sigma} = 2\gamma^{\sigma} \gamma^{\mu} \gamma^{\nu} \end{aligned}$$

$$\begin{aligned} \therefore \text{nemerator} &= 2k \gamma^{\mu} (k+q) + 4m(k+q)^{\mu} + 4m k^{\mu} + 2m^2 \gamma^{\mu} \\ &= 2 \left[ k \gamma^{\mu} (k+q) + 2m(2k+q)^{\mu} + m^2 \gamma^{\mu} \right] \end{aligned}$$

$$\Gamma_2^{\mu}(p, q) = 2e^2 \int \frac{d^4 k}{(2\pi)^2} \frac{k \gamma^{\mu} (k+q) + 2m(2k+q)^{\mu} + m^2 \gamma^{\mu}}{((k+q)^2+m^2)(k^2+m^2)(k-q)^2}$$

In the Pauli-Villars regularization we have been using, there is another 1-loop diagram where the regulator field  $B_\mu$  is involved:



As the coupling is  $ie\bar{\Psi}\not{B}\Psi$ , this will produce  $i^2\Gamma^M(p, q; \Lambda)$

where

$$\Gamma_2^M(p, q; \Lambda) := 2e^2 \int \frac{d^4k}{(2\pi)^4} \frac{k \gamma^M (k+q) + 2m(2k+q)^\mu + m^2 \gamma^M}{((k+q)^2 + m^2)(k^2 + m^2)((k-q)^2 + \Lambda^2)}$$

In total,

$$\begin{aligned} \Gamma_{2, PV}^M(p, q) &= \Gamma_2^M(p, q; 0) - \Gamma_2^M(p, q; \Lambda) \\ &= \sum_{i=0}^1 C_i \Gamma_2^M(p, q; \Lambda_i) \end{aligned}$$

where  $(C_0, \Lambda_0) = (1, 0)$  and  $(C_1, \Lambda_1) = (-1, \Lambda)$ .



Let us evaluate the integral.

$$\begin{aligned}
 & \frac{1}{((k+q)^2+m^2)(k^2+m^2)((k-p)^2+\Lambda_i^2)} \\
 &= 2 \int_0^{\infty} \frac{dx dy dz \delta(1-x-y-z)}{\left[ x(k^2+m^2) + y((k+q)^2+m^2) + z((k-p)^2+\Lambda_i^2) \right]^3} \\
 & \quad k^2 + 2yqh - 2zpk + yq^2 + zp^2 + (x+y)m^2 + z\Lambda_i^2 \\
 &= \underbrace{(k+yq-zp)^2}_{\ell} + \underbrace{y(1-y)q^2 + z(1-z)p^2 + 2yzpq}_{\Delta_i} + (1-z)m^2 + z\Lambda_i^2
 \end{aligned}$$

• numerator

$$\begin{aligned}
 & (\cancel{\ell - yq + zp}) \gamma^m (\cancel{\ell + (1-y)q + zp}) + 2m((1-2y)q + 2zp)^m + m^2 \gamma^m \\
 &= \cancel{\ell} \gamma^m \cancel{\ell} + (-yq + zp) \gamma^m ((1-y)q + zp) + 2m((1-2y)q + 2zp)^m + m^2 \gamma^m \\
 & \quad \text{+ } \ell\text{-linear} \\
 & \left[ \cancel{\ell} \gamma^m \cancel{\ell} \simeq \frac{1}{4} \cancel{\ell}^2 \gamma^\rho \gamma^\sigma \gamma_\rho = \frac{1}{2} \cancel{\ell}^2 \gamma^m \right. \\
 & \simeq \frac{1}{2} \cancel{\ell}^2 \gamma^m + \underbrace{(-yq + zp) \gamma^m ((1-y)q + zp) + 2m((1-2y)q + 2zp)^m + m^2 \gamma^m}_{\substack{!! \\ X^m}}
 \end{aligned}$$

$$\therefore \Gamma_{2PV}^M(p, q) = 4e^2 \int_0^\infty dx dy dz \delta(1-x-y-z) \underbrace{\int \frac{d^4 l}{(2\pi)^4} \sum_i C_i \frac{\frac{1}{2} l^2 \gamma^m + X^m}{(l^2 + \Delta_i)^3}}_{\text{depends only on } y, z}$$

$$= 4e^2 \int_{\triangle} dy dz \int \frac{d^4 l}{(2\pi)^4} \sum_i C_i \frac{\frac{1}{2} l^2 \gamma^m + X^m}{(l^2 + \Delta_i)^3}$$

where  $\triangle := \{ (y, z) \in \mathbb{R}^2 \mid y \geq 0, z \geq 0, y+z \leq 1 \}$

$$\begin{aligned} \cdot \int \frac{d^4 l}{(2\pi)^4} \frac{1}{(l^2 + \Delta)^3} &= \frac{1}{(4\pi)^2} \int_0^\infty \frac{t dt}{(t + \Delta)^3} = \frac{1}{(4\pi)^2} \int_0^\infty d \left( \frac{-1}{t + \Delta} + \frac{1}{2} \frac{\Delta}{(t + \Delta)^2} \right) \\ &= \frac{1}{(4\pi)^2} \left( \frac{1}{\Delta} - \frac{1}{2} \frac{\Delta}{\Delta^2} \right) = \frac{1}{2(4\pi)^2} \frac{1}{\Delta} \end{aligned}$$

$$\begin{aligned} \cdot \int \frac{d^4 l}{(2\pi)^4} \sum_i C_i \frac{l^2}{(l^2 + \Delta_i)^3} &= \frac{1}{(4\pi)^2} \int_0^\infty t dt \underbrace{\sum_i C_i \frac{t}{(t + \Delta_i)^3}}_{d \left( \sum_i C_i \log(t + \Delta_i) + 2 \sum_i \frac{C_i \Delta_i}{t + \Delta_i} - \frac{1}{2} \sum_i \frac{C_i \Delta_i^2}{(t + \Delta_i)^2} \right)} \\ &= \frac{1}{(4\pi)^2} \left( - \sum_i C_i \log \Delta_i \right) = \frac{1}{(4\pi)^2} \log \left( \frac{\Delta_1}{\Delta_0} \right) \end{aligned}$$

$$\therefore \Gamma_{2p\nu}^{\mu}(p, q) = 4e^2 \int_{\triangle} dy dz \left[ \frac{1}{(4\pi)^2} \log\left(\frac{\Delta_1}{\Delta_0}\right) \frac{1}{2} \gamma^{\mu} + \frac{1}{2(4\pi)^2} \left( \frac{X^{\mu}}{\Delta_0} - \frac{X^{\mu}}{\Delta_1} \right) \right]$$

$$= \frac{2e^2}{(4\pi)^2} \int_{\triangle} dy dz \left[ \log\left(\frac{z\Lambda^2}{\Delta_0}\right) \gamma^{\mu} + \frac{X^{\mu}}{\Delta_0} + \mathcal{O}\left(\frac{m^2}{\Lambda^2}, \frac{p^2}{\Lambda^2}, \frac{q^2}{\Lambda^2}, \frac{pq}{\Lambda^2}\right) \right]$$

where

$$\Delta_0 = y(1-y)q^2 + z(1-z)p^2 + 2yzpq + (1-z)m^2$$

$$X^{\mu} = (-y\not{q} + z\not{p})\gamma^{\mu}((1-y)\not{q} + z\not{p}) + 2m((1-2y)\not{q} + 2z\not{p})^{\mu} + m^2\gamma^{\mu}$$

Recall  $\Gamma[\Phi] = S_{\text{free}}[\Phi]$  - generating function of 1PI diagrams.

$$\text{1PI} \text{ (circle with wavy lines)} = \Pi^{\mu\nu}(p)$$

$$\text{1PI} \text{ (circle with straight lines)} = -\Sigma(p)$$

$$\text{1PI} \text{ (circle with wavy line and straight line)} = -e\Gamma^{\mu}(p, q)$$

} divergent 1PI vertices.

$\Gamma[A, \psi, \bar{\psi}]$

$$= \int \frac{d^4 p}{(2\pi)^4} \left\{ \frac{1}{2} A_{\mu}(-p) \left( \delta^{\mu\nu} p^2 - p^{\mu} p^{\nu} + \frac{1}{\xi} p^{\mu} p^{\nu} - \Pi^{\mu\nu}(p) \right) A_{\nu}(p) \right. \\ \left. + \bar{\psi}(-p) (-\not{p} + m + \Sigma(p)) \psi(p) \right.$$

$$\left. + \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} \bar{\psi}(-p-q) e\Gamma^{\mu}(p, q) A_{\mu}(q) \psi(p) \right.$$

+ higher powers of fields

↖ Here we recovered the general gauge parameter  $\xi$

## Digression: Ward identity for 1PI effective action

Let us consider a general QFT with variable  $\phi = (\phi_1, \dots, \phi_n)$  measure  $d\phi$  and action  $S(\phi)$ .

Suppose  $\phi \rightarrow \phi + \delta\phi$  is a symmetry,  $\delta(d\phi e^{-S(\phi)}) = 0$ .

Then, we have Ward identity

$$\begin{aligned} 0 &= \int \delta(d\phi e^{-S(\phi) + J \cdot \phi}) \\ &= \int d\phi e^{-S(\phi) + J \cdot \phi} J \cdot \delta\phi = e^{-W(J)} J \cdot \langle \delta\phi \rangle_J \end{aligned}$$

Set  $J = J(\phi)$  and use  $\frac{\partial \Gamma}{\partial \phi_i}(\phi) = J_i(\phi)$ . We obtain

$$\sum_i \langle \delta\phi_i \rangle_{J(\phi)} \frac{\partial \Gamma}{\partial \phi_i}(\phi) = 0. \quad \text{Slavnov-Taylor identity}$$

i.e.  $\Gamma(\phi)$  is invariant under  $\phi \rightarrow \phi + \langle \delta\phi \rangle_{J(\phi)}$ .

For an at most linear symmetry:  $\delta\phi_i = M_{ij} \phi_j + C_i$ ,

$$\langle \delta\phi_i \rangle_{J(\phi)} = M_{ij} \langle \phi_j \rangle_{J(\phi)} + C_i = M_{ij} \phi_j + C_i = \delta\phi_i.$$

So  $\Gamma(\phi)$  is invariant under the original symmetry.

A variant : non symmetry

$$\text{eg. } \delta(d\phi e^{-S(\phi)}) = d\phi e^{-S(\phi)} (-\delta S(\phi)).$$

$$\begin{aligned} \text{Then } 0 &= \int d\phi e^{-S(\phi) + J \cdot \phi} (-\delta S(\phi) + J \cdot \delta\phi) \\ &= e^{-W(J)} \left( -\langle \delta S(\phi) \rangle_J + J \cdot \langle \delta\phi \rangle_J \right) \end{aligned}$$

and setting  $J = J(\phi)$  we have

$$\sum_i \langle \delta\phi_i \rangle_{J(\phi)} \frac{\partial \Gamma}{\partial \phi_i}(\phi) = \langle \delta S(\phi) \rangle_{J(\phi)}.$$

If both  $\delta\phi$  &  $\delta S(\phi)$  are at most linear,

$$\delta\Gamma(\phi) = \delta S(\phi).$$

End of Digression

Let us apply this to the gauge fixed QED and the gauge transformation

$$\delta A_\mu = -\frac{1}{e} \partial_\mu \alpha, \quad \delta \psi = i\alpha \psi, \quad \delta \bar{\psi} = \bar{\psi}(-i\alpha).$$

This is at most linear, and also

$$\delta S = \int d^4x \frac{1}{\xi} \left(-\frac{1}{e} \partial^2 \alpha\right) \partial \cdot A \quad \text{is linear.}$$

Therefore (the variant of) Slavnov-Taylor identity reads

$$\delta \Gamma[A, \psi, \bar{\psi}] = \int d^4x \frac{1}{\xi} \left(-\frac{1}{e} \partial^2 \alpha\right) \partial \cdot A.$$

That is, if we set

$$\Gamma[A, \psi, \bar{\psi}] = \int d^4x \frac{1}{2\xi} (\partial \cdot A)^2 + \Gamma^{\text{inv}}[A, \psi, \bar{\psi}],$$

then,  $\Gamma^{\text{inv}}[A, \psi, \bar{\psi}]$  is gauge invariant,

$$\delta \Gamma^{\text{inv}}[A, \psi, \bar{\psi}] = 0.$$

In  $\Gamma[A, \psi, \bar{\psi}]$ , the gauge fixing term of the classical Lagrangian is the only term that breaks the gauge symmetry.

I.e. "Gauge fixing term is not renormalized."

Note:  $\Gamma^{inv}[A, \psi, \bar{\psi}]$

$$= \int \frac{d^4 p}{(2\pi)^4} \left\{ \frac{1}{2} A_\mu(-p) \left( \delta^{\mu\nu} p^2 - p^\mu p^\nu - \Pi^{\mu\nu}(p) \right) A_\nu(p) \right. \\ \left. + \bar{\psi}(-p) \left( -\not{p} + m + \Sigma(p) \right) \psi(p) \right. \\ \left. + \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 \ell}{(2\pi)^4} \bar{\psi}(-p-\ell) e \Gamma^M(p, \ell) A_\mu(\ell) \psi(p) \right. \\ \left. + \text{higher power} \right.$$

$$\delta A_\mu(p) = \frac{i}{e} p_\mu \alpha(p)$$

$$\delta \psi(p) = \int \frac{d^4 p_1}{(2\pi)^4} \frac{d^4 p_2}{(2\pi)^4} (2\pi)^4 \delta(p_1 + p_2 - p) i \alpha(p_1) \psi(p_2)$$

$$\delta \bar{\psi}(-p) = \int \frac{d^4 p_1}{(2\pi)^4} \frac{d^4 p_2}{(2\pi)^4} (2\pi)^4 \delta(p + p_1 - p_2) \bar{\psi}(-p_2) (-i \alpha(p_1))$$

$$\delta \Gamma^{inv}[A, \psi, \bar{\psi}] = 0$$

$$\Rightarrow \begin{cases} p_\mu \Pi^{\mu\nu}(p) = 0 \\ \not{\ell}_\mu \Gamma^M(p, \ell) - \not{p} + \Sigma(p+\ell) - \Sigma(p) = 0 \end{cases}$$

The identities are derived in another way (though the origin is the same).



## Another application

The (gauge fixed) Lagrangian of massless QED

$$\mathcal{L} = \frac{1}{4} F^{\mu\nu} F_{\mu\nu} + i \bar{\Psi} (-\gamma^\mu D_\mu) \Psi + \frac{1}{2\xi} (\partial \cdot A)^2$$

has axial symmetry ( $\gamma_5 = \gamma^1 \gamma^2 \gamma^3 \gamma^4$ ,  $\beta$ : constant)

$$A_\mu \rightarrow A_\mu, \quad \Psi \rightarrow e^{i\beta\gamma_5} \Psi, \quad \bar{\Psi} \rightarrow \bar{\Psi} e^{i\beta\gamma_5}$$

Indeed,  $\gamma^\mu \gamma_5 = -\gamma_5 \gamma^\mu$  and hence

$$e^{i\beta\gamma_5} \gamma^\mu e^{i\beta\gamma_5} = e^{i\beta\gamma_5} e^{-i\beta\gamma_5} \gamma^\mu = \gamma^\mu.$$

(A mass term  $\propto \bar{\Psi} \Psi$  would not be invariant.)

Suppose the path-integral measure is also invariant.

As the transformation is at most linear, by Slavnov-Taylor

identity,  $\Gamma[A, \Psi, \bar{\Psi}]$  is also invariant. In particular,

$$e^{i\beta\gamma_5} \Sigma(p) e^{i\beta\gamma_5} \stackrel{!}{=} \Sigma(p).$$

This requires  $\Sigma(p) \propto \not{p}$  in the massless theory.

If we recover  $m \neq 0$ ,  $\Sigma(p)$  must be of the form

$$\Sigma(p) = A(p^2) \not{p} + B(p^2) m.$$

## Structure of divergence

On dimensional ground, the divergence as the UV cutoff  $\Lambda$  is removed ( $\Lambda \rightarrow \infty$ ) must be of the form

$$\Pi^{\mu\nu}(p) = (\delta^{\mu\nu} p^2 - p^\mu p^\nu) a_1 \log \Lambda + \text{finite},$$

$$\Sigma(p) = a_2 \log \Lambda \not{p} + a_3 \Lambda + a_4 \log \Lambda m + \text{finite},$$

$$\Gamma^M(p, q) = a_5 \log \Lambda \gamma^M + \text{finite}$$

with some constants  $a_1, a_2, a_3, a_4, a_5$

- By  $q_\mu \Gamma^M(p, q) = \not{q} + \Sigma(p) - \Sigma(p+q)$ ,

we find  $a_5 = -a_2$ .

- By the axial symmetry of the  $m=0$  theory

we also find  $a_3 = 0$ .

Indeed, at the one loop level (with  $M_1 \sim M_2 \sim \Lambda$ ),

$$a_1 = -\frac{8e^2}{3(4\pi)^2}, \quad a_2 = -\frac{2e^2}{(4\pi)^2}, \quad a_3 = 0, \quad a_4 = \frac{8e^2}{(4\pi)^2},$$

$$a_5 = \frac{2e^2}{(4\pi)^2}. \quad a_2 + a_5 = a_3 = 0 \text{ is satisfied.}$$

## Renormalization

In view of the structure of divergence, after regularization, we can renormalize the theory as

$$\mathcal{L} = \frac{1}{4e_0^2} F_0^{\mu\nu} F_{0\mu\nu} + \bar{\Psi}_0 (-i\not{D}_{A_0} + m_0) \Psi_0 + \frac{1}{2e_0^2 \xi_0} (\partial \cdot A_0)^2$$

$$\frac{1}{e_0^2} = \frac{Z_3}{e^2}, \quad \Psi_0 = \sqrt{Z_2} \Psi, \quad \bar{\Psi}_0 = \sqrt{Z_2} \bar{\Psi}, \quad Z_2 m_0 = Z_m m$$

$$A_0 = eA, \quad e_0^2 \xi_0 = e^2 \xi$$

$$= \frac{Z_3}{4} F^{\mu\nu} F_{\mu\nu} + \bar{\Psi} (-iZ_2 \not{D} + Z_m m) \Psi + e Z_2 \bar{\Psi} A \Psi + \frac{1}{2\xi} (\partial \cdot A)^2$$

- $Z_3$  takes care of the divergence of  $\Pi^{\mu\nu}(p) \leftrightarrow F^{\mu\nu} F_{\mu\nu}$ .
- $Z_2$  takes care of the common divergence of  
   $\not{D}$  part of  $\Sigma(p) \leftrightarrow i\bar{\Psi} \not{D} \Psi$  and  $\Gamma^{\mu}(p, \xi) \leftrightarrow e\bar{\Psi} A \Psi$ .
- $Z_m$  takes care of the divergence of  
  m.id part of  $\Sigma(p) \leftrightarrow \bar{\Psi} m \Psi$ .

$$Z_3 = 1 + \delta_3^{(1)} + \delta_3^{(2)} + \dots$$

$$Z_2 = 1 + \delta_2^{(1)} + \delta_2^{(2)} + \dots$$

$$Z_n = 1 + \delta_n^{(1)} + \delta_n^{(2)} + \dots$$

We determine  $\delta_3^{(a)}$ ,  $\delta_2^{(a)}$ ,  $\delta_n^{(a)}$  order by order in perturbation theory so that

$$\begin{aligned} \Gamma_0[A_0, \psi_0, \bar{\psi}_0, e_0(\Lambda), m_0(\Lambda), \xi_0; \Lambda] \\ = \Gamma[A, \psi, \bar{\psi}, e, m, \xi; \Lambda] \end{aligned}$$

is finite as a function of  $A, \psi, \bar{\psi}, e, m, \xi$  as the UV cut-off  $\Lambda$  is removed.

There is an ambiguity in the choice of renormalized fields and couplings, but that is fixed by renormalization condition.

## Renormalization condition

$$\Pi^{\mu\nu}(q) = (\delta^{\mu\nu} q^2 - q^\mu q^\nu) \Pi(q^2)$$

$$\Sigma(p) = A(p^2) \not{p} + B(p^2) m$$

$$\left\{ \begin{array}{l} \Pi(0) = 0 \\ A(-m^2) = 0 \\ B(-m^2) = 0 \end{array} \right.$$

On shell renormalization

$$\left\{ \begin{array}{l} \Pi(\mu^2) = 0 \\ A(\mu^2) = 0 \\ B(\mu^2) = 0 \end{array} \right.$$

Another R.C.

$\mu$ : a mass scale.

## Solution at 1-loop

$$\Pi^{(1)}(q^2) = \Pi_2(q^2) - \delta_3^{(1)},$$

$$A^{(1)}(p^2) = A_2(p^2) - \delta_2^{(1)},$$

$$B^{(1)}(p^2) = B_2(p^2) + \delta_m^{(1)},$$

where (for  $j=1$ )

$$\Pi_2(q^2) = -\frac{8e^2}{(4\pi)^2} \int_0^1 dx x(1-x) \log\left(\frac{M^2}{m^2 + x(1-x)q^2}\right)$$

$$A_2(p^2) = -\frac{2e^2}{(4\pi)^2} \int_0^1 dx x \log\left(\frac{x\Lambda^2}{(1-x)m^2 + x(1-x)p^2}\right)$$

$$B_2(p^2) = \frac{4e^2}{(4\pi)^2} \int_0^1 dx \log\left(\frac{x\Lambda^2}{(1-x)m^2 + x(1-x)p^2}\right)$$

On shell  $\delta_3^{(1)} = \Pi_2(0)$ ,  $\delta_2^{(1)} = A_2(-m^2)$ ,  $\delta_m^{(1)} = -B_2(-m^2)$

Another  $\delta_3^{(1)} = \Pi_2(\mu^2)$ ,  $\delta_2^{(1)} = A_2(\mu^2)$ ,  $\delta_m^{(1)} = -B_2(\mu^2)$

The result (on shell R.C.):

$$\Pi^{(1)}(q^2) = \frac{8e^2}{(4\pi)^2} \int_0^1 dx x(1-x) \log\left(\frac{m^2 + x(1-x)q^2}{m^2}\right)$$

$$\Sigma^{(1)}(p) = \frac{2e^2}{(4\pi)^2} \int_0^1 dx (x\not{p} - 2m) \log\left(\frac{m^2 + xp^2}{(1-x)m^2}\right)$$

$$\Gamma^{(1)\mu}(p, q) = \gamma^\mu - \frac{2e^2}{(4\pi)^2} \int_{\triangle} dy dz \left[ \log\left(\frac{(1-z)\Delta_0}{z^3 m^2}\right) \gamma^\mu - \frac{X^\mu}{\Delta_0} \right]$$

$$\Delta_0 = y(1-y)q^2 + z(1-z)p^2 + 2yzpq + (1-z)m^2$$

$$X^\mu = (-y\not{p} + z\not{p})\gamma^\mu((1-y)\not{p} + z\not{p}) + 2m((1-2y)q + 2zp)^\mu + m^2\gamma^\mu$$

meaning of  $\Pi(q^2)$  :  $\Gamma$  enters into  $\Gamma[A, \psi, \bar{\psi}]$  as

$$\Gamma^{\text{inv}}[A, \psi, \bar{\psi}] = \int \frac{d^4 q}{(2\pi)^4} \frac{1}{2} A_\mu(-q) (\delta^{\mu\nu} q^2 - q^\mu q^\nu) (1 - \Pi(q^2)) A_\nu(q) + \dots$$


$$\leadsto e_{\text{eff}}^2(q^2) = \frac{e^2}{1 - \Pi(q^2)}$$

effective gauge coupling constant

at energy scale  $q^2 \approx$  length scale  $1/q^2$ .

$$\Pi(q^2) \rightarrow \begin{cases} O(q^2/m^2) & \text{as } q^2/m^2 \rightarrow 0 \\ \frac{4e^2}{3(4\pi)^2} \left[ \log\left(\frac{q^2}{m^2}\right) - \frac{5}{3} + O\left(\frac{m^2}{q^2}\right) \right] & \text{as } q^2 \gg m^2 \end{cases}$$

$$e_{\text{eff}}^2(q^2) \rightarrow \begin{cases} e^2 & \text{as } q^2/m^2 \rightarrow 0 \\ \frac{e^2}{1 - \frac{4e^2}{3(4\pi)^2} \left( \log\left(\frac{q^2}{m^2}\right) - \frac{5}{3} \right)} & \text{as } q^2/m^2 \gg 1 \end{cases}$$




$e_{\text{eff}}^2(q^2) \sim e^2$  at long distances  $|q|^{-1} \gg \frac{1}{m}$

$e_{\text{eff}}^2(q^2)$  grows at short distances  $|q|^{-1} \ll \frac{1}{m}$

Interpretation:

“charge screening by vacuum polarization”





## Renormalization group

Take the "Another renormalization condition", and denote the 1PI effective action as  $\Gamma[A, \psi, \bar{\psi}, e, m, \xi; \mu, \Lambda]$ .

Change the renormalization point  $\mu$  while fixing bare fields/couplings  $A_0, \psi_0, \bar{\psi}_0, e_0, m_0, \xi_0$ . Then, we have RGE

$$\begin{aligned} 0 &= \mu \frac{d}{d\mu} \Gamma[A_0, \psi_0, \bar{\psi}_0, e_0, m_0, \xi_0; \Lambda] \\ &= \mu \frac{d}{d\mu} \Gamma[A, \psi, \bar{\psi}, e, m, \xi; \mu, \Lambda] \end{aligned}$$

We denote

$$\mu \frac{d}{d\mu} \Phi_\Sigma = -\gamma_\Sigma \Phi_\Sigma \quad \text{for } \Phi_\Sigma = A, \psi, \bar{\psi}, m \quad \text{and}$$

$$\mu \frac{d}{d\mu} e = \beta.$$

As  $A_0 = eA$  and  $e_0^2 \xi_0 = e^2 \xi$ , we find

$$0 = \beta A + e(-\gamma_A A), \quad 0 = 2e\beta \xi + e^2 \mu \frac{d\xi}{d\mu}.$$

$$\text{i.e. } \gamma_A = \frac{1}{e} \beta \quad \& \quad \mu \frac{d\xi}{d\mu} = -2 \frac{1}{e} \beta \xi = -2\gamma_A \xi$$

Thus, the RGE (in the limit  $\Lambda \rightarrow \infty$ ) reads

$$\left( \mu \frac{\partial}{\partial m} - \frac{1}{e} \beta A \cdot \frac{\delta}{\delta A} - \gamma_{\psi} \psi \frac{\delta}{\delta \psi} - \gamma_{\bar{\psi}} \bar{\psi} \frac{\delta}{\delta \bar{\psi}} - \gamma_m m \frac{\partial}{\partial m} + \beta \frac{\partial}{\partial e} - \frac{2}{e} \beta \xi \frac{\partial}{\partial \xi} \right) \Gamma[A, \psi, \bar{\psi}, e, m, \xi; \mu] = 0.$$

As  $\frac{1}{e_0} = \frac{Z_3}{e^2}$ ,  $\psi_0 = \sqrt{Z_2} \psi$ ,  $\bar{\psi}_0 = \sqrt{Z_2} \bar{\psi}$ ,  $Z_2 m_0 = Z_m m$ ,

$$0 = -2e^{-3} \beta Z_3 + e^{-2} \mu \frac{d}{d\mu} Z_3,$$

$$0 = \mu \frac{d}{d\mu} \sqrt{Z_2} \psi + \sqrt{Z_2} (-\gamma_{\psi} \psi),$$

$$0 = \mu \frac{d}{d\mu} Z_2^{-1} Z_m m + Z_2^{-1} \mu \frac{d}{d\mu} Z_m m + Z_2^{-1} Z_m (-\gamma_m m).$$

$$\beta = \frac{1}{2} e \mu \frac{d}{d\mu} \log Z_3$$

$$\gamma_{\psi} = \frac{1}{2} \mu \frac{d}{d\mu} \log Z_2$$

$$\gamma_m = \mu \frac{d}{d\mu} \log Z_m - \mu \frac{d}{d\mu} \log Z_2$$

Computation at 1-loop (at  $\xi = ()$ ):

$$\beta^{(1)} = \frac{1}{2} e \mu \frac{d}{d\mu} \delta_3^{(1)} = \frac{4e^3}{(4\pi)^2} \int_0^1 dx x(1-x) \frac{x(1-x) 2\mu^2}{m^2 + x(1-x)\mu^2}$$

$$\gamma_4^{(1)} = \frac{1}{2} \mu \frac{d}{d\mu} \delta_2^{(1)} = \frac{e^2}{(4\pi)^2} \int_0^1 dx x \frac{x 2\mu^2}{m^2 + x\mu^2}$$

$$\gamma_m^{(1)} = \mu \frac{d}{d\mu} (\delta_m^{(1)} - \delta_2^{(1)}) = \frac{2e^2}{(4\pi)^2} \int_0^1 dx (2-x) \frac{x 2\mu^2}{m^2 + x\mu^2}$$

The result for  $\beta^{(1)}$  is valid for any  $\xi$

since  $\overbrace{\psi(x)\psi(y)}$  is independent of  $\xi$ .

The result for  $\gamma_4^{(1)}$  &  $\gamma_m^{(1)}$  depends on  $\xi$ .

(See the additional note for the expressions.)

Note that

$$\beta^{(1)} \approx \begin{cases} \frac{4e^3}{3(4\pi)^2} & \mu \gg m \\ \frac{4e^2}{15(4\pi)^2} \frac{\mu^2}{m^2} \sim 0 & \mu \ll m \end{cases}$$

Just like the  $\phi^4$  theory.

$$\underline{\mu \gg m}: \quad \beta \sim \frac{4e^3}{3(4\pi)^2} \quad \Leftrightarrow \quad \mu \frac{d}{d\mu} \frac{1}{e^2} \sim -\frac{8}{3(4\pi)^2}$$


$$\frac{1}{e^2(\mu)} - \frac{1}{e^2(\mu_0)} \sim -\frac{8}{3(4\pi)^2} \log(\mu/\mu_0)$$

$$e^2(\mu) \sim \frac{e^2(\mu_0)}{1 - \frac{8e^2(\mu_0)}{3(4\pi)^2} \log(\mu/\mu_0)}$$

The gauge coupling is weaker at lower energies  
or stronger at higher energies,

$$\underline{\mu \ll m}: \quad \beta \sim 0$$

The gauge coupling stops running.



As we've already seen.

The massless theory  $m = 0$  :

$$e^2(\mu) \sim \frac{e^2(\mu_0)}{1 - \frac{8 e^2(\mu_0)}{3(4\pi)^2} \log(\mu/\mu_0)} \quad \text{valid at low } \mu$$

$$\rightarrow 0 \quad \text{as} \quad \mu \rightarrow 0$$

infra-red free!

## A simple generalization

QED with electrons with charges  $Q_1, \dots, Q_N$

$$\mathcal{L} = \frac{1}{4e^2} F^{\mu\nu} F_{\mu\nu} + \sum_{i=1}^N \bar{\Psi}_i (-i \not{D}_A + m_i) \Psi_i$$

$$\not{D}_A \Psi_i = \gamma^\mu (\partial_\mu + i Q_i A_\mu) \Psi_i$$

$$\Pi_2(q) = \sum_i \text{Diagram}$$

$$= \sum_i - \frac{8(Q_i e)^2}{(4\pi)^2} \int_0^1 dx x(1-x) \log\left(\frac{M^2}{m_i^2 + x(1-x)q^2}\right)$$

With "another R.C."  $\Pi(\mu^2) = 0, \dots,$

$$\beta^{(1)} = \frac{1}{2} e \mu \frac{d}{d\mu} \beta^{(1)} = \frac{1}{2} e \mu \frac{d}{d\mu} \Pi_2(\mu^2)$$

$$= \sum_i e \frac{8(Q_i e)^2}{(4\pi)^2} \int_0^1 dx x(1-x) \frac{x(1-x)\mu^2}{m_i^2 + x(1-x)\mu^2}$$

Suppose the masses are well-separated

$$m_1 \ll m_2 \ll \dots \ll m_N.$$

At the energy scale  $m_i \ll \mu \ll m_{i+1}$ ,

$$\beta \sim \frac{4e^3}{3(4\pi)^2} \sum_{j=1}^i Q_j^2$$

The slope depends on the energy scale

