

1  
One thing I forgot to say in the last lecture.

-- particle interpretation.

For a real scalar field, we have seen that there is a unique ground state  $|0\rangle$  of  $E_0 = \int d^3p \frac{1}{2} \sqrt{p^2 + m^2} \delta^{(4)}(0)$  and other states are obtained from  $|0\rangle$  by operating  $a(p)^\dagger$ 's, each increasing the energy by  $\omega_p = \sqrt{p^2 + m^2}$ .

### Interpretation

$|0\rangle$  -- the vacuum

$a(p)^\dagger$  -- creation of a particle of mass  $m$   
and momentum  $p$  ( $\Rightarrow$  energy  $\sqrt{p^2 + m^2}$ )

e.g.

$a(p)^\dagger |0\rangle$  -- a one particle state

$a(p_1)^\dagger a(p_2)^\dagger |0\rangle$  -- a two particle state

$a(p_1)^\dagger a(p_2)^\dagger a(p_3)^\dagger |0\rangle$  -- a three particle state

$\vdots$

These respectively have

momentum

energy -  $E_0$

$P$

$$\sqrt{P^2 + m^2}$$

$P_1 + P_2$

$$\sqrt{P_1^2 + m^2} + \sqrt{P_2^2 + m^2}$$


$P_1 + P_2 + P_3$

$$\sqrt{P_1^2 + m^2} + \sqrt{P_2^2 + m^2} + \sqrt{P_3^2 + m^2}$$

;

;

Plot of momentum-energy spectrum



## Free fermions

A finite system:  $n$  pairs of anticommuting variables

$$\psi_1, \bar{\psi}^1, \psi_2, \bar{\psi}^2, \dots, \psi_n, \bar{\psi}^n$$

$$S_E = \sum_{i,j} \bar{\psi}^i A_{ij} \psi_j$$

$$d\bar{\psi} d\psi = d\bar{\psi}^n \dots d\bar{\psi}^1 d\psi_1 \dots d\psi_n = d\bar{\psi}^1 d\psi_1 \dots d\bar{\psi}^n d\psi_n$$

Partition function is

$$Z = \int d\bar{\psi} d\psi e^{-S_E} = \det A$$

To compute correlation functions, let us introduce

$$f(A, \bar{\eta}, \eta) := \int d\bar{\psi} d\psi e^{-S_E + \sum_i (\bar{\eta}^i \psi_i + \bar{\psi}^i \eta_i)}$$

Note:

$$\begin{aligned} & \frac{\partial}{\partial \bar{\eta}^{i_1}} \frac{\partial}{\partial \bar{\eta}^{i_2}} \dots \frac{\partial}{\partial \bar{\eta}^{i_s}} e^{\bar{\eta} \psi + \bar{\psi} \eta} \overset{\leftarrow}{\frac{\partial}{\partial \eta^{j_1}}} \overset{\leftarrow}{\frac{\partial}{\partial \eta^{j_2}}} \dots \overset{\leftarrow}{\frac{\partial}{\partial \eta^{j_t}}} \\ &= e^{\bar{\eta} \psi} \psi_{i_1} \dots \psi_{i_s} \bar{\psi}^{j_1} \dots \bar{\psi}^{j_t} e^{\bar{\psi} \eta} \end{aligned}$$

Thus,

$$\begin{aligned} & \frac{\partial}{\partial \bar{\eta}^{i_1}} \frac{\partial}{\partial \bar{\eta}^{i_2}} \dots \frac{\partial}{\partial \bar{\eta}^{i_s}} f(A, \bar{\eta}, \eta) \overset{\leftarrow}{\frac{\partial}{\partial \eta^{j_1}}} \overset{\leftarrow}{\frac{\partial}{\partial \eta^{j_2}}} \dots \overset{\leftarrow}{\frac{\partial}{\partial \eta^{j_t}}} \Big|_{\bar{\eta} = \eta = 0} \\ &= Z \langle \psi_{i_1} \dots \psi_{i_s} \bar{\psi}^{j_1} \dots \bar{\psi}^{j_t} \rangle \end{aligned}$$

$$f(A, \bar{\eta}, \eta) = \int d\bar{\psi} d\psi e^{-(\bar{\psi} - \bar{\eta} A^{-1}) A (\psi - A^{-1} \eta) + \bar{\eta} A^{-1} \eta}$$

$$= Z e^{\bar{\eta} A^{-1} \eta}$$

$$\therefore \langle \psi_{i_1} \dots \psi_{i_s} \bar{\psi}^{j_1} \dots \bar{\psi}^{j_t} \rangle$$

$$= \frac{\partial}{\partial \bar{\eta}^{i_1}} \frac{\partial}{\partial \bar{\eta}^{i_2}} \dots \frac{\partial}{\partial \bar{\eta}^{i_s}} e^{\bar{\eta} A^{-1} \eta} \overleftarrow{\frac{\partial}{\partial \eta^{j_1}}} \overleftarrow{\frac{\partial}{\partial \eta^{j_2}}} \dots \overleftarrow{\frac{\partial}{\partial \eta^{j_t}}} \Big|_{\bar{\eta} = \eta = 0}$$

$$= (A^{-1} \eta)_{i_1} \dots (A^{-1} \eta)_{i_s} \overleftarrow{\frac{\partial}{\partial \eta^{j_1}}} \overleftarrow{\frac{\partial}{\partial \eta^{j_2}}} \dots \overleftarrow{\frac{\partial}{\partial \eta^{j_t}}} \Big|_{\eta = 0}$$

This is non-zero only if  $s=t$ .

$$\text{e.g. } \langle \psi_i \bar{\psi}^j \rangle = \underbrace{(A^{-1} \eta)_i}_{A^{-1}{}^k{}_i \eta_k} \overleftarrow{\frac{\partial}{\partial \eta^j}} = A^{-1}{}^j{}_i$$

$$\langle \psi_i \psi_j \bar{\psi}^k \bar{\psi}^l \rangle = (A^{-1} \eta)_i (A^{-1} \eta)_j \overleftarrow{\frac{\partial}{\partial \eta^k}} \overleftarrow{\frac{\partial}{\partial \eta^l}}$$

$$= A^{-1}{}^l{}_i A^{-1}{}^k{}_j \text{---} A^{-1}{}^k{}_i A^{-1}{}^l{}_j$$

$\overleftarrow{\frac{\partial}{\partial \eta^k}}$  passes through  $\eta$  in  $(A^{-1} \eta)_j$



The result can also be presented as the sum of Wick contractions, with the understanding that a  $(-1)$  is produced each time two fermionic objects are swapped:

$$\langle \psi_i \bar{\psi}^j \rangle = \overbrace{\psi_i \bar{\psi}^j} = A_i^{-1 j}$$

$$\begin{aligned} \langle \psi_i \psi_j \bar{\psi}^k \bar{\psi}^l \rangle &= \overbrace{\psi_i \psi_j \bar{\psi}^k \bar{\psi}^l}^+ + \overbrace{\psi_i \psi_j \bar{\psi}^k \bar{\psi}^l}^- \\ &= \overbrace{\psi_i \bar{\psi}^l} \overbrace{\psi_j \bar{\psi}^k} - \overbrace{\psi_i \bar{\psi}^k} \overbrace{\psi_j \bar{\psi}^l} \\ &= A_i^{-1 l} A_j^{-1 k} - A_i^{-1 k} A_j^{-1 l} \end{aligned}$$

⋮

- We see that everything is determined by the two point functions

$$\langle \psi_i \bar{\psi}^j \rangle = \overbrace{\psi_i \bar{\psi}^j} = A_i^{-1 j}$$

- The logic holds also when  $n = \infty$ , e.g. in QFT in dimension  $d \geq 1$ . We now apply this to important examples.

The focus will thus be two point functions.

# Dirac fermions in d dimensions

## Gamma matrices

The algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$$

$$\text{ie. } \begin{cases} (\gamma^0)^2 = 1, & (\gamma^i)^2 = -1 \text{ for } i=1, \dots, d-1, \\ \gamma^\mu \gamma^\nu = -\gamma^\nu \gamma^\mu \text{ for } 0 \leq \mu < \nu \leq d-1 \end{cases}$$

.... called the Clifford algebra in dimension  $(d-1, 1)$

has an irreducible representation on a  $\mathbb{C}$ -vector space  $S$  of dimension  $d_S = 2^{\lfloor \frac{d}{2} \rfloor} = \begin{cases} 2^n & d=2n \text{ even} \\ 2^n & d=2n+1 \text{ odd.} \end{cases}$

•  $S$  has a hermitian inner product s.t.

$$\gamma^{0\dagger} = \gamma^0 \text{ hermitian, } \gamma^{i\dagger} = -\gamma^i \text{ antihermitian.}$$

•  $\text{tr}_S(\gamma^{\mu_1} \dots \gamma^{\mu_\ell}) = 0$  for distinct  $\mu_1, \dots, \mu_\ell$ 's

except  $\{\mu_1, \dots, \mu_\ell\} = \{0, \dots, d-1\}$  for  $d$  odd.

In particular,  $\text{tr}_S \gamma^\mu = 0$  if  $d > 1$ .

•  $S$  is a representation of Lorentz group  $SO(d-1,1)$

$$e^{\omega} \in SO(d-1,1) : \varphi \in S \mapsto e^{\frac{1}{2}\gamma(\omega)} \varphi \in S$$

$$\text{where } \gamma(\omega) := \frac{1}{2} \omega_{\mu\nu} [\gamma^{\mu}, \gamma^{\nu}] =: \omega_{\mu\nu} \gamma^{\mu\nu}.$$

$\left\{ \begin{array}{l} d \text{ odd: irreducible} \end{array} \right.$

$\left\{ \begin{array}{l} d \text{ even: splits to two irreducibles, distinguished by the action} \\ \text{of } \gamma^0 \gamma^1 \dots \gamma^{d-1}. \end{array} \right.$

• For  $\varphi \in S$ , define  $\bar{\varphi} \in S^*$  by

$$\bar{\varphi} \varphi' := \varphi^{\dagger} \gamma^0 \varphi' \quad \text{for } \varphi' \in S.$$

Then,  $\bar{\varphi} \varphi'$  is a Lorentz scalar and

$\bar{\varphi} \gamma^{\mu} \varphi'$  is a Lorentz vector.

$$\odot (\gamma^0 \gamma^i)^{\dagger} = -\gamma^i \gamma^0 = \gamma^0 \gamma^i, \quad (\gamma^i \gamma^j)^{\dagger} = \gamma^j \gamma^i = -\gamma^i \gamma^j \quad (i \neq j)$$

$$\therefore (\gamma^{\mu\nu})^{\dagger} \gamma^0 = -\gamma^0 \gamma^{\mu\nu} \quad \therefore (e^{\frac{1}{2}\gamma(\omega)})^{\dagger} \gamma^0 = \gamma^0 e^{-\frac{1}{2}\gamma(\omega)}$$

$$\begin{aligned} \left[ \frac{1}{2}\gamma(\omega), \gamma^{\mu} \right] &= \frac{1}{4} \omega_{\rho\lambda} \left[ \gamma^{\rho} \gamma^{\lambda}, \gamma^{\mu} \right] = \gamma^{\rho} \left\{ \frac{\gamma^{\lambda} \gamma^{\mu} - \gamma^{\mu} \gamma^{\lambda}}{2\eta^{\lambda\mu}} \right\} - \gamma^{\lambda} \left\{ \frac{\gamma^{\rho} \gamma^{\mu} - \gamma^{\mu} \gamma^{\rho}}{2\eta^{\rho\mu}} \right\} \\ &= -\omega^{\mu\nu} \gamma^{\nu}. \end{aligned}$$

$$\therefore e^{-\frac{1}{2}\gamma(\omega)} \gamma^{\mu} e^{\frac{1}{2}\gamma(\omega)} = (e^{\omega})^{\mu}{}_{\nu} \gamma^{\nu} \quad //$$

## The system

Variable : an  $S$ -valued anticommuting function  $\psi(x)$  of  $d$  dimensional spacetime.

Lagrangian :  $\mathcal{L} = i\bar{\psi} \not{\partial} \psi - m\bar{\psi}\psi$

where  $\cdot \not{\partial} := \gamma^\mu \partial_\mu$

$\cdot m$  is a real parameter  $m^* = m$ .

- $\mathcal{L}$  is real modulo total derivative

$$\mathcal{L}^* = -i \partial_\mu \psi^\dagger \underbrace{(\gamma^0 \gamma^\mu)^\dagger}_{\gamma^0 \gamma^\mu} \psi - m \psi^\dagger \underbrace{\gamma^0}_{\gamma^0} \psi \quad \bar{\psi}\psi$$

$$\underbrace{-i \partial_\mu \bar{\psi} \gamma^\mu \psi}_{-i \partial_\mu \bar{\psi} \gamma^\mu \psi}$$

$$= \partial_\mu (-i \bar{\psi} \gamma^\mu \psi) + i \bar{\psi} \gamma^\mu \partial_\mu \psi - m \bar{\psi} \psi$$

- The system has Poincaré symmetry (translations & Lorentz) and phase rotation symmetry  $\psi \rightarrow e^{-i\alpha} \psi$ .

Charge densities for translations & phase rotation are

$$T_0^\alpha = -i \bar{\psi} \boldsymbol{\gamma} \cdot \nabla \psi + m \bar{\psi} \psi, \quad T_j^\alpha = i \psi^\dagger \partial_j \psi$$

$$J^\alpha = \psi^\dagger \psi.$$

Wick rotation  $x^0 \rightarrow -ix^d$

$$\gamma^0 \rightarrow -i\gamma_E^d, \quad \gamma_E^i = \gamma^i \quad i=1, \dots, d-1.$$

$$\{\gamma_E^M, \gamma_E^N\} = -2\delta^{M,N} \quad 1 \leq M, N \leq d.$$

$$\mathcal{L}_E = -i\bar{\Psi}\gamma_E^M\partial_M\Psi + m\bar{\Psi}\Psi = \bar{\Psi}(-i\not{\partial}_E + m)\Psi.$$

$$\langle \Psi(x)\bar{\Psi}(y) \rangle_E = (-i\not{\partial}_E + m)^{-1}_{xy}$$

$$= \int \frac{d^d p_E}{(2\pi)^d} \frac{e^{-ip_E(x-y)}}{-\not{p}_E + m}$$

$$\left[ (-\not{p}_E + m)(\not{p}_E + m) = -\not{p}_E^2 + m^2 = p_E^2 + m^2 \right]$$

$$= \int \frac{d^d p_E}{(2\pi)^d} \frac{e^{-ip_E(x-y)}}{p_E^2 + m^2} (\not{p}_E + m)$$

Here, we regard  $\langle \Psi(x)\bar{\Psi}(y) \rangle_E$  as  $\text{End}(S)$  valued function.

In components with respect to a basis  $\{e^\alpha\} \subset S$ , it reads

$$\langle \Psi_\alpha(x)\bar{\Psi}^\beta(y) \rangle_E = \int \frac{d^d p_E}{(2\pi)^d} \frac{e^{-ip_E(x-y)}}{p_E^2 + m^2} (\not{p}_E + m)_\alpha^\beta$$


$$(\gamma_E^M)_\alpha^\beta p_{E\mu} + m\delta_\alpha^\beta$$

## Minkowski limit

Proceeding just as in the case of the scalar field, we find that under the reverse Wick rotation

$$x^d \rightarrow ix^0, \quad p_d \rightarrow -ip_0, \quad \gamma^d \rightarrow i\gamma^0,$$

$\langle \psi(x) \bar{\psi}(y) \rangle_E$  goes to:

$$\langle \psi(x) \bar{\psi}(y) \rangle = \int_{\mathbb{R}^{d-1} \times \mathcal{C}} \frac{d^{d-1}p dp_0}{(2\pi)^d} \frac{i e^{-iP \cdot (x-y) - ip_0(x^0-y^0)}}{(p_0 - \omega_p)(p_0 + \omega_p)} (\not{p} + m)$$


$$\text{where } \omega_p = \sqrt{p^2 + m^2}$$

$$\text{or } = \int \frac{d^d p}{(2\pi)^d} \frac{i e^{-ip(x-y)}}{p^2 - m^2 + i0} (\not{p} + m)$$

$$= \int \frac{d^{d-1}p}{(2\pi)^{d-1} 2\omega_p} e^{-i\omega_p|x^0-y^0| - iP \cdot (x-y)} \left( \text{sgn}(x^0-y^0) \omega_p \gamma^0 + \not{P} + m \right)$$

## Canonical quantization ( $d > 1$ )

$$\psi(x) = \int \frac{d^{d-1}p}{(2\pi)^{d-1}} e^{i p \cdot x} \psi(p)$$

$$\begin{aligned} L &= \int d^{d-1}x \left( i \psi^\dagger (\partial_0 + \gamma^0 \boldsymbol{\gamma} \cdot \nabla) \psi - m \bar{\psi} \psi \right) \\ &= \int \frac{d^{d-1}p}{(2\pi)^{d-1}} \left( i \psi(p)^\dagger \dot{\psi}(p) - \psi(p)^\dagger \underbrace{\gamma^0 (\boldsymbol{\gamma} \cdot p + m)} \psi(p) \right) \end{aligned}$$

$$\underline{\Delta_p := \gamma^0 (\boldsymbol{\gamma} \cdot p + m) : \mathcal{S} \rightarrow \mathcal{S}}$$

• It is hermitian, hence diagonalizable

$$\bullet \Delta_p^2 = \gamma^0 (\boldsymbol{\gamma} \cdot p + m) \gamma^0 (\boldsymbol{\gamma} \cdot p + m)$$

$$= \underbrace{\gamma^0 \gamma^0}_1 \underbrace{(-\boldsymbol{\gamma} \cdot p + m)(\boldsymbol{\gamma} \cdot p + m)}_{p^2 + m^2} = p^2 + m^2 = \omega_p^2$$

Thus, eigenvalues of  $\Delta_p$  are  $\pm \omega_p$

$$\bullet \text{tr}_{\mathcal{S}} \Delta_p = 0 \quad \text{since } \text{tr}_{\mathcal{S}}(\gamma^0 \gamma^i) = \text{tr}_{\mathcal{S}}(\gamma^i \gamma^0) = -\text{tr}_{\mathcal{S}}(\gamma^0 \gamma^i)$$

$$\& \text{tr}_{\mathcal{S}} \gamma^0 = 0.$$

Thus, half of the eigenvalues are  $+\omega_p$

and the other halves are  $-\omega_p$ .

Let  $S_{\pm}(P) \subset S$  be the  $\pm \omega_P$  eigen space of  $\Delta_P$ .

Then  $S = S_+(P) \dot{+} S_-(P)$  orthogonal decomposition, and

$$\dim S_+(P) = \dim S_-(P) = \frac{1}{2} \dim S = \frac{1}{2} d_S = 2^{\lfloor d/2 \rfloor - 1}.$$

Let  $\{U_{\pm}^s(P)\}_{s=1}^{d_S/2} \subset S_{\pm}(P)$  be an orthonormal basis.

The elements satisfy

$$\bullet U_{\varepsilon}^s(P)^\dagger U_{\varepsilon'}^{s'}(P) = \delta^{ss'} \delta_{\varepsilon\varepsilon'}$$

$$\bullet \sum_s U_{\pm}^s(P) U_{\pm}^s(P)^\dagger = \text{projection operator to } S_{\pm}(P)$$

$$= \frac{1}{2} \left( 1 \pm \frac{1}{\omega_P} \Delta_P \right)$$

$$\bullet \sum_s U_{\pm}^s(P) \overline{U_{\pm}^s(P)} = \frac{1}{2} \left( 1 \pm \frac{1}{\omega_P} \Delta_P \right) \gamma^0$$

$$= \frac{1}{2\omega_P} \left( \omega_P \gamma^0 \pm \gamma^0 (\not{D} \cdot P + m) \gamma^0 \right)$$

$$\left[ \gamma^0 \gamma^i = -\gamma^i \gamma^0, \gamma^0 \gamma^0 = 1 \right]$$

$$= \frac{1}{2\omega_P} \left( \omega_P \gamma^0 \pm \not{D} \cdot P \pm m \right).$$



Let us expand  $\Psi(\mathbb{P})$  w.r.t. the basis  $\{u_+^s(\mathbb{P})\}_{s=1}^{d_S/2} \cup \{u_-^s(\mathbb{P})\}_{s=1}^{d_S/2}$  of  $S$  as

$$\Psi(\mathbb{P}) = \sum_s \left( u_+^s(\mathbb{P}) b_{+s}(\mathbb{P}) + u_-^s(\mathbb{P}) b_{-s}(\mathbb{P}) \right) \cdot \sqrt{(2\pi)^{d-1}}$$

Then

$$L = \int d^{d-1}\mathbb{P} \sum_{\epsilon, s} \left( b_{\epsilon s}(\mathbb{P})^\dagger \dot{b}_{\epsilon s}(\mathbb{P}) - \epsilon \omega_{\mathbb{P}} b_{\epsilon s}(\mathbb{P})^\dagger b_{\epsilon s}(\mathbb{P}) \right)$$

We know how to quantize such a system (Lecture 3):

$$\{ b_{\epsilon s}(\mathbb{P}), b_{\epsilon' s'}(\mathbb{P}')^\dagger \} = \delta_{\epsilon\epsilon'} \delta_{ss'} \delta^{d-1}(\mathbb{P} - \mathbb{P}'),$$

$$\{ b_{\epsilon s}(\mathbb{P}), b_{\epsilon' s'}(\mathbb{P}') \} = \{ b_{\epsilon s}(\mathbb{P})^\dagger, b_{\epsilon' s'}(\mathbb{P}')^\dagger \} = 0,$$

$$H = \int d^{d-1}\mathbb{P} \sum_{\epsilon s} \epsilon \omega_{\mathbb{P}} b_{\epsilon s}(\mathbb{P})^\dagger b_{\epsilon s}(\mathbb{P}).$$

$$[H, b_{\epsilon s}(\mathbb{P})] = -\epsilon \omega_{\mathbb{P}} b_{\epsilon s}(\mathbb{P}), \quad [H, b_{\epsilon s}(\mathbb{P})^\dagger] = \epsilon \omega_{\mathbb{P}} b_{\epsilon s}(\mathbb{P})^\dagger$$

$\therefore b_{+s}(\mathbb{P})^\dagger, b_{-s}(\mathbb{P})$  : creation operator

$b_{+s}(\mathbb{P}), b_{-s}(\mathbb{P})^\dagger$  : annihilation operator

The state  $|0\rangle$  annihilated by  $b_{+s}(\mathbb{P})$  and  $b_{-s}(\mathbb{P})^\dagger \quad \forall s, \forall \mathbb{P}$

is the unique ground state, with energy

$$E_0 = \int d^{d-1}\mathbb{P} \left( -\frac{d_S}{2} \omega_{\mathbb{P}} \delta^{d-1}(0) \right).$$

Other states are obtained from  $|0\rangle$  by operating  $b_{+s}(p)^\dagger$  &  $b_{-s}(p)$ ,  
each operation increasing energy by  $\omega_p$ .

Remarks • There is no negative norm states.

e.g.  $|p; +s\rangle := b_{+s}(p)^\dagger |0\rangle$ ,  $|p; -s\rangle := b_{-s}(p) |0\rangle$ .

Assuming  $\langle 0|0\rangle = 1$ ,

$$\langle p; +s | p'; +s' \rangle = \langle 0 | b_{+s}(p) b_{+s'}(p')^\dagger | 0 \rangle = \delta_{ss'} \delta^{d-1}(p-p')$$

$$\langle p; -s | p'; -s' \rangle = \langle 0 | b_{-s}(p) b_{-s'}(p') | 0 \rangle = \delta_{ss'} \delta^{d-1}(p-p')$$

$$\cdot Q = \int d^{d-1}p \sum_{\epsilon s} b_{\epsilon s}(p)^\dagger b_{\epsilon s}(p)$$

$$[Q, b_{\epsilon s}(p)] = -b_{\epsilon s}(p), \quad [Q, b_{\epsilon s}(p)^\dagger] = b_{\epsilon s}(p)^\dagger.$$

### Interpretation

$b_{+s}(p)^\dagger / b_{+s}(p)$  is creation/annihilation of a particle  
of mass  $m$ , momentum  $p$ , charge  $+1$

$b_{-s}(p) / b_{-s}(p)^\dagger$  is creation/annihilation of a particle  
of mass  $m$ , momentum  $-p$ , charge  $-1$ .

$S=1, \dots, ds/2$  is the label of representation of the subgroup

of Lorentz group that fixes  $(\omega_p, p)$ :  $\underset{m \neq 0}{SO(d-1)}$  or  $\underset{m=0}{SO(d-2)}$ .

$$\psi(x) = \int \frac{d^4 p}{\sqrt{(2\pi)^{d-1}}} e^{i p \cdot x} \sum_s \left( u_+^s(p) b_{+s}(p) + u_-^s(p) b_{-s}(p) \right)$$

$$\psi(t, x) = e^{i t H} \psi(x) e^{-i t H}$$

$$= \int \frac{d^4 p}{\sqrt{(2\pi)^{d-1}}} e^{i p \cdot x} \sum_s \left( u_+^s(p) e^{-i \omega_p t} b_{+s}(p) + u_-^s(p) e^{i \omega_p t} b_{-s}(p) \right)$$

$$\bar{\psi}(t, x) = \int \frac{d^4 p}{\sqrt{(2\pi)^{d-1}}} e^{-i p \cdot x} \sum_s \left( \overline{u_+^s(p)} e^{+i \omega_p t} b_{+s}(p)^\dagger + \overline{u_-^s(p)} e^{-i \omega_p t} b_{-s}(p)^\dagger \right)$$

$$\langle 0 | \psi(x) \bar{\psi}(y) | 0 \rangle$$

$$= \int \frac{d^4 p_1, d^4 p_2}{(2\pi)^{d-1}} e^{i p_1 \cdot x - i p_2 \cdot y} \sum_{s_1, s_2} e^{-i \omega_{p_1} x^0 + i \omega_{p_2} y^0} u_+^{s_1}(p_1) \overline{u_+^{s_2}(p_2)} \times$$

$$\langle 0 | b_{+s_1}(p_1) b_{+s_2}(p_2)^\dagger | 0 \rangle$$

$\delta_{s_1, s_2} \delta^d(p_1 - p_2)$

$$= \int \frac{d^4 p}{(2\pi)^{d-1}} e^{i p \cdot (x-y) - i \omega_p (x^0 - y^0)} \sum_s \overline{u_+^s(p)} u_+^s(p)$$

$\frac{1}{2\omega_p} (\omega_p y^0 - \vec{\gamma} \cdot \vec{p} + m)$

$$\langle 0 | \bar{\psi}(y) \psi(x) | 0 \rangle \leftarrow \text{regarded as End}(S) \text{ element}$$

$$= \int \frac{d^4 p_1, d^4 p_2}{(2\pi)^{d-1}} e^{i p_1 \cdot x - i p_2 \cdot y} \sum_{s_1, s_2} e^{-i \omega_{p_2} y^0 + i \omega_{p_1} x^0} u_-^{s_1}(p_1) \overline{u_-^{s_2}(p_2)} \times$$

$$\langle 0 | b_{-s_2}(p_2)^\dagger b_{-s_1}(p_1) | 0 \rangle$$

$\delta_{s_2, s_1} \delta^d(p_2 - p_1)$

$$= \int \frac{d^4 p}{(2\pi)^{d-1}} e^{i p \cdot (x-y) - i \omega_p (x^0 - y^0)} \sum_s \overline{u_-^s(p)} u_-^s(p)$$

$\frac{1}{2\omega_p} (\omega_p y^0 + \vec{\gamma} \cdot \vec{p} - m)$

$$\langle 0 | T \psi(x) \bar{\psi}(y) | 0 \rangle = \begin{cases} \langle 0 | \psi(x) \bar{\psi}(y) | 0 \rangle & x^0 > y^0 \\ -\langle 0 | \bar{\psi}(y) \psi(x) | 0 \rangle & y^0 > x^0 \end{cases}$$

$$= \int \frac{d^4 p}{(2\pi)^4} e^{-i\omega_p |x^0 - y^0| + i\mathbf{p} \cdot (\mathbf{x} - \mathbf{y})} \frac{1}{2\omega_p} \left( \text{sgn}(x^0 - y^0) \omega_p \gamma^0 - \boldsymbol{\gamma} \cdot \mathbf{p} + m \right)$$

$$= \int \frac{d^4 p}{(2\pi)^4 2\omega_p} e^{-i\omega_p |x^0 - y^0| - i\mathbf{p} \cdot (\mathbf{x} - \mathbf{y})} \left( \text{sgn}(x^0 - y^0) \omega_p \gamma^0 + \boldsymbol{\gamma} \cdot \mathbf{p} + m \right)$$

--- match with  $\langle \psi(x) \bar{\psi}(y) \rangle$ .

## Ghost system

The gauge fixed Maxwell theory (B-eliminated):

$$\tilde{\mathcal{L}}_E = \frac{1}{4e^2} \sum_{\mu\nu} F_{\mu\nu}^2 + \frac{1}{2e^2} (\partial \cdot A)^2 + \underbrace{\bar{c} \cdot \partial^2 c}_{\text{Consider this part}}$$


Consider this part

$$\langle C(x) \bar{C}(y) \rangle_E = \partial^{-2}_{x,y} = \int \frac{d^d p_E}{(2\pi)^d} \frac{e^{-i p_E (x-y)}}{-p_E^2}$$

reverse Wick rotation:  $x^d \rightarrow i x^d$ ,  $p_d \rightarrow -i p_0$ ,  $\delta^{\mu\nu} \rightarrow -\eta^{\mu\nu}$

Also  $\bar{c} \rightarrow -i \bar{c}$  (sign error in Lec 5)

$$\langle C(x) (-i) \bar{C}(y) \rangle = \int_{\mathbb{R}^{d-1} \times \mathbb{C}} \frac{d^{d-1} p \, d p_0}{(2\pi)^{d-1}} \frac{i e^{-i p \cdot (x-y) - i p_0 (x^0 - y^0)}}{-(p_0 - |p|)(p_0 + |p|)}$$



or

$$\langle C(x) \bar{C}(y) \rangle = \int_{\mathbb{R}^d} \frac{d^d p}{(2\pi)^d} \frac{e^{-i p (x-y)}}{p^2 + i0}$$

$$= -i \int \frac{d^{d-1} p}{(2\pi)^{d-1} 2|p|} e^{-i |p| |x^0 - y^0| - i p \cdot (x - y)}$$

## Canonical quantization

In Minkowski space,

$$S[C, \bar{C}] = \int d^d x (-i \bar{C} \partial^2 C) = \int d^d x i \partial^\mu \bar{C} \partial_\mu C$$

Reality of fields:  $C^* = C$ ,  $\bar{C}^* = \bar{C}$ .

Let us describe the system in momentum space.

$$C(x) = \int \frac{d^{d-1} P}{(2\pi)^{d-1}} e^{iP \cdot x} C(P), \quad C(P)^* = C(-P)$$

$$\bar{C}(x) = \int \frac{d^{d-1} P}{(2\pi)^{d-1}} e^{iP \cdot x} \bar{C}(P), \quad \bar{C}(P)^* = \bar{C}(-P)$$

$$L = \int \frac{d^{d-1} P}{(2\pi)^{d-1}} \left( i \dot{\bar{C}}(-P) \dot{C}(P) - i P^2 \bar{C}(-P) C(P) \right)$$

$$H = \int \frac{d^{d-1} P}{(2\pi)^{d-1}} \left( i \dot{\bar{C}}(-P) \dot{C}(P) + i P^2 \bar{C}(-P) C(P) \right)$$

A system of this type was discussed in Lecture 3, Exercise (C).

By Ward identity, we find

$$\{C(P), \dot{\bar{C}}(-P')\} = (2\pi)^{d-1} \delta^{d-1}(P-P'),$$

$$\{\bar{C}(-P), \dot{C}(P')\} = -(2\pi)^{d-1} \delta^{d-1}(P-P'),$$

all other anticommutators of  $C, \bar{C}, \dot{C}, \dot{\bar{C}} = 0$ .

Let us introduce

$$b(\mathbf{p}) := \sqrt{\frac{|\mathbf{p}|}{2(2\pi)^{d-1}}} c(\mathbf{p}) + \frac{i}{\sqrt{2(2\pi)^{d-1}|\mathbf{p}|}} \dot{c}(\mathbf{p}),$$

$$\bar{b}(\mathbf{p}) := -i \sqrt{\frac{|\mathbf{p}|}{2(2\pi)^{d-1}}} \bar{c}(\mathbf{p}) + \frac{i}{\sqrt{2(2\pi)^{d-1}|\mathbf{p}|}} \dot{\bar{c}}(\mathbf{p}).$$

Then

$$\{b(\mathbf{p}), \bar{b}(\mathbf{p}')^\dagger\} = \delta^{d-1}(\mathbf{p} - \mathbf{p}')$$

$$\{\bar{b}(\mathbf{p}), b(\mathbf{p}')^\dagger\} = \delta^{d-1}(\mathbf{p} - \mathbf{p}')$$

all other anticommutators of  $b, b^\dagger, \bar{b}, \bar{b}^\dagger = 0$ .

$$H = \int d^{d-1}\mathbf{p} |\mathbf{p}| (b(\mathbf{p})^\dagger \bar{b}(\mathbf{p}) + \bar{b}(\mathbf{p})^\dagger b(\mathbf{p}) - \delta^{d-1}(\mathbf{0}))$$

$$[H, b(\mathbf{p})] = -|\mathbf{p}| b(\mathbf{p}), \quad [H, b(\mathbf{p})^\dagger] = |\mathbf{p}| b(\mathbf{p})^\dagger$$

$$[H, \bar{b}(\mathbf{p})] = -|\mathbf{p}| \bar{b}(\mathbf{p}), \quad [H, \bar{b}(\mathbf{p})^\dagger] = |\mathbf{p}| \bar{b}(\mathbf{p})^\dagger$$

$\therefore b(\mathbf{p})^\dagger, \bar{b}(\mathbf{p})^\dagger$  : creation operators,

$b(\mathbf{p}), \bar{b}(\mathbf{p})$  : annihilation operators.

The state  $|0\rangle$  annihilated by  $b(\mathbf{p})$  &  $\bar{b}(\mathbf{p}) \forall \mathbf{p}$  is

the unique ground state, with energy  $E_0 = -\int d^{d-1}\mathbf{p} |\mathbf{p}| \delta^{d-1}(\mathbf{0})$ .

Other states are obtained from  $|0\rangle$  by operating  $b(p)^\dagger$  &  $\bar{b}(p)^\dagger$ , each increasing energy by  $|p|$ .

Remark There are zero & negative norm states.

$$|p, \phi\rangle := b(p)^\dagger |0\rangle, \quad |\phi, p\rangle := \bar{b}(p)^\dagger |0\rangle$$

$$|p_1, p_2\rangle := b(p_1)^\dagger \bar{b}(p_2)^\dagger |0\rangle$$

$$\begin{aligned} \langle p_1, \phi | p_2, \phi \rangle &= \langle 0 | b(p_1) b(p_2)^\dagger |0\rangle = 0 \\ \langle \phi, p_1 | \phi, p_2 \rangle &= \langle 0 | \bar{b}(p_1) \bar{b}(p_2)^\dagger |0\rangle = 0 \end{aligned} \quad \left. \vphantom{\begin{aligned} \langle p_1, \phi | p_2, \phi \rangle \\ \langle \phi, p_1 | \phi, p_2 \rangle \end{aligned}} \right\} \text{zero norm}$$

$$\begin{aligned} \langle p_1, p_2 | p_3, p_4 \rangle &= \langle 0 | \bar{b}(p_2) b(p_1) b(p_3)^\dagger \bar{b}(p_4)^\dagger |0\rangle \\ &= -\delta^{d-1}(p_1 - p_4) \delta^{d-1}(p_2 - p_3) \quad \leftarrow \text{negative norm} \end{aligned}$$

Interpretation?

$b(p)^\dagger$  &  $\bar{b}(p)^\dagger$  are interpreted as creating massless particles.

However, they create states of zero/negative norm.

Thus these particles are unphysical "ghost" particles.

Computation of  $\langle 0 | T C(x) \bar{C}(y) |0\rangle$

$$C(p) = \sqrt{\frac{(2\pi)^{d-1}}{2|p|}} (b(p) + b(-p)^\dagger),$$

$$\bar{C}(p) = i \sqrt{\frac{(2\pi)^{d-1}}{2|p|}} (\bar{b}(p) + \bar{b}(-p)^\dagger).$$



$$C(x) = \int \frac{d^{d-1}P}{\sqrt{(2\pi)^{d-1} 2|P|}} \left( e^{iP \cdot x} b(P) + e^{-iP \cdot x} b(P)^\dagger \right)$$

$$\bar{C}(x) = \int \frac{d^{d-1}P}{\sqrt{(2\pi)^{d-1} 2|P|}} i \left( e^{iP \cdot x} \bar{b}(P) - e^{-iP \cdot x} \bar{b}(P)^\dagger \right)$$

$$C(t, \mathbf{x}) = e^{itH} C(x) e^{-itH}$$

$$= \int \frac{d^{d-1}P}{\sqrt{(2\pi)^{d-1} 2|P|}} \left( e^{iP \cdot x - i|P|t} b(P) + e^{-iP \cdot x + i|P|t} b(P)^\dagger \right)$$

$$\bar{C}(t, \mathbf{x}) = e^{itH} \bar{C}(x) e^{-itH}$$

$$= \int \frac{d^{d-1}P}{\sqrt{(2\pi)^{d-1} 2|P|}} i \left( e^{iP \cdot x - i|P|t} \bar{b}(P) - e^{-iP \cdot x + i|P|t} \bar{b}(P)^\dagger \right)$$

$$\langle 0 | C(x) \bar{C}(y) | 0 \rangle = -i \int \frac{d^{d-1}P}{(2\pi)^{d-1} 2|P|} e^{-i|P|(x^0 - y^0) + iP \cdot (x - y)}$$

$$\langle 0 | \bar{C}(y) C(x) | 0 \rangle = i \int \frac{d^{d-1}P}{(2\pi)^{d-1} 2|P|} e^{-i|P|(y^0 - x^0) + iP \cdot (y - x)}$$

$$\therefore \langle 0 | T C(x) \bar{C}(y) | 0 \rangle = -i \int \frac{d^{d-1}P}{(2\pi)^{d-1} 2|P|} e^{-i|P||x^0 - y^0| - iP \cdot (x - y)}$$

... match with  $\langle C(x) \bar{C}(y) \rangle$ .

## Total gauge fixed Maxwell theory

$$\tilde{\mathcal{L}} = -\frac{1}{4e^2} F^{\mu\nu} F_{\mu\nu} - \frac{1}{2e^2 \xi} (\partial \cdot A)^2 - i \bar{c} \partial^2 c$$

(Set  $\xi = e = 1$  below)

BRST symmetry

$$\delta_B A_\mu = \partial_\mu c, \quad \delta_B c = 0, \quad \delta_B \bar{c} = i \partial \cdot A$$

$$\leadsto Q_B = \int d^d x (-F^{0\nu} \partial_\nu c - \dot{c} \partial \cdot A)$$

Quantization: (Notation change:  $a_\nu(P) \leftrightarrow a_\nu(-P)^\dagger$ )

$$\leftarrow P_\mu x = (|P|t - \mathbf{P} \cdot \mathbf{x})$$

$$A_\mu(x) = \int \frac{d^{d-1} P}{\sqrt{(2\pi)^{d-1} 2|P|}} \left( e^{-iP_\mu x} a_\mu(P) + e^{iP_\mu x} a_\mu(P)^\dagger \right)$$

$$C(x) = \int \frac{d^{d-1} P}{\sqrt{(2\pi)^{d-1} 2|P|}} \left( e^{-iP_\mu x} b(P) + e^{iP_\mu x} b(P)^\dagger \right)$$

$$\bar{c}(x) = \int \frac{d^{d-1} P}{\sqrt{(2\pi)^{d-1} 2|P|}} i \left( e^{-iP_\mu x} \bar{b}(P) - e^{iP_\mu x} \bar{b}(P)^\dagger \right)$$

$$[a_\mu(P), a_\nu(P')^\dagger] = -\eta_{\mu\nu} \delta^{d-1}(P-P'),$$

$$\{b(P), \bar{b}(P')^\dagger\} = \{\bar{b}(P), b(P')^\dagger\} = \delta^{d-1}(P-P'),$$

Other commutators/anti commutators = 0

$$H = \int d^{d-1}P |P| \left( -\eta^{\mu\nu} a_\mu(P)^\dagger a_\nu(P) + b(P)^\dagger \bar{b}(P) + \bar{b}(P)^\dagger b(P) + \frac{d-2}{2} \delta^{d-1}(0) \right)$$

$$[H, \mathcal{Q}_P] = -|P| \mathcal{Q}_P, \quad [H, \mathcal{Q}_P^\dagger] = |P| \mathcal{Q}_P^\dagger$$

$$\text{for } \mathcal{Q}_P = a_\mu(P), b(P), \bar{b}(P).$$

The state  $|0\rangle$  annihilated by all such  $\mathcal{Q}_P$ 's  $\forall P$  is the unique ground state. Other states are obtained from  $|0\rangle$  by operating  $\mathcal{Q}_P^\dagger$ 's each creating energy  $|P|$ .

$$Q_B = - \int d^{d-1}P \left\{ b(P) \left( \sum_i P_i a_i(P)^\dagger + |P| a_0(P)^\dagger \right) + b(P)^\dagger \left( \sum_i P_i a_i(P) + |P| a_0(P) \right) \right\}$$

$$[Q_B, a_i(P)] = P_i b(P), \quad [Q_B, a_0(P)] = -|P| b(P),$$

$$[Q_B, a_i(P)^\dagger] = -P_i b(P)^\dagger, \quad [Q_B, a_0(P)^\dagger] = |P| b(P)^\dagger,$$

$$\{Q_B, b(P)\} = \{Q_B, b(P)^\dagger\} = 0,$$

$$\{Q_B, \bar{b}(P)\} = - \sum_i P_i a_i(P) - |P| a_0(P),$$

$$\{Q_B, \bar{b}(P)^\dagger\} = - \sum_i P_i a_i(P) - |P| a_0(P)^\dagger.$$

## BRST cohomology

$$\mathcal{H} = \{ \text{products of } a_r^\dagger, b^\dagger, \bar{b}^\dagger \text{ on } |0\rangle \}$$

$$\text{ghost number : } \frac{a_r \quad a_r^\dagger \quad b \quad b^\dagger \quad \bar{b} \quad \bar{b}^\dagger \quad |0\rangle}{0 \quad 0 \quad 1 \quad 1 \quad -1 \quad -1 \quad \underbrace{0}_{\text{definition.}}}$$

$$\mathcal{H}^i = \{ \text{state of ghost number} = i \}$$

$$\dots \xrightarrow{Q_B} \mathcal{H}^i \xrightarrow{Q_B} \mathcal{H}^{i+1} \xrightarrow{Q_B} \mathcal{H}^{i+2} \rightarrow \dots \quad Q_B^2 = 0$$

$\leadsto$  BRST cohomology

$$H^i(\mathcal{H}, Q_B) = \text{Ker}(Q_B: \mathcal{H}^i \rightarrow \mathcal{H}^{i+1}) / \text{Im}(Q_B: \mathcal{H}^{i-1} \rightarrow \mathcal{H}^i).$$

The basic proposal was :

physical states are BRST cohomology classes

$$\mathcal{H}_{\text{phys}} = \bigoplus_i H^i(\mathcal{H}, Q_B).$$

What is  $H^i(\mathcal{H}, Q_B)$  then ?

Let us compute !

Warm-up: Examine low lying states.

$N_c :=$  number of creation operators

$$N_c = 0 \quad \text{on } |0\rangle$$

$N_c = 1$  spanned by  $a_p(p)^\dagger |0\rangle$ ,  $b(p)^\dagger |0\rangle$ ,  $\bar{b}(p)^\dagger |0\rangle$  (all  $p$ 's).

$$\mathcal{H}_{N_c} = \{ N_c \text{ creation ops on } |0\rangle \}$$

$Q_B$  does not change  $N_c$

$$\mathcal{H}_{N_c}^i = \{ N_c \text{ creation ops on } |0\rangle, \text{ ghost number} = i \}$$

$$\xrightarrow{Q_B} \mathcal{H}_{N_c}^i \xrightarrow{Q_B} \mathcal{H}_{N_c}^{i+1} \xrightarrow{Q_B} \dots \quad \text{subcomplex.}$$

$$H^i(\mathcal{H}, Q_B) = \bigoplus_{N_c} H^i(\mathcal{H}_{N_c}, Q_B)$$

$N_c = 0$

$$\begin{array}{ccccccc} & & \mathcal{H}_0^{-1} & & \mathcal{H}_0^0 & & \mathcal{H}_0^1 \\ & & \parallel & & \parallel & & \parallel \\ \dots & 0 & \rightarrow & 0 & \rightarrow & \mathbb{C}|0\rangle & \rightarrow & 0 & \rightarrow & 0 & \dots \end{array}$$

$$\therefore H^i(\mathcal{H}_0, Q_B) = \begin{cases} \mathbb{C}|0\rangle & i=0 \\ 0 & i \neq 0 \end{cases} \quad \leftarrow \text{vacuum}$$

$$\underline{N_c = 1}$$

$$\begin{array}{cccccc} \mathcal{H}_2^{-2} & \mathcal{H}_1^{-1} & \mathcal{H}_1^0 & \mathcal{H}_1^1 & \mathcal{H}_1^2 & \\ \parallel & \parallel & \parallel & \parallel & \parallel & \\ \rightarrow 0 \rightarrow & \{ \bar{b}(p)^\dagger |0\rangle \}_p & \{ a_\mu(p)^\dagger |0\rangle \}_p & \{ b(p)^\dagger |0\rangle \}_p & \rightarrow 0 \rightarrow & \dots \end{array}$$

$$Q_B \bar{b}(p)^\dagger |0\rangle = \left( -\sum_i p_i a_i(p)^\dagger - |p| a_0(p)^\dagger \right) |0\rangle$$

$$\begin{aligned} Q_B \epsilon^\mu a_\mu(p)^\dagger |0\rangle &= -\sum_i \epsilon^i p_i b(p)^\dagger |0\rangle + \epsilon^0 |p| b(p)^\dagger |0\rangle \\ &= \underbrace{\left( -\sum_i \epsilon^i p_i + \epsilon^0 |p| \right)}_{\epsilon \cdot P_p} b(p)^\dagger |0\rangle \end{aligned}$$

$$\epsilon \cdot P_p ; P_p := (|p|, p) \begin{cases} P_p^0 = |p| \\ P_p^i = p_i \end{cases}$$

$$\therefore H^i(\mathcal{H}_1, Q_B) = 0 \quad \text{if } i \neq 0$$

$$H^0(\mathcal{H}_1, Q_B) = \frac{\left\{ \epsilon^\mu(p) a_\mu(p)^\dagger |0\rangle \mid \epsilon(p) \cdot P_p = 0 \right\}_p}{\epsilon(p) \sim \epsilon(p) + \text{const } P_p}$$

$$\cong \left\{ \sum_i \epsilon^i(p) a_i(p)^\dagger |0\rangle \mid \sum_i \epsilon^i(p) p_i = 0 \right\}_p$$

↑  
a particle with a transversal polarization.

(d-2 choices for each p)

Let us put

$$\mathcal{H}_{\text{transv}} := \{ \text{product of transversal } a^\dagger \text{'s on } |0\rangle \}$$

Then

Theorem

$$H^i(\mathcal{H}, \mathcal{Q}_B) \cong \begin{cases} \mathcal{H}_{\text{transv}} & i=0 \\ 0 & i \neq 0. \end{cases}$$

Therefore,  $\mathcal{H}_{\text{phys}} \cong \mathcal{H}_{\text{transv}}$ .

Remarks

- (1) a positive definite inner product is induced on  $\mathcal{H}_{\text{phys}}$ .
- (2) Canonical quantization also gives the same result.