

Course Plan.

★ QFT II / QFT. ^{2024/25} ~~2020/21~~ Autumn Semester.

- §1. Path Integral for Quantum Mechanics. [2 weeks] 10/7, 10/21
- §2. Introduct'n to QFT. [2 weeks] 10/28, 11/11,
- §3. Tree-level Processes & QED. [3 weeks] 11/13 (Wed), 11/18, 25
- §4. Bound States [2 weeks] 12/2, 12/9
- §5. Unitarity [1 week] 12/16
- §6. Low-energy Effective Theories. [1 week] 12/23
- §7. Path Integral for QFT. [3 weeks] 1/6, 20, 23 (Thu)
total
- §8. Introduct'n to 1-loop Calculat'ns. [1 week] 1/27. 15 weeks.

★ Theory of Elementary Particles. ²⁰²⁵ ~~2024~~ Summer Semester. (expected)

UV divergence & Renormalization. [several weeks]

IR divergence & Factorization. [several weeks]

★ Subjects that are not covered by QFT II + TEP.

- quantizati'n of vector fields & gauge symmetry.
- Standard Model of Particle Physics \Rightarrow $\left\{ \begin{array}{l} \text{Elementary Particle Phys (II) Autumn.S} \\ \text{III. Summer.S} \end{array} \right.$
- Anything beyond Perturbation.
- Topology & Anomaly.

§1. Path Integral Formulation in Quantum Mechanics

§1.1 Bosonic Quantum Mechanical Systems

Derivation

Consider a quantum mechanical system

H : Hamiltonian
 (q, p) canonical conjugate pair
 initial state $\psi_0(q_{in}, t_{in})$ @ $t = t_{in}$

Time evolution of the wave function

$$\psi(q, t; \psi_0) = e^{-i \int_{t_{in}}^t H dt} \psi_0(q_{in}, t_{in}) \quad \left(\begin{array}{l} \text{time-ordered} \\ \text{exp. to be more} \\ \text{precise} \end{array} \right)$$

(already $t_1 = t_2$ convention.)

This time evolution can be rewritten as follows:

$$\psi(q, t; \psi_0) = \langle q, t | e^{-iHT} | \psi_0 \rangle = \langle q, t | e^{-iHT} | q_{in}, t_{in} \rangle \psi_0(q_{in}, t_{in})$$

$H(x)$ for simplicity $\int dq_{in}$ $T = (t - t_{in})$

$$\langle q, t | e^{-iHT} | q_{in}, t_{in} \rangle$$

$$= \int dq_{N-1} \dots \int dq_1 \langle q, t | e^{-iH\Delta t} | q_{N-1}, t_{N-1} \rangle \dots$$

$$\langle q, t_2 | e^{-iH\Delta t} | q_1, t_1 \rangle \langle q_1, t_1 | e^{-iH\Delta t} | q_{in}, t_{in} \rangle$$

inserted a complete system of states at a time slice t_0

$$(t = t_N) > t_{N-1} > \dots > t_0 > t_1 > (t_{in} = t_0) \quad \text{interval } \Delta t = \frac{T}{N}$$

In cases with $H = \frac{p^2}{2m} + V(q)$ — (**)

$$\langle q_k, t_k | e^{-iH\Delta t} | q_{k-1}, t_{k-1} \rangle = \int \frac{dp_k}{2\pi} \langle q_k, t_k | p_k, t_k \rangle \langle p_k, t_k | e^{-iH\Delta t} | q_{k-1}, t_{k-1} \rangle$$

use $e^{-iH\Delta t} = e^{-i\frac{p^2}{2m}\Delta t} e^{(0(\Delta t)^2 + p\Delta q)} e^{-iV(q)\Delta t}$

$$= \int \frac{dp_k}{2\pi} e^{i p_k \dot{q}_k - i \frac{p_k^2}{2m} \Delta t - i (0(\Delta t)^2 + p_k \Delta q_k - i V(q_k) \Delta t)}$$

$$= \int \frac{dp_k}{2\pi} e^{i \Delta t \left\{ p_k \left(\frac{\partial q_k - \partial q_{k-1}}{\Delta t} \right) - H(q_{k-1}) \right\} + O((\Delta t)^2)}$$

Therefore

$$\langle q, t | e^{-iHT} | q_i, t_i \rangle = \lim_{N \rightarrow \infty} \int \frac{dp_N}{2\pi} \prod_{k=1}^{N-1} \left(\int \frac{dq_k dp_k}{2\pi} \right) e^{i \Delta t \sum_{k=1}^N \left\{ p_k \left(\frac{\partial q_k - \partial q_{k-1}}{\Delta t} \right) - H(q_{k-1}) \right\}}$$

(drop $(\Delta t)^2$ in the exp. $(N \rightarrow \infty)$)

denote it by $\int_{\substack{\delta = \dot{q}_i @ t = t_i \\ \delta = \dot{q} @ t = t}} \mathcal{D}q(t) \mathcal{D}p(t) e^{i \int dt (p \dot{q} - H)}$

As long as (**) holds: do Gaussian integral

$$\int \frac{dp_k}{2\pi} e^{-i \Delta t \frac{p_k^2}{2m} + i(\Delta t) p_k \dot{q}_k} = \int \frac{dp_k}{2\pi} e^{-i \frac{\Delta t}{2m} (p_k - m \dot{q}_k)^2 + i(\Delta t) \frac{m}{2} (\dot{q}_k)^2}$$

$$= \sqrt{\frac{m}{2\pi i(\Delta t)}} e^{i(\Delta t) \frac{m}{2} (\dot{q}_k)^2}$$

So, there is an alternative expression

$$\langle q_f, t_f | e^{-iHT} | q_i, t_i \rangle \propto \int_{\substack{\delta(t_i) = \delta_i \\ \delta(t_f) = \delta_f}} \mathcal{D}q(t) e^{i \int dt L}$$

(the Gaussian integral factors $\sqrt{\frac{mN}{2\pi i T}}$ is indep. of δ_i or δ_f)

Example 1 free particle $V(q) = 0$

$$\left[\begin{array}{l} q = q_{in} @ t = t_{in} \\ q = q_f @ t = t_{fin} \end{array} \right] \xrightarrow{(*)} \text{take } t_i = 0$$

$$T = (t_{fin} - t_{in})$$

$$\langle q_f, t_f | e^{-iHT} | q_{in}, t_i \rangle = \left(\frac{mN}{2\pi i T} \right)^{\frac{N}{2}} \int_{\text{bdry cond}} \left(\oint q(t) = dq_2 dq_3 \dots dq_{N-1} \right) e^{i \int_0^T dt' \frac{m}{2} (\dot{q})^2}$$

$$q_k = q(t' - (t_{in} - t) + (t - t') \cdot k) = \left(q_{in} + \frac{(\Delta t) \cdot k}{T} (q_f - q_{in}) \right) + \delta q_k$$

$$\Rightarrow q_{cl}(t')$$

} bdry cond $(*)$ satisfied
 } split into (classical solution q_{cl}) and (fluctuation around it)

$$S' = \frac{m}{2} \int_0^T dt' (\dot{q}_{cl} + \dot{\delta q})^2 = \frac{m}{2} \int_0^T dt' \left\{ (\dot{q}_{cl})^2 + (\dot{\delta q})^2 \right\}$$

$$\left(m \int_0^T dt' (\dot{q}_{cl} \dot{\delta q}) = m \int_0^T dt' \frac{d}{dt'} (q_{cl} \delta q) = m [q_{cl} \delta q]_0^T = 0 \right)$$

eg. of motion

so

$$\langle q_f, t_f | e^{-iHT} | q_{in}, t_i \rangle = \left(\frac{mN}{2\pi i T} \right)^{\frac{N}{2}} e^{i \left(\frac{m}{2} \frac{(q_f - q_{in})^2}{T} = S_{cl} \right)} \int dq_2 \dots dq_{N-1} e^{i S_{fl}}$$

$$i S_{fl} = i \frac{m}{2} \sum_{k=2}^N \left(\frac{q_k - q_{k-1}}{\Delta t} \right)^2 (\Delta t) \cong i \frac{mN}{2T} (q_{N-1}, \dots, q_2) \begin{bmatrix} 2 & -1 & & \\ -1 & 2 & & \\ & & \ddots & \\ & & & 2 & -1 \\ & & & -1 & 2 \end{bmatrix} \begin{pmatrix} q_{N-1} \\ \vdots \\ q_2 \end{pmatrix}$$

$$= \left(\frac{mN}{2\pi i T} \right)^{\frac{N}{2}} e^{i \frac{(q_f - q_{in})^2 m}{2T}} \times \left(\frac{2\pi T}{-i m N} \right)^{\frac{N-1}{2}} \frac{1}{\sqrt{\det \begin{bmatrix} 2 & -1 & & \\ -1 & 2 & & \\ & & \ddots & \\ & & & 2 & -1 \\ & & & -1 & 2 \end{bmatrix}}} = \sqrt{\frac{m}{2\pi i T}} e^{i \frac{m}{2} \frac{(q_f - q_{in})^2}{T}}$$

$$\left(\det \begin{bmatrix} 2 & -1 & & \\ -1 & 2 & & \\ & & \ddots & \\ & & & 2 & -1 \\ & & & -1 & 2 \end{bmatrix} = N \right)$$

$\det [2] = 2, \det \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} = 3$ use recursion rel'n

[X.G. Wen's textbook p. 24 ~ 25]

We have reproduced $\int \frac{d^N p}{(2\pi)^N} \langle q_f, T | e^{-i \frac{p^2 T}{2m}} | q_{in}, 0 \rangle = \int \frac{d^N p}{2\pi} e^{i p (q_f - q_{in})} e^{-i \frac{p^2 T}{2m}} = \sqrt{\frac{m}{2\pi i T}} e^{i \frac{m}{2} \frac{(q_f - q_{in})^2}{T}}$ plus

Example 2 harmonic oscillator $H = \frac{1}{2m} p^2 + \frac{1}{2} m \omega^2 q^2$

$$\left. \begin{aligned} q &= q_i \text{ @ } t = t_i = 0 \\ q &= q_f \text{ @ } t = t_f = T \end{aligned} \right\} \text{ bdy condition } \text{---} (*)$$

Expand. (parametrize.)

$$\begin{aligned} q(t) &= q_d(t) + q_{pl}(t) \\ &= A \cos(\omega t + \delta) + \sum_{n=1}^{N-1} X_n \sin\left(\frac{t}{T} \pi n\right) \end{aligned}$$

$$q_i = A \cos(\delta)$$

$$q_f = A \cos(\omega T + \delta) = A [\cos(\omega T) \cos(\delta) - \sin(\omega T) \sin(\delta)]$$

$$\Rightarrow \tan(\delta) = \frac{\cos(\omega T) - (q_f/q_i)}{\sin(\omega T)}$$

$$J^q_d = \int_0^T dt \frac{m}{2} A^2 \omega^2 \{ \sin^2(\omega t + \delta) - \cos^2(\omega t + \delta) \}$$

$$= -\frac{m}{2} A^2 \omega^2 \int_0^T dt \cos(2(\omega t + \delta))$$

$$= \frac{m\omega}{2} \frac{A^2}{2} \{ \sin(2\delta) - \sin(2\omega T + 2\delta) \}$$

$$= \frac{m\omega}{2} \frac{A^2}{2} [\sin(2\delta) \{ 1 - \cos(2\omega T) \} - \cos(2\delta) \sin(2\omega T)]$$

$$= \frac{m\omega}{2} A^2 \sin(\omega T) [\sin(2\delta) \sin(\omega T) - \cos(2\delta) \cos(\omega T)]$$

~~use~~ use $\sin(2\theta) = \frac{2 \tan \theta}{1 + \tan^2 \theta}$ $\cos(2\theta) = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$

$$= \frac{m\omega}{2} \sin(\omega T) (q_i)^2 \left[2 \left(\cos(\omega T) - \frac{q_f}{q_i} \right) - \frac{\{ \sin^2(\omega T) - (\cos(\omega T) - \frac{q_f}{q_i})^2 \} \cos(\omega T)}{\sin^2(\omega T)} \right]$$

$$= \frac{m\omega}{2} \frac{(q_i)^2}{\sin(\omega T)} \left[\left(\frac{q_f}{q_i} \right)^2 \cos(\omega T) + \cos(\omega T) - 2 \frac{q_f}{q_i} \right]$$

$$= \frac{m\omega}{2} \frac{1}{\sin(\omega T)} [\cos(\omega T) (q_i^2 + q_f^2) - 2 q_i q_f]$$

$$S_{fe} = \frac{m}{2} \sum_{n=1}^{N-1} \int_0^T dt' \left\{ \left(\frac{\pi}{T} n \right)^2 \cos^2 \left(\frac{t'}{T} \pi n \right) - \omega^2 \sin^2 \left(\frac{t'}{T} \pi n \right) \right\} (X_n)^2$$

$$= \sum_n \frac{mT}{4} \left\{ \left(\frac{\pi}{T} n \right)^2 - \omega^2 \right\} (X_n)^2$$

Gaussian integral over the fluctuations:

$$\left| \det \left(\frac{\partial^2 k}{\partial X_n^2} \right) \right| \prod_{n=1}^{N-1} \int dx_n e^{i \frac{mT}{4} \left\{ \left(\frac{\pi}{T} n \right)^2 - \omega^2 \right\} (X_n)^2} = \left| \det \left(\frac{\partial^2 k}{\partial X_n^2} \right) \right| \cdot \prod_{n=1}^{N-1} \sqrt{\frac{4\pi}{-imT \left\{ \left(\frac{\pi}{T} n \right)^2 - \omega^2 \right\}}} \quad (***)$$

The ($\omega=0$) case is the free particle case.

$$(***)_{\omega} = (***)_{\omega=0} \times \frac{1}{\prod_{n=1}^{N-1} \left\{ 1 - \left(\frac{\omega T}{\pi n} \right)^2 \right\}} \xrightarrow{\text{lim}_{N \rightarrow \infty}} (***)_{\omega=0} \times \sqrt{\frac{\omega T}{\sin(\omega T)}}$$

So,

$$\langle g_f @ T | e^{-iHT} | g_i @ 0 \rangle = e^{iS_{cl}} \underbrace{\left(\frac{mN}{2\pi i T} \right)^{\frac{N}{2}}}_{\text{free particle}} \times (***)_{\omega=0} \times \sqrt{\frac{\omega T}{\sin(\omega T)}}$$

$$= e^{iS_{cl}} \sqrt{\frac{m}{2\pi i T}} \times \sqrt{\frac{\omega T}{\sin(\omega T)}} \quad (\text{use the result in p.5})$$

$$= e^{iS_{cl}} \sqrt{\frac{m\omega}{2\pi i \sin(\omega T)}}$$

Note: worked out in p.5

Quantum states from path integral

If $\langle q_f \text{ at } t_f | e^{-iH(t_f-t_i)} | q_i \text{ at } t_i \rangle = f_m(q_f, q_i, t_f-t_i)$ is given.

$$f_m(q_f, q_i; t_f-t_i)$$

$$= \int \frac{d\omega}{2\pi} e^{-i\omega(t_f-t_i)} \psi_\omega(q_f) [\psi_\omega(q_i)]^*$$

So, the spectrum and wavefunctions can be extracted from $f_m(q_f, q_i; t_f-t_i)$ by Fourier transformation.

Time-ordered product expectation value

Consider

$$\int \frac{d^N p}{(2\pi)^N} \frac{d^N q}{(2\pi)^N} \dots \frac{d^N q_1}{(2\pi)^N} \frac{d^N p_1}{(2\pi)^N} e^{i(\Delta t) \sum_{k=1}^N \left[p_k \dot{q}_k - H(p_k, q_{k-1}) \right]}$$

$$\times \left(f_{i_1}(q_{i_1}) f_{i_2}(q_{i_2}) \dots \tilde{f}_{j_1}(p_{j_1}) \tilde{f}_{j_2}(p_{j_2}) \dots \right)$$

$$= \langle \mathcal{O}_f \text{ at } t_f \mid T \left\{ \prod_i f_i(q_i) \prod_j \tilde{f}_j(p_j) e^{-i \int_{t_i}^{t_f} H(t')} \right\} \mid \mathcal{O}_i \text{ at } t_i \rangle$$

instead of $\langle \mathcal{O}_f \text{ at } t_f \mid e^{-iHT} \mid \mathcal{O}_i \text{ at } t_i \rangle$

The commutation relation $[q, p] = i$ (equal time)

$$0 = \int \prod_{k=1}^N \left[dq_k \frac{dp_k}{2\pi} \right] \frac{\partial}{\partial p_i} \left(p_i e^{i\Delta t \sum_{k=1}^N \{ p_k \dot{q}_k - H(p_k, q_{k-1}) \}} \right)$$
$$= \int \prod_{k=1}^N \left[dq_k \frac{dp_k}{2\pi} \right] \left(\delta_{ij} + i p_j (\dot{q}_i - \dot{q}_{i-1}) - (\Delta t) \frac{\partial H(p_i, q_{i-1})}{\partial p_i} p_j \right) e^{i\Delta t \sum_{k=1}^N \{ p_k \dot{q}_k - H \}}$$

In the $N \rightarrow \infty$, $\Delta t \rightarrow 0$ limit.

$$\langle \dot{q}_i p_i - p_i \dot{q}_{i-1} \rangle = i \quad (\text{equal time commutator } [q, p] = i)$$

is reproduced from the path integral formulation.