

Nonperturbative Contributions to D3-brane Potentials

Liam McAllister
Cornell

Based on:

D. Baumann, A. Dymarsky, S. Kachru, I. Klebanov, L.M., 1001.5028


Strings 2010, Texas A&M

March 19, 2010




Motivation

- Inflation provides a beautiful causal mechanism to generate the observed CMB anisotropies and distribution of large-scale structure.
- Inflation is sensitive to Planck-scale physics. Planck-suppressed operators generically make critical contributions to the dynamics.



Motivation

- Inflation provides a beautiful causal mechanism to generate the observed CMB anisotropies and distribution of large-scale structure.
- Inflation is sensitive to Planck-scale physics. Planck-suppressed operators generically make critical contributions to the dynamics.
- Therefore, we should understand inflation in string theory, and compute, or at least characterize, the Planck-suppressed contributions.



Physics of slow roll inflation

Scalar field with a potential,


$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} R + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right]$$

Potential drives acceleration,

$$ds^2 = -dt^2 + e^{2Ht} d\vec{x}^2 \quad H \approx \text{const.}$$

- Acceleration prolonged if V is flat in Planck units:

$$\eta \equiv M_p^2 \frac{V''}{V} \quad \epsilon \equiv M_p^2 \left(\frac{V'}{V} \right)^2$$



Planck-sensitivity of inflation

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} R + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V_0(\phi) \right]$$

$$\Delta V \equiv \mathcal{O}_6 = V_0 \frac{\phi^2}{M_p^2} \quad \longrightarrow \quad \begin{array}{l} \Delta\eta \sim 1 \\ N_e \sim 1 \end{array} \quad \eta \equiv M_p^2 \frac{V''}{V}$$

For **small** inflaton excursions, $\Delta\phi \lesssim M_{pl}$, one must control corrections \mathcal{O}_Δ with $\Delta \lesssim 6$.

For **large** inflaton excursions, $\Delta\phi \gg M_{pl}$, one must control an infinite series of corrections, with arbitrarily large Δ .

Options for dealing with the sensitivity to Planck-scale physics.

I. Invoke a symmetry strong enough to forbid all such contributions.

- i.e., forbid the inflaton from coupling to massive d.o.f.

Freese, Frieman, Olinto 1990

Arkani-Hamed, Cheng, Creminelli, Randall 2003

Kalosh, Hsu, Prokushkin 2004

Dimopoulos, Kachru, McGreevy, Wacker 2005

Conlon & Quevedo 2005

L.M., Silverstein, Westphal 2008

Flauger, L.M., Pajer, Westphal, Xu 2008

II. Enumerate all relevant contributions and determine whether fine-tuned inflation can occur.

- i.e., arrange for cancellations.


Baumann, Dymarsky, Klebanov, L.M., 2007

Haack, Kalosh, Krause, Linde, Lüster, Zagermann, 2008

Baumann, Dymarsky, Kachru, Klebanov, L.M., 2008, 2009, 2010


Status report

- Inflaton *candidates* continue to proliferate.
- Many novel phenomenological models inspired by string theory, with distinctive predictions.
- Still very few examples where present techniques admit systematic study of the Planck-suppressed contributions.
- Why is it hard?



Motivation

- Inflationary solutions of string theory are only possible in vacua with stabilized moduli.
 - significant progress in the past decade.
- Integrating out the massive moduli induces interactions that are typically no more than Planck-suppressed.
- Thus, to characterize the inflaton action, we must work in a stabilized vacuum and carefully incorporate the couplings of the inflaton candidate to the moduli.




What's the problem?

$N=0$ supersymmetry.

Compactness crucial.

Effects that stabilize the moduli rarely decouple from inflation.



This talk

- Work in a carefully-chosen corner where powerful tools (noncompact approximation, AdS/CFT) are applicable.
- Goal: characterize the action for a spacetime-filling D3-brane in a type IIB flux compactification, including nonperturbative contributions.
- Nonperturbative effects
 - are crucial in some of the best-studied scenarios for Kähler moduli stabilization
 - make dominant contributions to the inflationary dynamics of a D3-brane in such vacua.
- Our analysis provides a toy model of inflation in quantum gravity.
 - A small step towards a more comprehensive and systematic understanding

Results presented here:

- Structure of the potential for a D3-brane in a conifold attached to a general compact space. All significant contributions to the D3-brane potential incorporated in 10D supergravity.
- Gaugino condensation on D7-branes wrapping a four-cycle sources IASD flux.

I.

D3-branes in flux compactifications

D3-branes in warped compactifications

$$ds^2 = \underline{e^{2A(y)}} g_{\mu\nu} dx^\mu dx^\nu + e^{-2A(y)} \tilde{g}_{mn}(y) dy^m dy^n$$



$$\tilde{F}_5 = (1 + \star_{10}) d\underline{\alpha(y)} \wedge \sqrt{-\det g_{\mu\nu}} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3$$

DBI+CS:

$$V = T_3 \Phi_-$$

$$\Phi_\pm \equiv e^{4A} \pm \alpha$$

D3-brane



\tilde{g}_{mn} : CY at leading order

cf. talk by Lü

D3-branes in warped flux compactifications

$$ds^2 = e^{2A(y)} g_{\mu\nu} dx^\mu dx^\nu + e^{-2A(y)} \tilde{g}_{mn}(y) dy^m dy^n$$

$$\tilde{F}_5 = (1 + \star_{10}) d\alpha(y) \wedge \sqrt{-\det g_{\mu\nu}} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3$$

$$G_\pm \equiv (i \pm \star_6) G_3 \quad \Phi_\pm \equiv e^{4A} \pm \alpha \quad V = T_3 \Phi_-$$

$$\text{ISD solutions: } G_- = \Phi_- = 0$$

GKP 2001

D3-branes in warped flux compactifications

$$ds^2 = e^{2A(y)} g_{\mu\nu} dx^\mu dx^\nu + e^{-2A(y)} \tilde{g}_{mn}(y) dy^m dy^n$$

$$\tilde{F}_5 = (1 + \star_{10}) d\alpha(y) \wedge \sqrt{-\det g_{\mu\nu}} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3$$

$$G_\pm \equiv (i \pm \star_6) G_3 \quad \Phi_\pm \equiv e^{4A} \pm \alpha \quad V = T_3 \Phi_-$$

$$\text{ISD solutions: } G_- = \Phi_- = 0$$

GKP 2001

D3-branes feel no potential in ISD solutions ('no-scale'), but nonperturbative stabilization of Kähler moduli spoils this.


Nonperturbative contributions

No-scale symmetry $\rho \rightarrow \rho + \text{const.}$ is broken by Euclidean D3-branes, or gaugino condensation on N_c D7-branes, wrapping suitable four-cycles Σ_i .

Witten, 1996 KKL, 2003

$$W = \int G_3 \wedge \Omega + A(y) e^{-a\rho}, \quad \mathcal{K} = -3 \log(\rho + \bar{\rho} - k(y, \bar{y}))$$


$$a = \frac{2\pi}{N_c}$$



Euclidean D3-brane,
or stack of $N > 1$ D7-branes

cf. talk by Cvetič

Nonperturbative contributions



If divisor is defined by $h(z_\alpha) = 0$ in local coordinates, then

$$W_{\text{np}}(z_\alpha) = \mathcal{A}_0 h(z_\alpha)^{1/N_c} e^{-a\rho}$$

Ganor, 1996

Berg, Haack, Körs, 2004

Baumann, Dymarsky, Klebanov, Maldacena, L.M., Murugan, 2006

Koerber & Martucci, 2007

$$W = \int G_3 \wedge \Omega + W_{\text{np}}(z_\alpha)$$

D3-brane vacua are generically isolated.
We want the potential in between.

DeWolfe, L.M.,
Shiu, Underwood, 2007

D3-branes in warped flux compactifications

$$ds^2 = e^{2A(y)} g_{\mu\nu} dx^\mu dx^\nu + e^{-2A(y)} \tilde{g}_{mn}(y) dy^m dy^n$$

$$\tilde{F}_5 = (1 + \star_{10}) d\alpha(y) \wedge \sqrt{-\det g_{\mu\nu}} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3$$

$$G_\pm \equiv (i \pm \star_6) G_3 \quad \Phi_\pm \equiv e^{4A} \pm \alpha \quad V = T_3 \Phi_-$$

$$\text{ISD solutions: } G_- = \Phi_- = 0$$

GKP 2001

D3-branes feel no potential in ISD solutions ('no-scale'), but nonperturbative stabilization of Kähler moduli spoils this.

We will **expand around ISD solutions**, $\Phi_-^{(0)} = G_-^{(0)} = 0$

and find that nonperturbative effects source IASD flux.

Equations of motion expanded around an ISD background

$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2 + \mathcal{R}_4$$

$$d\Lambda = 0 \quad \star_6 \Lambda = -i\Lambda \quad \Lambda \equiv \Phi_+ G_-$$

Metric \tilde{g}_{mn} and dilaton:

zeroth-order solutions suffice to determine leading contributions to D3-brane potential, Φ_- .

We only need to know the **background metric**.

Similarly, Bianchi identity for G must be solved, but is not relevant for determining leading contributions.

General compact model still intractable


$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2 + \mathcal{R}_4$$

$$d\Lambda = 0 \quad \star_6 \Lambda = -i\Lambda$$

We do need the internal background (zeroth-order) metric.


General structure of the D3-brane potential?

Hard to compute in full generality.



General structure of the D3-brane potential?


Hard to compute in full generality.



Idea: begin with a noncompact CY cone, and systematically incorporate compactification effects.

General structure of the D3-brane potential?


Hard to compute in full generality.



Idea: begin with a noncompact CY cone, and systematically incorporate compactification effects.

Compactification effects

e.g.:



Warped throats in compact spaces

We will obtain a further handle on the problem by taking the cone to be **warped**, such as the warped conifold or a more general warped CY cone.

Concrete example: a finite-length KS throat, which we approximate by $\text{AdS}_5 \times T^{1,1}$.


$$ds^2 = e^{2A_{(0)}(r)} \eta_{\mu\nu} dx^\mu dx^\nu + e^{-2A_{(0)}(r)} (dr^2 + r^2 d\Omega_{T^{1,1}}^2)$$

$$e^{-4A_{(0)}(r)} = \frac{L^4}{r^4} \quad L^4 \equiv \frac{27\pi}{4} g_s N (\alpha')^2$$


A warped CY cone attached to a stabilized compactification is precisely the configuration of interest in warped D-brane inflation.

Warped D-brane inflation

warped throat
(e.g. Klebanov-Strassler)



CY orientifold, with
fluxes and nonperturbative W
(KKLT 2003)



Dvali&Tye 1998

Dvali,Shafi,Solganik 2001

Burgess,Majumdar,Nolte,Quevedo,Rajesh,Zhang 2001

Kachru, Kallosh, Linde, Maldacena, L.M., Trivedi, 2003

Filtering in the throat

The warped geometry **filters** the compactification effects: the dominant effects in the IR are those with the smallest dimensions Δ_i .

$$V = \sum_i c_i \phi^{\Delta_i} h_i(\Psi)$$

By determining the spectrum of Δ_i we can extract the **leading terms** in the potential.

Double expansion:

around ISD backgrounds $\Phi_-, G_- \ll 1$

and in distance from the UV $r_{D3} \ll r_{UV}$
(hierarchy of scales)

Noncompact approximation

The D3-brane potential comes from Φ_- alone. At leading order in an expansion around ISD backgrounds, the only relevant 10d source for Φ_- is IASD flux Λ . $\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2$

Arbitrary **compactification effects** can be represented by specifying boundary conditions for Φ_- and Λ in the UV of the throat, i.e. by allowing arbitrary **non-normalizable profiles**.

Noncompact approximation

The D3-brane potential comes from Φ_- alone. At leading order in an expansion around ISD backgrounds, the only relevant 10d source for Φ_- is IASD flux Λ .
$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2$$


Arbitrary **compactification effects** can be represented by specifying boundary conditions for Φ_- and Λ in the UV of the throat, i.e. by allowing arbitrary **non-normalizable profiles**.



Fluxes are the leading source

$$d\Lambda = 0 \quad \star_6 \Lambda = -i\Lambda$$

$$\Lambda \equiv \Phi_+ G_-$$



$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2 + \mathcal{R}_4$$

To solve for the potential, we must first solve for the IASD flux.



Three classes of IASD flux solutions in Calabi-Yau cones

$$\boxed{d\Lambda = 0}$$

$$\boxed{\star_6 \Lambda = -i\Lambda}$$

$$\Lambda_{\text{III}} = f_3 \Omega \quad (3,0)$$

Three classes of IASD flux solutions in Calabi-Yau cones

$$\boxed{d\Lambda = 0}$$

$$\boxed{\star_6 \Lambda = -i\Lambda}$$

$$\Lambda_{\text{II}} = \partial f_2 \wedge J$$

$$(2,1)_{\text{NP}}$$

$$\Lambda_{\text{III}} = f_3 \Omega$$

$$(3,0)$$

Three classes of IASD flux solutions in Calabi-Yau cones

$$\boxed{d\Lambda = 0}$$

$$\boxed{\star_6 \Lambda = -i\Lambda}$$

$$(\Lambda_{\text{I}})_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_{\alpha}\nabla_{\sigma}f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}} \quad (1,2)$$

$$\Lambda_{\text{II}} = \partial f_2 \wedge J \quad (2,1)_{\text{NP}}$$

$$\Lambda_{\text{III}} = f_3 \Omega \quad (3,0)$$

Here f_i are holomorphic functions. Easy to generalize to harmonic functions.

D3-brane potential from holomorphic fluxes

$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2 \quad \left| \quad \begin{aligned} \Lambda_{\text{I}} &= \nabla \nabla f_1 \cdot \bar{\Omega} \\ \Lambda_{\text{II}} &= \partial f_2 \wedge J \\ \Lambda_{\text{III}} &= f_3 \Omega \end{aligned} \right.$$

$$\Phi_- = \frac{g_s}{32} \left[g^{\alpha\bar{\beta}} \nabla_{\alpha} f_1 \overline{\nabla_{\beta} f_1} + 2|f_2|^2 + 2 \nabla^{-2} |f_3|^2 \right]$$

Simple when fluxes are holomorphic.



Harmonics on $\text{AdS}_5 \times \text{T}^{1,1}$

$$\nabla^2 f = 0 \quad f = r^{\Delta[j_1, j_2, R]} Y_{[j_1, j_2, R]}(\Psi)$$

$$\begin{aligned} \Delta_f &\equiv -2 + \sqrt{H(j_1, j_2, R_f) + 4} & H(j_1, j_2, R_f) &\equiv 6 \left[j_1(j_1 + 1) + j_2(j_2 + 1) - R_f^2/8 \right] \\ &= \frac{3}{2}, 2, 3, \sqrt{28} - 2, \dots \end{aligned} \quad \text{Ceresole, Dall'Agata, D'Auria, Ferrara 1999}$$

$$(\Lambda_{\text{I}})_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_{\alpha} \nabla_{\sigma} f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}} \quad \Lambda_{\text{I}} \sim r^4 G_{-} \sim r^{\delta_{\text{I}}}$$

$$\delta_{\text{I}} = 1 + \Delta_f = -1 + \sqrt{H(j_1, j_2, R + 2) + 4}$$

D3-brane potential from non-holomorphic fluxes

$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2$$

$$\Phi_- = \sum_{\delta_i, \delta_j} r^{\Delta(\delta_i, \delta_j)} h_{(\delta_i, \delta_j)}(\Psi) \quad \Delta \equiv \delta_i + \delta_j - 4$$

$$\delta_I = 1 + \Delta_f = -1 + \sqrt{H(j_1, j_2, R + 2) + 4}$$

Spectrum of the D3-brane potential

$$\Delta_{\mathcal{H}} = \frac{3}{2}, 2, 3, \dots$$

$$\Delta_{\Lambda} = 1, 2, \frac{5}{2}, \sqrt{28} - \frac{5}{2}$$


$$\Delta_{\mathcal{R}} = 2_s, 3, \frac{7}{2}, 4, \dots$$

$$\nabla^2 \Phi_- = \frac{g_s}{96} |\Lambda|^2 + \mathcal{R}_4$$

$$V = \sum_i c_i \phi^{\Delta_i} h_i(\Psi)$$

$$V(\phi) = V_0 + b_1 j_1(\Psi) \phi^1 + a_{3/2} h_{3/2}(\Psi) \phi^{3/2} + \left(c_2 + a_2 h_2(\Psi) + b_2 j_2(\Psi) \right) \phi^2 + b_{5/2} j_{5/2}(\Psi) \phi^{5/2} + b_{2.79} j_{2.79}(\Psi) \phi^{2.79} + \dots$$

Baumann, Dymarsky, Kachru, Klebanov, L.M., 1001.5028




General structure attested in examples

Special case is explicitly computable: assume the moduli-stabilizing D7-branes hang into the throat region

Can then compute superpotential.

Baumann, Dymarsky, Klebanov, Maldacena, L.M., & Murugan, 2006.



Resulting potential:

$$V(\phi) = V_0 + b_1 j_1(\Psi) \phi^1 + a_{3/2} h_{3/2}(\Psi) \phi^{3/2} + \left(c_2 + a_2 h_2(\Psi) + b_2 j_2(\Psi) \right) \phi^2 + \dots$$

Identical structure!

II.

Gauge theory description

Gauge theory version

Arbitrary compactification effects can be represented by incorporating arbitrary perturbations of the CFT [Lagrangian](#), including coupling it to 4D gravity and to hidden sector degrees of freedom.

$$\mathcal{L}_0 + \delta\mathcal{L} = \int d^2\theta d^2\bar{\theta} (K_0 + \delta K) + \int d^2\theta (W_0 + \delta W) + h.c.$$

We must first classify all operators in the CFT that are dual to the IASD flux modes of interest.

Gauge theory version

Some IASD flux perturbations are dual to supersymmetric perturbations of the CFT Lagrangian,

$$\int d^2\theta \mathcal{O}_{CFT}$$

while others have supersymmetry broken by a hidden sector spurion,

$$\int d^2\theta \mathcal{O}_{CFT} X \quad X = \theta^2 F_X$$

Analogous formulae for non-chiral contributions.

Gauge theory version

Klebanov-Witten SCFT:

SU(N) x SU(N) gauge group $W_{\alpha}^{\pm} \equiv W_{\alpha}^{(1)} \pm W_{\alpha}^{(2)}$

SU(2) x SU(2) x U(1)_R global symmetry $[j_1, j_2, R]$

bifundamentals A_i, B_i

Flux	Operator	Δ	R	j_1	j_2
$\nabla\nabla f_1 \cdot \bar{\Omega}$	$[\text{Tr}(AB)^k]_{\theta^2}$	$\frac{3}{2}k + 1$	$k - 2$	$\frac{1}{2}k$	$\frac{1}{2}k$
$\partial f_2 \wedge J$	$[\text{Tr}[W_+^{\alpha}(AB)^k]]_{\theta}$	$\frac{3}{2}k + 2$	k	$\frac{1}{2}k$	$\frac{1}{2}k$
$f_3 \Omega$	$[\text{Tr}[(W_+^2(AB)^k)]_b$	$\frac{3}{2}k + 3$	$k + 2$	$\frac{1}{2}k$	$\frac{1}{2}k$

Table 7: Matching between supergravity G_- flux modes and CFT operators.

δ	j_1	j_2	R	Operator		Multiplet	Type	Flux Series
$\frac{5}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	-1	$[S^1]_{\theta^2}$	$[\text{Tr}(AB)]_{\theta^2}$	V.I	chiral	I
3	0	0	2	$[\Phi_+^0]_b$	$[\text{Tr}(W_{(1)}^2 + W_{(2)}^2)]_b$	V.IV	chiral	III
$\frac{7}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	$[T_\alpha^1]_\theta$	$[\text{Tr}(W_\alpha(AB))]_\theta$	G.I	chiral	II
4	0	0	0	$[\Phi_-^0]_{\theta^2}$	$[\text{Tr}(W_{(1)}^2 - W_{(2)}^2)]_{\theta^2}$	V.III	chiral	★
4	0	1	0	$[{}_a L_\alpha^{2,0}]_\theta$	$[\text{Tr}(W_\alpha J_a)]_\theta$	G.I+G.III	semi-long	II
4	1	0	0	$[{}_b L_\alpha^{2,0}]_\theta$	$[\text{Tr}(W_\alpha J_b)]_\theta$	G.I+G.III	semi-long	II
4	1	1	0	$[S^2]_{\theta^2}$	$[\text{Tr}(AB)^2]_{\theta^2}$	V.I	chiral	I
$\sqrt{28} - 1$	1	1	-2	–	$[\text{Tr}(f)]_{\theta^2}$	V.I	long	I
$\frac{9}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	3	$[\Phi_+^1]_b$	$[\text{Tr}(W_{(1)}^2 + W_{(2)}^2)(AB)]_b$	V.IV	chiral	III
$\frac{9}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	$[\bar{\Phi}_+^1]_b$	$[\text{Tr}(W_{(1)}^2 + W_{(2)}^2)(\overline{AB})]_b$	V.IV	–	III
$\frac{9}{2}$	$\frac{1}{2}$	$\frac{3}{2}$	-1	$[{}_a J^1]_{\theta^2}$	$[\text{Tr}(J_a(AB))]_{\theta^2}$	V.I	semi-long	I
$\frac{9}{2}$	$\frac{3}{2}$	$\frac{1}{2}$	-1	$[{}_b J^1]_{\theta^2}$	$[\text{Tr}(J_b(AB))]_{\theta^2}$	V.I	semi-long	I
5	1	1	2	$[T_\alpha^2]_\theta$	$[\text{Tr}(W_\alpha(AB)^2)]_\theta$	G.I	chiral	II
5	0	1	2	$[{}_a I^0]_b$	$[\text{Tr}((W_{(1)}^2 + W_{(2)}^2)J_a)]_b$	V.IV	semi-long	III
5	1	0	2	$[{}_b I^0]_b$	$[\text{Tr}((W_{(1)}^2 + W_{(2)}^2)J_b)]_b$	V.IV	semi-long	III
$\sqrt{28}$	1	1	0	–	$[\text{Tr}(W_\alpha f)]_\theta$	G.I+G.III	long	II
$\sqrt{40} - 1$	0	2	-2	–	$[\text{Tr}(f_a)]_{\theta^2}$	V.I	long	I
$\sqrt{40} - 1$	2	0	-2	–	$[\text{Tr}(f_b)]_{\theta^2}$	cf. Ceresole, Dall'Agata, D'Auria 1999		

Gauge theory version

The **leading (r^1) term** comes from a superpotential perturbation by the **lowest-dimension gauge invariant operator** in the Klebanov-Witten SCFT,

$$\int d^2\theta \operatorname{Tr}(AB)$$

exactly as one would expect.

(relation to G-flux: cf. [Graña & Polchinski 2000.](#))

For general perturbations by **chiral** operators, we reproduce the gravity-side potential.


We study perturbations by **non-chiral** operators exclusively in the dual gravity description, as modes of IASD flux (with Δ irrational in general).

III.


**IASD fluxes sourced by
nonperturbative effects**

Four-dimensional effects in ten dimensions?


cf. Frey and Lippert 2005, Koerber and Martucci 2007



Four-dimensional effects in ten dimensions?




Four-dimensional effects in ten dimensions?



Four-dimensional effects in ten dimensions?

- Gaugino condensate is a 4d IR effect, but:
- In a suitably anisotropic compactification, can have 10d description valid locally (near D3-brane) even as D7-brane theory undergoes gaugino condensation.
- Equivalently, non-locality due to 4d nonperturbative effects is localized near the corresponding four-cycle.




Noncompact example

$$W_{\text{np}}(z_\alpha) = \mathcal{A}_0 h(z_\alpha)^{1/N_c} e^{-a\rho}$$

$$V = g^{\alpha\bar{\beta}} \nabla_\alpha W \overline{\nabla_\beta W}$$

N_c D7-branes

$$\Sigma : h(z_\alpha) = 0$$




Noncompact example

$$W_{\text{np}}(z_\alpha) = \mathcal{A}_0 h(z_\alpha)^{1/N_c} e^{-a\rho}$$

$$V = g^{\alpha\bar{\beta}} \nabla_\alpha W \overline{\nabla_\beta W}$$

N_c D7-branes

$$\Sigma : h(z_\alpha) = 0$$



Gaugino condensation sources IASD flux

First, work out complete 10d EOM for fluxes.
The D7-brane gaugino mass comes from the coupling

$$\mathcal{L} = 16 c_0 \zeta \int_{\Sigma} \sqrt{g} G_3 \cdot \Omega \bar{\lambda} \lambda + c.c. \quad \zeta \equiv T_3 \sqrt{\frac{g_s}{32}}$$

Cámara, Ibáñez, Uranga 2004

$$m_{\lambda} \propto G_{0,3} \Leftrightarrow F_{\rho}$$

Gaugino condensation sources IASD flux

First, work out complete 10d EOM for fluxes.
The D7-brane gaugino mass comes from the coupling

$$\mathcal{L} = 16 c_0 \zeta \int_{\Sigma} \sqrt{g} G_3 \cdot \Omega \bar{\lambda} \lambda + c.c. \quad \zeta \equiv T_3 \sqrt{\frac{g_s}{32}}$$

Cámara, Ibáñez, Uranga 2004

This yields a **local source term** in the 10d EOM,

$$d\Lambda = d\left(\frac{2\pi}{\zeta} \bar{\Omega} \lambda \lambda \delta^{(0)}\right) \quad \delta^{(0)} = \frac{1}{2\pi} \nabla^2 \text{Re}(\log h)$$
$$\star_6 \Lambda = -i\Lambda$$

Heretofore omitted because $\langle \text{Tr} \lambda \lambda \rangle = 0$ in classical solutions. But we must include this source term!

Gaugino condensation sources IASD flux

To solve

$$d\Lambda = d\left(\frac{2\pi}{\zeta} \bar{\Omega} \lambda\lambda \delta^{(0)}\right) \quad \delta^{(0)} = \frac{1}{2\pi} \nabla^2 \text{Re}(\log h)$$
$$\star_6 \Lambda = -i\Lambda$$

Gaugino condensation sources IASD flux

To solve

$$d\Lambda = d\left(\frac{2\pi}{\zeta} \bar{\Omega} \lambda\lambda \delta^{(0)}\right) \quad \delta^{(0)} = \frac{1}{2\pi} \nabla^2 \text{Re}(\log h)$$
$$\star_6 \Lambda = -i\Lambda$$

we note that for $(\Lambda_I)_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_\alpha \nabla_\sigma f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}}$,

$$d\Lambda = d\left(\frac{1}{2} \nabla^2 f_1 \wedge \bar{\Omega}\right)$$

Gaugino condensation sources IASD flux

To solve

$$d\Lambda = d\left(\frac{2\pi}{\zeta} \bar{\Omega} \lambda\lambda \delta^{(0)}\right) \quad \delta^{(0)} = \frac{1}{2\pi} \nabla^2 \text{Re}(\log h)$$
$$\star_6 \Lambda = -i\Lambda$$

we note that for $(\Lambda_I)_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_\alpha \nabla_\sigma f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}}$,

$$d\Lambda = d\left(\frac{1}{2} \nabla^2 f_1 \wedge \bar{\Omega}\right)$$

so the solution is $f_1 = 2\zeta^{-1} \lambda\lambda \text{Re}(\log h(z_\alpha))$

Gaugino condensation sources IASD flux

- With (1,2) flux given by this solution,

$$(\Lambda_I)_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_\alpha \nabla_{\bar{\sigma}} f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}} \quad f_1 = 2\zeta^{-1} \lambda \lambda \operatorname{Re}(\log h(z_\alpha))$$

Gaugino condensation sources IASD flux

- With (1,2) flux given by this solution,

$$(\Lambda_I)_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_\alpha \nabla_{\bar{\sigma}} f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}} \quad f_1 = 2\zeta^{-1} \lambda \lambda \operatorname{Re}(\log h(z_\alpha))$$

we use
$$\Phi_- = \frac{g_s}{32} \left[g^{\alpha\bar{\beta}} \nabla_\alpha f_1 \overline{\nabla_{\bar{\beta}} f_1} \right]$$

to find a **10d DBI+CS potential**

Gaugino condensation sources IASD flux

- With (1,2) flux given by this solution,

$$(\Lambda_I)_{\alpha\bar{\beta}\bar{\gamma}} = \nabla_\alpha \nabla_\sigma f_1 g^{\sigma\bar{\zeta}} \bar{\Omega}_{\bar{\zeta}\bar{\beta}\bar{\gamma}} \quad f_1 = 2\zeta^{-1} \lambda \lambda \operatorname{Re}(\log h(z_\alpha))$$

we use
$$\Phi_- = \frac{g_s}{32} \left[g^{\alpha\bar{\beta}} \nabla_\alpha f_1 \overline{\nabla_\beta f_1} \right]$$

to find a **10d DBI+CS potential**

that precisely coincides with

the 4d F-term potential

computed with $W = \mathcal{A}_0 h(z_\alpha)^{1/N_c} e^{-a\rho}$

The sourced IASD flux ‘geometrizes’ the gaugino condensate superpotential.

Conclusions

- Obtained structure of the potential for a D3-brane in a conifold attached to a general compact space.
- All significant contributions to the D3-brane potential captured in 10d supergravity.
- Results consistent with computation in the dual gauge theory and in 4d supergravity.
- Gaugino condensation on D7-branes sources IASD flux.
 - The flux ‘geometrizes’ the nonperturbative superpotential.